



Bulk Density of Soils from Oil Palm Agroforestry Systems in Kogi East, Nigeria

**S. O. Amhakhian^a, I. J. J. Otene^{a*}, I. O. Adava^a, B. Muhammed^a, E. C. Are^a
and N. O. Ozovehe^a**

^a *Department of Soil and Environmental Management, Faculty of Agriculture, Kogi State University, P.M.B. 1008, Anyigba, Kogi State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2021/v11i1230590

Editor(s):

(1) Dr. Wen-Cheng Liu, National United University, Taiwan.

Reviewers:

(1) Abdelaali RAHMOUNI, Sidi Mohamed Ben Abdellah University, Morocco.

(2) M. Nuri ÖNER, Çankırı Karatekin University, Turkey.

(3) Avtar Singh Bimbraw, Punjab Agricultural University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/78103>

Original Research Article

Received 15 October 2021

Accepted 10 December 2021

Published 20 December 2021

ABSTRACT

This study was conducted to assess the bulk density and textural class of soils of selected oil palm agroforestry systems in Kogi East (Ankpa, Dekina, Ofu, Olamaboro and Omala local government areas- LGAs), Nigeria. Random soil sampling was used to collect a total of 100 core samples from the five LGAs (20 from each of the LGA). Soil bulk density was determined using core sampler technique. Data collected were analysed using Analysis of Variance (ANOVA) of GENSTAT Discovery Software. Significant means were separated using Duncan Multiple Range Test at 5 % level of probability. The results revealed that the soil bulk density of the oil palm agroforestry systems in Kogi East, Nigeria were within moderate range of 1.15-1.47 g/cm³. Most of the soils in the studied locations were observed to be sandy clay: Ofu and Olamaboro LGAs were sandy clay while Dekina, Ankpa, Omala were loamy sand, sand, and sandy loam respectively.

Keywords: Agroforestry; soil physical properties; soil management; tillage.

1. INTRODUCTION

Bulk density of soils within thresholds can contribute to the capacity of the soils to function in terms of soil aeration, structural support, water and solute movement. This implies that bulk density of soils that are above thresholds could lead to impaired soil function. Based on soil texture, ideal bulk density for plant growth are < 1.60 , < 1.40 , and $< 1.10 \text{ g/cm}^3$ for sandy, silty, and clayey soils respectively while bulk density that restricts plant growth are >1.80 , >1.65 , and >1.47 for sandy, silty, and clayey soils respectively [1, 2,3]. High values of soil bulk density can result to soil compaction [4]. Soil compaction can limit biochemical processes and microbial activities, restrict root growth, negatively affect the movement of air and water through the soil profile, decrease plant growth and crop yield including reduction of vegetative cover available to protect soil from erosion [5], Keesstra et al., 2016; [4].

Agroforestry is a land use system that combines agriculture (crops and/or livestock) and forestry (trees and shrubs) on the same land management unit [6]. Agroforestry as an integrated and sustainable land use system can provide food and income for the farmer as well as can contribute to environmental sustainability. Agroforestry systems are source of nutrition as well as additional income for farmers engaged in it - the farmers practicing agroforestry are gainfully employed with reduced level of poverty and enhanced livelihoods. The sales of food crops as well as timber and non-timber products by the farmers in AFS can promote sustainable development [7,8,6]. Furthermore, the diversification of farm outputs in an agroforestry system is helpful in the reduction of risks from total crop failure compared to monoculture system in periods of extreme weather events including floods and droughts [9]. The environmental benefits of agroforestry systems (AFS) include water quality management - the roots of trees in AFS can reduce nitrogen and phosphorus residues in soils and also the reduction in the use of inputs such as fertilizers and pesticides in AFS can reduce water contamination and eutrophication [10]; conservation of biodiversity by providing habitat for biodiversity to live and breed [11]; soil quality improvement through carbon sequestration, enhancement of soil nutrient cycling, soil structure, soil moisture retention [12]. Oil palm AFS have been reported to improve soil physical properties such as bulk density [13]. On the

other hand, the findings of Khasanah et al. [14] and Rahmani et al. [15] revealed that oil palm Agroforestry AFS can provide economic and environmental benefits. This study was conducted to assess the soil bulk density and textural class of soils from selected oil palm agroforestry systems in Kogi East. Knowledge from this study will provide insights on the ability of the soils of the selected oil palm agroforestry systems in Kogi East Nigeria to provide structural support, soil aeration, water and solute movement.

2. METHODOLOGY

2.1 Study Location

This study was conducted in selected oil palm agroforestry systems in Kogi East (5 local government areas: Ankpa, Dekina, Ofu, Olamaboro and Omala). Study locations were comprised of 4 communities in each LGA. The communities within Ankpa were Odagbo, Oje Elanyi, Ojogobi Olaji, and Okabo. For Dekina LGA, the following communities were selected: Anyigba, Egume, Dekina, and Odu Ogbaloto. The communities selected for Ofu LGA were Ejule, Ochadamu, Ogbulu, and Ugwolawo. Olamaboro was comprised of Ejoka, Igoti-Ade, Ubalu, and Unobe while communities selected from Omala LGA were Ajedibo, Ajomakoji, Odumukpo, and Okugba.

2.2 Soil Sampling and Analysis

Random soil sampling technique was employed for the collection of samples. For bulk density, core samplers of known weight, height and diameter were used to collect a total of 100 core samples from the five LGAs (20 from each of the LGA). For particle size analysis, soil auger was used to collect surface samples (0-15 cm depth). Also, a total of 100 samples (20 from each of the LGA) were collected for particle size analysis but were bulked to 60 samples (12 composite samples per LGA). The prepared soil samples were analyzed at the Department of Soil and Environment Management, Faculty of Agriculture, Kogi State University Anyigba for bulk density and particle size distribution.

2.2.1 Determination of bulk density

The soil bulk density was determined using core sampler technique. Where samples contained in the core rings of known weight, height and diameter were weighed and the fresh weight

Ankpa Local Government Area: The values of bulk density of soils from oil palm agroforestry systems in Ankpa were within 1.35-1.42 g/cm³. Odagbo, Oje-Elanyi, Ojogobi Olaji, and Okabo had bulk density of 1.36, 1.42, 1.35, and 1.39 g/cm³ respectively.

Dekina Local Government Area: Bulk density of Dekina soils were within 1.27-1.40 g/cm³. Anyigba, Dekina, Egume, and Odu Ogbaloto had bulk density of 1.36, 1.40, 1.27, and 1.40 g/cm³ respectively.

Omala Local Government Area: The oil palm agroforestry systems in Omala LGA had bulk density within 1.41-1.47 g/cm³. Ajedibo, Ajomakoji, Odumukpo, and Okugba had bulk density of 1.44, 1.47, 1.41, and 1.45 g/cm³ respectively.

Ofu Local Government Area: The bulk density recorded from Ogbulu, Ugwolawo, and Ejule were 1.43 g/cm³, 1.37 g/cm³, and 1.32 g/cm³ respectively in Ofu local government area. There was no significant difference between the soils of the three locations. Although Ochadamu had the lowest Bd (1.23 g/cm³), there was no significant difference between the value and that of Ejule soils (1.32 g/cm³). The oil palm agroforestry

systems in Ofu LGA had bulk density within 1.23-1.43 g/cm³.

Olamaboro Local Government Area: Bulk density results obtained from soils of Ejoka, Igoti Ade, Unobe, and Ubalu in Olamaboro local government area were 1.38 g/cm³, 1.28 g/cm³, 1.16 g/cm³, and 1.15 g/cm³ respectively. There was no significant difference in the bulk density of Ejoka and Igoti Ade. Lowest and no significant difference in bulk densities were observed in soils in Ubalu and Unobe. Bulk densities of Olamaboro soils were within 1.15-1.38 g/cm³.

3.2 Discussion

3.2.1 Texture of soils of oil palm agroforestry systems in Kogi East, Nigeria

Texture is an important soil property as it greatly affects crop production and soil quality – it plays a key role on soil nutrient and water retention including biological processes in the soil [19,20,21]. Most of the soils in the studied locations were observed to be sandy clay. The chemical and physical properties of sand soils are low [22]. Sandy soils have low organic matter content (OMC), cation exchange capacity (CEC) and therefore low nutrient retention capacity [23].

Table 3. Bulk density of soils from oil palm agroforestry systems in Kogi East, Nigeria

Local Government Area	Location of Oil Palm Agroforestry System	Bulk Density (g/cm ³)	Statistics			
			LSD	Max	Min	SEM
Ankpa	Odagbo	1.36	0.1231NS	1.494	1.175	0.0399
	Oje Elanyi	1.42				
	Ojogobi Olaji	1.35				
	Okabo	1.39				
Dekina	Ayingba	1.36	0.1447 NS	1.509	1.152	0.0470
	Dekina	1.40				
	Egume	1.27				
	Odu Ogbaloto	1.40				
Ofu	Ogbulu	1.43a	0.1256	1.520	1.092	0.0408
	Ugwolawo	1.37a				
	Ejule	1.32ab				
	Ochadamu	1.23b				
Olamaboro	Ejoka	1.38a	0.1007	1.450	1.009	0.0327
	Igoti Ade	1.28a				
	Unobe	1.16b				
	Ubalu	1.15b				
Omala	Ajedibo	1.44	0.1397 NS	1.565	1.143	0.0453
	Ajomakoji	1.47				
	Odumukpo	1.41				
	Okugba	1.45				

Note: Means in a column with different letters are statistically significant at 5 % level of probability. NS = Not significant; LSD = Least significant difference; Max = Maximum; Min = Minimum; SEM = Standard Error of Mean.

On the other hand, the positive contribution of clay content of soils to soil bulk density have been reported in previous study [24]. Nonetheless, soil compaction arising from higher clay content affects the growth, distribution, and function of roots, and crop productivity [25,26].

3.2.2 Bulk density of soils from oil palm agroforestry systems in Kogi East, Nigeria

The results obtained from all the studied locations revealed that the soil bulk density of oil palm agroforestry systems in Kogi East were within the range of 1.15 -1.47 g/cm³ which further indicate that the soils have a range of bulk density required for arable crop production. The critical value of bulk density for restricting root growth varies with soil type – Soil texture [27] but in general, bulk densities > 1.6 g/cm³ tends to restrict root growth [28]. Based on soil texture, ideal bulk density for plant growth are < 1.60, < 1.40, and < 1.10 g/cm³ for sandy, silty, and clayey soils respectively while bulk density that restricts plant growth are >1.80, >1.65, and >1.47 for sandy, silty, and clayey soils respectively [1,2,3]. It is generally desirable to have soils with low bulk density (<1.5 g/cm³) [27] for optimum movement of air and water through the soil as well as suitability for root growth [29, 28]. Furthermore, bulk density of soils within acceptable thresholds contribute to water and solute transport which are vital for evaluating soil carbon as well as nutrient reserves [30].

4. SUMMARY AND CONCLUSION

This study was conducted to assess the bulk density of soils from selected oil palm agroforestry systems in Kogi East (5 local government areas: Ankpa, Dekina, Ofu, Olamaboro and Omala). Study locations were comprised of 4 communities in each LGA. Random soil sampling was used to collect a total of 100 core samples from the five LGAs (20 from each of the LGA). The bulk density of soils were determined using core sampler technique. Where samples contained in the core rings of known weight, height and diameter were weighed and the fresh weight recorded, then oven dried at 105⁰C for 24hours which the dried weight were also recorded. Data collected were analysed using Analysis of Variance (ANOVA) of GENSTAT Discovery Software. Significant means were separated using Duncan Multiple Range Test at 5 %level of probability.

The results of the analysis of the bulk density of the AFS showed moderate values of 1.15-1.47g/cm³. This implies that the soils of the agroforestry systems in Kogi East can enhance soil bulk density for crop production. Most of the soils in the studied locations were observed to be sandy clay: Ofu and Olamaboro LGAs were sandy clay while Dekina, Ankpa, Omala were loamy sand, sand, and sandy loam respectively.

ACKNOWLEDGEMENTS

This work was funded by grant under Tertiary Education Trust Fund (TETFund) institutional based research (2014-2015 merged intervention), Kogi State University Anyigba, Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aubertin GM, Kardos LT. Root growth through porous media under controlled conditions: I. Effect of pore size and rigidity. *Soil science society of America Journal*. 1965;29(3):290-293.
2. Jones CA. Effect of soil texture on critical bulk densities for root growth. *Soil Science* Keesstra, S., Pereira, P., Novara, A., Brevik, E.C., Azorin-Molina, C., Parras-Alcántara, L., Jordan, A., Cerda, A. (2016). Effects of soil management techniques on soil water erosion in apricot orchards. *Sci Total Environ*. 1983;551-552: 357–366.
3. Arshad MA, Lowery B, Grossman B. Physical tests for monitoring soil quality. *Methods for assessing soil quality*. 1997; 49:123-141.
4. Al-Shammary AAG, Kouzani AZ, Kaynak A, Khoo SY, Norton M, Gates W. Soil bulk density estimation methods: A review. *Pedosphere*. 2018;28(4):581-596.
5. Chaudhari PR, Ahire DV, Ahire VD, Chkravarty M, Maity S. Soil bulk density as related to soil texture, organic matter content and available total nutrients of Coimbatore soil. *International Journal of Scientific and Research Publications*. 2013;3(2):1-8.

6. World Agroforestry Centre. People and Agroforestry. Trees on Farms: 2011 International Year of Forests Agroforestry; 2012.
Available:<http://treesonfarms.com/agroforestry/people-and-agroforestry>
7. Amadu FO, Miller DC, McNamara PE. Agroforestry as a pathway to agricultural yield impacts in climate-smart agriculture investments: Evidence from southern Malawi. *Ecological Economics*. 2020;167:106443.
8. Elagib NA, Al-Saidi M. Balancing the benefits from the water–energy–land–food nexus through agroforestry in the Sahel. *Science of the Total Environment*. 2020;742:140509.
9. Cornell JD, Miller M. Agroforestry. In: *Encyclopedia of Earth*. Eds. Cutler J. Cleveland (Washington, D.C.: Environmental Information Coalition, National Council for Science and the Environment); 2011.
Available:<http://www.eoearth.org/article/Agroforestry>
10. Pavlidis G, Tsihrintzis VA. Environmental benefits and control of pollution to surface water and groundwater by agroforestry systems: A review. *Water Resources Management*. 2018;32(1):1-29.
11. Nair PKR. Agroforestry systems and environmental quality: Introduction. *Journal of Environmental Quality*. 2011;40:784–790.
12. Pardon P, Reubens B, Reheul D, Mertens J, De Frenne P, Coussement T, Verheyen K. Trees increase soil organic carbon and nutrient availability in temperate agroforestry systems. *Agriculture, Ecosystems and Environment*. 2017;247:98-111.
13. Agriani FF, Ranuda R, Wawan W, Qomar N. The physical properties of soil in palm oil agroforetrial system (*Elaeis guineensis* Jacq.) with Aloes (*Aquilaria malacensis* Lamk.) and in palm oil monoculture system. *Jurnal Agronomi Tanaman Tropika (JUATIKA)*. 2021;3(1):15-28.
14. Khasanah N, Van Noordwijk M, Slingerland M, Sofiyudin M, Stomph D, Migeon AF, Hairiah K. Oil palm agroforestry can achieve economic and environmental gains as indicated by multifunctional Land Equivalent Ratios. *Frontiers in Sustainable Food Systems*. 2020;3:122.
15. Rahmani TA, Nurrochmat DR, Hero Y, Park MS, Boer R, Satria A. Evaluating the feasibility of oil palm agroforestry in Harapan Rainforest, Jambi, Indonesia. *Forest and Society*. 2021;458-477.
16. Ali H. *Fundamentals of irrigation and on-farm water management: Volume 1 (Vol. 1)*. Springer Science and Business Media; 2010.
17. Bauder J. *methods of soil analysis Part 2. (Soil Science Society of America Monograph no 9)*. (Edited by Miller, R. H. and Keeney, D. R). Published by: American Society of Agronomy, Madison, Wisconsin, USA. 1985;891–901.
18. USDA. *Texture*. United State Department of Agriculture (USDA). Conservation service; 2008.
19. Jaja N. *Understanding the texture of your soil for agricultural productivity*; 2016.
20. Alotaibi KD, Cambouris AN, St. Luce M, Ziadi N, Tremblay N. Economic optimum nitrogen fertilizer rate and residual soil nitrate as influenced by soil texture in corn production. *Agronomy Journal*. 2018;110(6):2233-2242.
21. Kurniawan S, Corre MD, Matson AL, Schulte-Bisping H, Utami SR, Straaten OV, Veldkamp E. Conversion of tropical forests to smallholder rubber and oil palm plantations impacts nutrient leaching losses and nutrient retention efficiency in highly weathered soils. *Biogeosciences*. 2018;15(16):5131-5154.
22. Hazelton P, Murphy B. *Interpreting soil test results: What do all the numbers mean?* CSIRO publishing; 2016.
23. Walpola BC, Arunakumara KKIU. Decomposition of *Gliricidia* leaves: the effect of particle size of leaves and soil texture on carbon mineralization. *Tropical Agriculture Research*. 2010;13:19-23.
24. Heuscher SA, Brandt CC, Jardine PM. Using soil physical and chemical properties to estimate bulk density. *Soil Science Society of America Journal*. 2005;69(1):51-56.
25. Gregorich EG, Lapen DR, Ma BL, McLaughlin NB, Vanden Bygaart AJ. Soil and crop response to varying levels of compaction, nitrogen fertilization, and clay content. *Soil Science Society of America Journal*. 2011;75(4):1483-1492.
26. Gregorich EG, McLaughlin NB, Lapen DR, Ma BL, Rochette P. Soil compaction, both an environmental and agronomic culprit: Increased nitrous oxide emissions and

- reduced plant nitrogen uptake. Soil Science Society of America Journal. 2014; 78(6):1913-1923.
27. Hunt N, Gilkes R. Farm monitoring handbook. The University of Western Australia: Nedlands, WA; 1992.
 28. McKenzie NJ, Jacquier D, Isbell R, Brown K. Australian soils and landscapes – An illustrated compendium. (CSIRO Publishing: Melbourne); 2004.
 29. Cresswell HP, Hamilton. Particle size analysis. In: Soil Physical Measurement and Interpretation for Land Evaluation. (Eds. NJ McKenzie, HP Cresswell and KJ Coughlan) CSIRO Publishing: Collingwood, Victoria. 2002;224-239.
 30. Yang Q, Luo W, Jiang Z, Li W, Yuan D. Improve the prediction of soil bulk density by cokriging with predicted soil water content as auxiliary variable. Journal of soils and sediments. 2016;16(1):77-84.

© 2021 Amhakhian et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78103>