



Increased Beneficial Outcome of Insulin by Medical Ozone against Experimentally Diabetic Rats

Shaymaa Z. AL Rumaidh ^{a*} and Bushra F. Hasan ^b

^a *Biology Department, Collage of Science, University of Thi-Qar, Iraq.*

^b *Department of Physiology, Pharmacology and Biochemistry, College of Veterinary Medicine, University of Basrah, Iraq.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.56557/UPJOZ/2024/v45i73974

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.mbimph.com/review-history/3345>

Original Research Article

Received: 14/01/2024

Accepted: 18/03/2024

Published: 26/03/2024

ABSTRACT

The present research aimed to determine the effective dose of medical ozone on some blood parameters in experimental diabetic rats. 48 adult female rats with (a 180 - 200 g) weight average were randomly divided into (6) groups (n=8): control (T1), Medical ozone (MO) (T2), diabetic (T3), ozone-treated diabetic (T4), insulin-treated diabetic (T5), and MO, insulin-treated diabetic (T6). Diabetes was induced by an intraperitoneal injection of a single dose of Alloxan (150 mg/kg of the body weight). The diabetic rats were left for 2 days to assure the stability of diabetes in them, before continuing experiments. The results of RBCs in groups (T2, T4, T5, and T6) showed a significant increase compared with the diabetic group and control group, and Hb and WBCs parameters showed a significant decrease in (T3, T4, and T5) when compared with other groups. The results of the platelet count there was a significant increase in (T3, T4, and T5) compared to the control group. Clotting time and calcium ions recorded a significant decrease in all treated diabetic groups (T3, T4, T5, and T6) when compared with (T1 and T2), at the same time, the results of bleeding time recorded a significant decrease in (T3 and T4) compared with the other

*Corresponding author: Email: shymaa1980@utq.edu.iq;

groups while (T4 and T5) showed non-significant difference compared to (T1 and T2). The current study concluded that the treatment combination of medical ozone and insulin has ameliorative effects on some blood parameters except clotting time and calcium, ions.

Keywords: Medical Ozone; insulin; alloxan; hematological; parameters.

1. INTRODUCTION

“Diabetes mellitus (DM) is a chronic metabolic disorder in which the body's ability to produce or respond to the hormone insulin is impaired, resulting in abnormal metabolism of carbohydrates and elevated levels of glucose in the blood and urine. DM has been diagnosed in canine and feline families after humans. The clinical features described and investigated are rarely observed in other domestic large animals such as horses, cattle, buffalo, swine, and other small ruminants” [1-3]. “The classification of diabetes differs for large and small animals, although it has similarities to humans. The common and general forms of DM are known to be insulin-dependent DM (IDDM) Type-1 and non-IDDM (NIDDM) Type-2 in animals. Also, the secondary DM or Type-3 has been identified, which is a complication of insulin antagonisms. This occurs due to pancreatic islet damage by pancreatic necrosis, tumor progression, and pancreatitis. Metabolic DM is the specific experimental expression of this form, primarily described in dogs and cats” [2,4].

“Ozone, a gas composed of three atoms of oxygen with a cyclic structure, was initially discovered as an oxidant and a disinfectant in 1834, exerting medical affectivity first for gunshot gangrene” [5]. “Evidence supports ozone has been used for the treatment of cutaneous wounds with satisfactory healing results” [6]. “Ozone was used in many methods, tent, bag, even injection, and systematic applications referring to rectal insufflation as well as autohemotherapy” [7].

“Ozone has been used as a therapeutical agent and beneficial effects have been observed. However, so far only a few biochemical and pharmacodynamic mechanisms have been

elucidated. many studies demonstrated that controlled ozone administration may promote oxidative preconditioning or adaptation to oxidative stress, preventing the damage induced by ROS” [1]. Taking into account that diabetes is a disorder associated with oxidative stress, we postulate that ozone treatment might protect antioxidant systems and maintain, at a physiological level, other markers of damage associated with diabetic complications.

2. MATERIALS AND METHODS

The study was carried out on (48) adult female albino rats weighing (180-200g). Animals were housed in temperature and light- controlled rooms and allowed free access to normal diet pellets and tap water. Medical Ozone (MO) was generated with (Medical Ozone generator Aqua plus, model No: AOT-MD-520). MO given to each animal was adjusted to a final dose of 1.1 mg/kg BW intraperitoneal [8]. Huminsulin R India was used at a dose of 0.75 IU/100 g body weight [9].

2.1 Induction of Diabetes Mellitus

Diabetes mellitus (DM) was induced in overnight fasting rats by a single injection of alloxan (alloxan monohydrate) at a dose of 150 mg/kg body weight into intraperitoneal(I.P). After injection alloxan. Rats were given a 5% glucose solution for 24 hours with drinking water to prevent initial drug-induced hypoglycemic mortality.

2.2 Experimental Design

The mature female rats were randomly divided into six groups each containing eight animals as follows:

-
- | | |
|-----------|---|
| T1 | Animals were daily administered (I.P) with citrate buffer (0.1 M, pH 5). daily for six weeks. |
| T2 | Animals daily administered 1.1 mg/kg of MO (I.P), daily for six weeks. |
| T3 | Animals Untreated diabetic group will induced with single (I.P) injections of alloxan (150 mg/kg) |
| T4 | Diabetic rats will receive 1.1 mg/kg of MO (I.P), for six weeks. |
| T5 | Diabetic rats will receive the dose of Mixtard insulin and will be injected subcutaneously at a dose of 0.75 IU/100 g body weight, for six weeks. |
| T6 | Diabetic rats will receive 1.1 mg/kg of MO (I.P) and Mixtard insulin will injected subcutaneously at a dose of 0.75 IU/100 g body weight, respectively, for six weeks |
-

2.3 Determination of Bleeding Time (BT)

The rat tail was warmed for one minute in water at 40°C and then dried. A small cut was made in the middle of the tail with a scalpel. Bleeding time started when the first drop touched the circular filter paper. It was checked at 30-second intervals until the paper was no longer stained with blood. The time taken for bleeding to stop was recorded for each rat and the average was taken as bleeding time [10].

2.4 Clotting time Estimation (CT)

Clotting time was determined using the capillary glass tube method. A standard incision was made in the skin of the ear and the blood was taken into a capillary glass tube and the time of collection was noted. Pieces of capillary glass were broken from one end every thirty seconds and the appearance of fibrin threads was used as the endpoint and the time was noted in seconds [11].

2.5 Collection of Blood Samples

On the 60th day of the experiment, Each of the rats was anesthetized. 10 ml blood samples were collected from each sacrificed rat by cardiac

puncture and 5 ml of blood was discharged into blood tube EDTA K for hematological analysis and the residual amount was discharged into a gel tube to obtain the serum for biochemical analysis. Hematological parameters that were analyzed include Red Blood Cells (RBCs), White Blood Cells (WBCs), Haemoglobin (Hb), and Platelets. They were determined using Automated Hematology Analyzer Genex(COUNT 60). A calcium test kit is used to measure in serum(BIOLABO kit, 052010A).

The results obtained from the current study recorded a significant ($p > 0.001$) rise in RBC count in the (T2) treated group when compared with the control group and the rest of the diabetic groups. as well as noticed that there was no significant difference in the number of RBC between the groups (T1, T4, T5, and T6), also a significant decrease was recorded in the diabetic group when it was compared with the control group. As for the results of white blood cells, no statistically significant difference was recorded between the (T2) group and control group, As for the treated diabetes groups, no statistical difference was recorded when comparing them, except (T6) group, which came close to the value of the control group.

Table 1. Effects of Medical Ozone on some hematological parameters of diabetic female rats

Groups n=8	Parameters		
	RBCs ($10^6/\mu\text{L}$)	Hb(g/dl)	WBCs($10^3/\mu\text{L}$)
T1	7.8642 ± 0.2364 ^b	13.8250 ± 0.2246 ^b	7.71 ± 0.47 ^{ab}
T2	9.0783 ± 0.1595 ^a	14.4250 ± 0.1605 ^a	8.05 ± 0.47 ^a
T3	6.0575 ± 0.2153 ^c	10.1917 ± 0.1367 ^d	6.00 ± 0.48 ^c
T4	7.4225 ± 73.976 ^b	10.6583 ± 0.1389 ^{cd}	6.37 ± 0.66 ^c
T5	7.3983 ± 73.9788 ^b	10.3583 ± 0.2224 ^d	6.26 ± 0.62 ^c
T6	7.8892 ± 0.1807 ^b	10.9000 ± 0.1692 ^c	7.52 ± 0.73 ^b
LSD	0.60	0.50	0.47

Values expressed in small letters mean significant differences at p . value < 0.001

Table 2. Effects of Medical Ozone on hemostatic parameters of diabetic female Rats

Groups n=8	Parameters			
	Platelets count x $10^3/\text{MI}$	Clotting Time	Bleeding Time	Calcium
T1	457.0833 ± 55.9 ^d	2.0333 ± 0.22293 ^a	3.5833 ± 0.41090 ^a	7.1917 ± 0.69473 ^a
T2	473.5000 ± 48.9 ^{cd}	1.8833 ± 0.15859 ^a	3.2792 ± 0.33345 ^b	7.3833 ± 0.67532 ^a
T3	769.4167 ± 58.6 ^a	0.9542 ± 0.13049 ^d	1.9583 ± 0.37005 ^c	5.0375 ± 0.55478 ^c
T4	571.0000 ± 39.6 ^b	1.1058 ± 0.16082 ^c	2.0117 ± 0.42000 ^c	5.1083 ± 0.46993 ^c
T5	553.8333 ± 46.6 ^b	1.9833 ± 0.14035 ^b	3.2750 ± 0.249930 ^b	5.1500 ± 0.61274 ^c
T6	493.0000 ± 30.6 ^c	1.8750 ± 0.171230 ^b	3.2708 ± 0.33671 ^b	5.6417 ± 0.47760 ^b
LSD	38.8	0.13	0.29	0.47

Values expressed in small letters mean significant differences at p . value > 0.001

The results of hemoglobin value in the current study noticed the T2 group recorded a statistical increase in the level of hemoglobin when compared with the T1 group. While the groups of diabetics were given doses of (MO)T2 or (Insulin) T4, they recorded a significant increase when compared with other diabetic groups

The platelet count results showed a significant increase in the (T3)group compared to the other groups, while the (T2) group did not show any significant difference when compared with the control group. At the same time, there was no significant difference between (T4) and (T5), but compared to(T3) there was a significant decrease. López [12] while the (T6)group recorded a significant decrease compared to the (T3) group In addition to being close to reading of the (T2) group.

Results showed a significant decrease ($p > 0.001$) in bleeding time in the (T2) and (T3) groups when compared to the (T1), at the same time showed the (T4) group non-significant difference in bleeding time compared to (T3) group while the (T5 and T6) recorded significant increase when compared to (T3) group.

The results of clotting time showed a significant decrease in the (T2) group when compared with the control group, while the (T3) group showed a significant decrease compared to the control group.

On the other hand, all treated diabetic groups recorded a significant increase in clotting time compared to the (T3) group and at the same time, both groups (T5 and T6) were approached to the reading of the (T2) group. The obtained results are in the tab. (2) revealed a non-significant change in Ca^{++} in serum of female rats when compared between the (T2) group and (T1) group while the results showed a significant decrease in serum Ca^{++} of diabetic group (T3) compared with the control group. At the same time, there was a non-significant difference between (T4), and (T5) compared to (T3) group.

3. DISCUSSION

In a study by Lopez [13] that used medical ozone autohemotherapy in horses, a significant increase in red blood corpus was detected; this result is similar to observations in the present study. "Ozone therapy acts as a form of oxygen therapy that stimulates mitochondria and gives them a bath of fresh air. Oxygen is the most vital

element required for human life and it is the key to good health, the best way to optimize health is to oxygenate every cell in the body" [14].

The results of the present study supported by Bocci [15], and Mahmoud et al. [14] found that ozone therapy increases the oxygen level in the blood, oxygenates the tissues in the whole body, and can activate biochemical pathways in erythrocytes, hemoglobin, and leukocytes without acute or chronic toxicity. "during prolonged ozone therapy, the bone marrow may release a cohort (about 0.9% of the pool) of new erythrocytes with improved biochemical characteristics" [16].

The appreciable decrease in RBCS count, Hb and WBCS count in the untreated alloxan-induced diabetic rats correlates with the findings of Mansi [17] and Mounce and AL-Saeed, [18]. "Reactive oxygen species have also been implicated in the mechanism of RBCS damage" [19]. "The cytotoxic action of diabetogenic agents such as alloxan is mediated by reactive oxygen species" [20].

"In diabetic rats, red blood cell (RBC) count was found to decrease significantly, due to the non-enzymatic glycosylation of RBCS membrane protein which is directly associated with the hyperglycemic condition. High glucose level leads to the generation of toxic product alteration which causes many such as reduced production of bone marrow" [21]. "Hyperglycemia results in glycosylated hemoglobin, thus total hemoglobin level is decreased in alloxan-induced diabetic rats" [22]. "Reduction in hemoglobin may be accompanied by a fall in the red blood corpus count and packed cell volume" [23].

"The hyperglycemic situation leads to the production of lipid peroxides. These peroxides increase membrane stiffness, decrease cellular deformability, decrease the life span of RBCs and lipid fluidity, and also cause hemolysis of RBCs and low hemoglobin" [24]. "The previous study recorded that WBCs which play a pivotal role in initiating immunity showed a significant decrease ($P > 0.05$ and $P > 0.01$ respectively) In untreated diabetic rats" [25].

"According to the current results, observed that the addition of MO with daily doses of insulin does not affect the action of insulin, as a significant increase in the number of RBCs and WBCs. The combination of metabolic actions of insulin and the ability of ozone to enhance antioxidant endogenous systems, in addition to

oxidative stress by oxidative preconditioning mechanisms, resulted in a reduction in lipid and glucose oxidation and increased NO oxidation caused by insulin alone plus antioxidant supplementation of ozone can normalize all parameters" [26]. "Ozone may increase the endogenous antioxidant defense mechanism in rats, thereby protecting them from cranial and spinal tissue neuropathies. Moreover, the combination of ozone and Insulin therapy has a better prospectus for treating DM" [27].

"Diabetes-associated reductions in the antioxidant state can be ameliorated by insulin combined with antioxidant therapy. These results indicate that the combination of insulin treatment with ozone in calculated doses leads to a more significant enhancement in metabolic oxidative states and vascular complications in diabetic rats" [28].

"Administration of insulin and exogenous antioxidants significantly normalized the RBC count because of the ability of antioxidants to scavenge the peroxides, reduce membrane damages, and simultaneously insulin competes for the regulation of glucose metabolism and hence prevents glycosylation of proteins" [29].

Ozone causes a decrease in blood and plasma viscosity (plasma macromolecules are decreased). Moreover, ozone may induce a hypo-coagulatory state thus decreasing the tendency to clot [30], This agrees with the results of the current study.

In the present study diabetes rats (DM) group showed an increment in platelet count. It is reported that platelet hyperactivity in patients with hyperglycemia results from dysregulated signaling pathways that lead to an increased tendency to activate and aggregate response to a given stimulus [31], Platelet activation therefore triggers thrombus formation, microcapillary embolization and facilitates the development of other cardiovascular diseases. Akingbami et al. [32] added that diabetes mellitus is characterized by enhanced platelets activation and coagulation proteins and reduced fibrinolytic activity which usually precede the development of cardiovascular complications.

As shown in Table (2), It was observed that alloxan decreased the bleeding time and clotting time. The increased platelet counts in the diabetic rats may have been responsible for the observed decrease in bleeding and clotting times

and may increase the risk of intravascular blood clotting and associated diseases [29]. As for the results of the calcium level in the serum, a significant decrease was recorded in the diabetic groups when it was compared with the (control and MO) groups this is consistent with a previous study [33].

"In many research studies, a relationship between trace elements and diabetes mellitus was observed, in different studies an alteration in the metabolism of these trace elements (calcium and magnesium was noted" [34]. "Calcium ion plays an important role in glycemic control by affecting the biosynthesis and release of insulin from beta cells of the pancreas. There is a significant correlation between diabetes and hypocalcemia" [35].

The current study found that after ozone therapy there are no statistically significant changes in hemostatic parameters, at the same time MO plus insulin might provide a simpler, safer, and more effective means for preventing and ameliorating chronic complications of DM than each given alone perhaps because of their action as an antioxidant.

"The combination of metabolic actions of insulin and the ability of ozone to enhance antioxidant endogenous systems, in addition to oxidative stress by oxidative preconditioning mechanisms, resulted in up-regulation of NOS isoforms and reduction in lipid and glucose oxidation and increased NO oxidation caused by insulin alone plus antioxidant supplementation of ozone can normalize all parameters" [36,37].

4. CONCLUSION

Finally, from the previous discussion of these results and according to reports of other investigators in similar studies: It can be explained that ozone therapy has a significant effect on the improvement of diabetic patients undergoing insulin treatment as evidenced by the improvement in hematological and hemostatic parameters. The results of the current study would introduce a scientifically applicable protocol to help, physicians dealing with diabetes, organize a plan of care to overcome this problem and prevent complications of it and the development of associated morbidities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Catchpole B, Adams JP, Holder AL, Short AD, Ollier WE, Kennedy LJ. Genetics of canine diabetes mellitus: Are the diabetes susceptibility genes identified in humans involved in breed susceptibility to diabetes mellitus in dogs? *Vet. J.* 2013;195:139-147.
2. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2014;37:S81-S90.
3. Robinson WF, Robinson NA, Maxie MG. Jubb, Kennedy, and Palmar's Pathology of Domestic Animals. Elsevier, St. Louis, MO. 2016:1-101.
4. Niaz K, Maqbool F, Khan F, Hassan FI, Momtaz S, Abdollahi M. Comparative occurrence of diabetes in canine, feline, and few wild animals and their association with pancreatic diseases and ketoacidosis with a therapeutic approach. *Veterinary world,* 2018;11(4):410.
5. Bocci V. Ozone as Janus: this controversial gas can be either toxic or medically useful. *Mediat Inflamm.* 2004;13(1):3- 11
6. Martínez-Sánchez G, Al-Dalain SM, Menéndez S, et al. Therapeutic efficacy of ozone in patients with diabetic foot. *Eur J Pharmacol.* 2005;523(1-3):151- 161.
7. Bocci V, Zanardi I, Huijberts MS, Travagli V. An integrated medical treatment for type-2 diabetes. *Diabetes Metab Syndr.* 2014;8(1):57- 61
8. Morsy MD, Hassan WN, Zalat SI. Improvement of renal oxidative stress markers after ozone administration in diabetic nephropathy in rats. *Diabetology & metabolic syndrome.* 2010;2(1):1-7.
9. Saleh S, El-Ridi M, Zalat S, El-Kotb S, Donia S. Additive effect of ozone therapy to insulin in the treatment of diabetic rats. *Menoufia Medical Journal.* 2014;27(1):85.
10. García-Manzano A, González-Llaven J, Lemini C, Rubio-Póo C. Standardization of rat blood clotting tests with reagents used for humans. *Proceedings of the Western Pharmacology Society.* 2001;44:153-155
11. Harris DT, Gilding HP, Smart WAM. *Experimental Physiology for Medical Students.* 6th Edn., Sagar Publications, New Del; 1956.
12. Hajam YA, Rai S, Ghosh H, Basheer M. Combined administration of exogenous melatonin and insulin ameliorates streptozotocin induced toxic alteration on hematological parameters in diabetic male Wistar rats. *Toxicology reports.* 2020;7: 353-9.
13. López LG. Efecto de la Ozonoterapia sobre los parámetros hematológicos y constantes fisiológicas en equinos; 2007.
14. Mahmoud Adel Abdel-Hakim, Emad Tawfik Ahmed, Fady Samy Faltaous, Haidy Nady Asham Effect of Ozone Therapy on Anemic Cancer Patients Undergoing Chemotherapy, *European Journal of Applied Sciences.* 2020;12(2):56-61
15. Bocci V. The case for oxygen-ozone therapy. *British journal of biomedical science.* 2007;64(1):44-49.
16. Bocci V: *Ozone. A new medical drug.* Dordrecht, The Netherlands: Springer; 2011.
17. Mansi KMS. Effects of administration of alpha-melanocyte-stimulating hormone (α -MSH) on some hematological values of alloxan-induced diabetic rats. *Am J Pharmacol Toxicol.* 2006;1:5-10.
18. Mounce FS, AL-Saeed MH. Study the effect of phytoesterol of ceratoina siliqua fruit and insulin on hematological and biochemical parameters in diabetic pregnant female rabbits induced by Alloxan. *Basrah Journal of Veterinary Research.* 2017;16(1):219-242.
19. Rao GU, Kamath C, Raghothama, KSP, Rao P. Maternal and fetal indicators of oxidative stress in various obstetric complications. *Ind J Clin Biochem.* 2003; 18:80-86.
20. Szkudelski T. The mechanism of alloxan and streptozotocin action B cells of the rat pancreas. *Physiol Res.* 2001;50:536-546.
21. Abbas M, Siddiqi MH, Khan K, Zahra K. Hematological evaluation of sodium fluoride toxicity in *oryctolagus cuniculus*. *Toxicology Reports.* 2017;4:450-454.
22. Sheela CG, Augusti KT. Antidiabetic effects of S-allyl cysteine sulphoxide isolated from garlic *Allium sativum* Linn. *Ind J Exp Biol.* 1992;30:523-6.
23. Muhammad NO, Oloyede OB. Hematological parameters of broiler chicks fed *Aspergillus niger* - fermented *Terminalia catappa* seed meal-based diet. *Global J Biotechnol Biochem.* 2009;4:179-183.
24. Kolanjiappan K, Manoharan S, Kayalvizhi M. Measurement of erythrocyte lipids, lipid peroxidation, antioxidants, and osmotic fragility in cervical cancer patients. *Clinica Chimica Acta.* 2002;326(1-2):143-149.

25. Helal EG, Abd-Elwahab SM, Mohammad AA. Comparison between the effect of ozone and vitamin C in the treatment of diabetes mellitus. *The Egyptian Journal of Hospital Medicine*. 2013;51(1):434-447.
26. Sindhu RK, Koo JR, Roberts CK, Vaziri ND. Dysregulation of hepatic superoxide dismutase, catalase, and glutathione peroxidase in diabetes: response to insulin and antioxidant therapies. *Clinical and experimental hypertension*. 2004;26(1):43-53.
27. Karademir MUSTAFA, Sönmez M, Akcılar RAZİYE, Koçak FATMA, Yay A, Eser O. Evaluation of the therapeutic potential of intraperitoneal ozone gas in combination with insulin above cranial and spinal neuropathy in rats with diabetes mellitus. *bratislava medical journal-bratislavske lekarske listy*. 2018;119(10).
28. Saleh S, El-Ridi M, Zalat S, El-Kotb S, Donia S. Additive effect of ozone therapy to insulin in the treatment of diabetic rats. *Menoufia Medical Journal*. 2014;27(1):85.
29. Hajam YA, Rai S, Ghosh H, Basheer M. Combined administration of exogenous melatonin and insulin ameliorates streptozotocin-induced toxic alteration on hematological parameters in diabetic male Wistar rats. *Toxicology reports*. 2020;7: 353-359
30. Van Der Zee H, De Monte A. Ozone auto-chemotherapy in lower limb ulcerations. In *Proceedings of the 15th Ozone World Congress*. London: Speedprint MacMedia Ltd (London). 2001:148-157.
31. Kakouros N, Rade JJ., Kourliouros A, Resar JR. Platelet function in patients with diabetes mellitus: From a theoretical to a practical perspective. *International Journal of Endocrinology*; 2011.
32. Akingbami A, Dada AO, John OS, Ushanaiki O, Adediran A, Odesanya M, Ogbara A, Uche E, Okunoye O, Arogundade O, Aile K. Mean platelets volume and platelets count in type 2 diabetes mellitus patients on treatment and nondiabetes mellitus controls in Lagos, Nigeria. *The Pan African Medical Journal*. 2014;18.
33. Tahir IM, Akhter N, Parveen A, Mehboob H, Saleem S, Munir N, Akram M. Effects of methanolic and aqueous extracts of Ipomoea batatas L on mineral contents level (calcium and magnesium) in alloxan-induced diabetic rats. *Pakistan Journal of Pharmaceutical Sciences*; 2018.
34. Hans P, Sialy R and Bansal D. Magnesium deficiency and diabetes mellitus. *Curr. Sci.*, 2002;83:12-25.
35. Al-Yaseen, Nimnim T, Challab D Calcium and diabetes mellitus type Two a prospective study done on people with type 2 diabetes in Diwaniya Teaching Hospital. *Kufa Med. J*. 2009;12(1):468-475.
36. Sindhu RK, Koo JR, Roberts CK, Vaziri ND. Dysregulation of hepatic superoxide dismutase, catalase, and glutathione peroxidase in diabetes: response to insulin and antioxidant therapies. *Clinical and experimental hypertension*. 2004;26(1):43-53
37. Falih S, HASAN BF. Medical Ozone treatment alleviate blood oxidative stress and pancreas damage in an alloxan-induced diabetes model in Rats. *University of Thi-Qar Journal of Science*. 2023; 10(2):1-9.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://prh.mbimph.com/review-history/3345>