



Growth Analysis of Coriander (*Coriandrum sativum* L.) as Influenced by Different Sowing Dates and Planting Geometry in Chhattisgarh Plains Zone

Sandeep Kumar Painkara ^a, D. K. Chandrakar ^b,
Golmei Langangmeilu ^{b*}, P. M. Paraye ^a,
A. K. Verma ^b and Anjali Patel ^b

^a College of Agriculture and Research Station, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, 492012, India.

^b College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, 492012, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i34088

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113760>

Original Research Article

Received: 21/12/2023

Accepted: 26/02/2024

Published: 01/04/2024

ABSTRACT

An experiment was conducted to assess the effect of different sowing dates and planting geometry on coriander during the winter seasons of 2020-21 and 2021-22 at Instructional farm, College of Agriculture and Research Station, IGKV, Raigarh, Chhattisgarh. The experiment was laid out in split-plot design, comprised of four date of sowing viz., D1: 25th October, D2: 10th November, D3: 25th November and D4: 10th December as main plot and four planting geometries viz., S1: 30 x 5 cm, S2: 30 x 7.5 cm, S3: 30 x 10 cm and S4: 30 x 12.5 cm as sub-plot, and replicated thrice. Influence of above treatments on crop was studied regarding growth parameters of coriander and analyzed statistically. The result showed that the crop sown on 25th October recorded significantly

*Corresponding author: E-mail: langangmeilu2@gmail.com;

higher growth parameters viz., plant height, number of branches, dry matter accumulation, crop growth rate and relative growth rate. Among planting geometry, sowing of coriander at 30 x 12.5 cm spacing produced significantly taller plants, higher number of branches plant-1, dry matter accumulation plant-1, crop growth rate and relative growth rate. Coriander sown on 25th October interacted with the 30 x 12.5 cm spacing and registered significantly higher dry matter accumulation.

Keywords: Coriander; date of sowing; planting geometry.

1. INTRODUCTION

Coriander (*Coriandrum sativum* L.), which belongs to the family of umbelliferae (Apiaceae) is one of the most important annual spice and medicinal herb, native to the Eastern Europe and Asia. Coriander is commercially grown in India, Morocco, Romania, USSR, Mexico, Argentina, Hungary, Poland, Bulgaria and USA. In India it is mainly grown in the states of the Madhya Pradesh, Rajasthan, Gujrat, Assam, Odisha, West Bangal, Andhra Pradesh and Uttar Pradesh in a limited extent, it is also grown in Karnataka, Uttar Pradesh, Bihar, Tamil Nadu, Chhattisgarh and Punjab. [1].

It is grown as a field crop for its seeds commonly known as "*Dhania*" which has a fragrant odor and aromatic test. The dry seeds are said to have carminative diuretic, stomachic and aphrodisiac properties. On stem distillation, coriander seeds yield 0.2 to 1.2% essential oil. The major components of essential oil are linalool (67.7 %) followed by 1-pipen (10.5 %), 1-terpinin (9.0 %), geranyl acetate (4 %) and geraniol (1.9 %). The success of coriander production is influenced by genetic, weather and agronomic factors [2]. The maximum seed and essential oil yields are attained only when an appropriate combination of these factors is provided for the plant [3,4]. Date of sowing is an important management factor for almost all seed spices including coriander. Coriander exploits the environment most favorably when it is sown at the optimum time [5], since the sowing date significantly affects the photoperiodic response of plants and determines yields and qualities [6]. Change in sowing time leads to significant changes in weather microclimate and subsequently the performance of the crop. In addition, the physical environment has a profound influence on growth, partitioning and ultimately the yield of coriander. Temperature, humidity, rainfall and other meteorological factors may individually or collectively limit the plant growth and production. The establishment of an appropriate spacing for maintaining the optimum plant population per unit area is the main pre-requisite to obtaining

maximum yields from any crop. Optimal planting arrangements also contribute to enhanced crop growth and development, thereby resulting in increased production yields. To increase the productivity of local and improved varieties of coriander, it is important that farmers adopt the recommended sowing date and planting geometry.

2. MATERIALS AND METHODS

The experiment was laid out for two consecutive *Rabi* seasons of 2020-21 and 2021-22 at Instructional Farm, College of Agriculture and Research Station, Raigarh, IGKV, Raipur, Chhattisgarh, which is situated at 21.9' North latitude, 83.4' East longitude and at an altitude of 215 meters above mean sea level. The experimental soil was loamy sand in texture, slightly acidic in reaction, low in organic carbon and available nitrogen, phosphorus, sulphur and zinc and medium in available potassium, having low moisture retention capacity. The experiment was carried out in split-plot design with sixteen treatment combinations of main and sub-plots, replicated thrice. Four Dates of sowing viz. D₁: 25th October, D₂: 10th November, D₃: 25th November and D₄: 10th December were arranged in main plot and four planting geometries viz., S₁: 30 x 5 cm, S₂: 30 x 7.5 cm, S₃: 30 x 10 cm and S₄: 30 x 12.5 cm were taken as sub-plot treatments. Recommended dose of fertilizer of 80:60:40 N: P₂O₅: K₂O kg ha⁻¹ was applied. Data on the growth behavior of coriander i.e. plant height, number of branches, dry matter accumulation, crop growth rate and relative growth rate were collected and analyzed statistically using *F*-test, the procedure given by Gomez and Gomez [7]. Critical difference (CD) values at *P*=0.05 were used to determine the significance of mean differences between treatments.

3. RESULTS AND DISCUSSION

Coriander sown on 25th October recorded significantly higher plant population in both seasons, which were not significantly different

Table 1. Growth of coriander as influenced by different date of sowing and planting geometry

Treatment	Plant population (No. m ⁻²)						Plant height (cm)									Number of branches plant ⁻¹									
	20 DAS			At harvest			30 DAS			60 DAS			90 DAS			At harvest			60 DAS			At harvest			
	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	
Date of sowing																									
25 th October	42.62	43.03	42.82	39.92	40.33	40.12	11.13	11.63	11.38	30.19	34.66	32.42	90.77	96.24	93.50	114.40	121.41	117.90	5.83	6.39	6.11	8.32	8.69	8.50	8.50
10 th November	41.69	42.64	42.17	38.99	39.94	39.47	9.74	10.23	9.98	27.93	30.82	29.37	85.68	90.04	87.86	105.89	111.80	108.84	5.27	5.82	5.50	7.73	7.98	7.85	7.85
25 th November	40.78	41.48	41.13	38.08	38.78	38.43	8.80	9.20	9.00	24.89	26.55	25.72	80.12	83.68	81.90	90.98	95.15	93.06	4.72	5.12	4.92	7.26	7.47	7.37	7.37
10 th December	38.61	40.06	39.34	35.91	37.36	36.64	8.13	8.49	8.31	22.77	24.07	23.42	74.83	77.68	76.26	81.04	84.51	82.78	4.38	4.74	4.56	6.72	6.88	6.80	6.80
SEm±	0.71	0.54	0.61	0.63	0.53	0.57	0.17	0.21	0.18	0.72	0.92	0.70	2.49	2.21	2.22	2.86	2.84	3.01	0.15	0.13	0.12	0.17	0.23	0.19	0.19
CD (P=0.05)	2.47	1.86	1.76	2.13	1.81	1.93	0.60	0.72	0.63	2.23	3.18	2.41	8.61	7.63	7.69	9.88	9.82	9.36	0.51	0.44	0.42	0.59	0.79	0.65	0.65
Planting geometry																									
30 cm x 5 cm	63.37	65.65	64.51	59.37	61.65	60.51	10.37	10.78	10.58	31.05	34.53	32.79	87.86	93.13	90.49	105.47	112.18	108.82	4.25	4.55	4.39	6.58	6.86	6.72	6.72
30 cm x 7.5 cm	43.24	43.77	43.51	40.24	40.77	40.51	9.94	10.38	10.16	28.80	31.55	30.18	85.36	90.15	87.76	103.58	108.45	106.02	4.82	5.17	4.98	7.22	7.38	7.30	7.30
30 cm x 10 cm	31.68	32.22	31.95	29.38	29.92	29.65	9.04	9.49	9.27	24.75	26.99	25.87	81.47	85.19	83.33	94.08	99.22	96.65	5.46	6.03	5.73	7.98	8.24	8.11	8.11
30 cm x 12.5 cm	25.40	25.58	25.49	23.90	24.08	23.99	8.44	8.89	8.66	21.18	23.03	22.11	76.71	79.17	77.94	89.18	93.02	91.10	5.68	6.32	5.98	8.25	8.55	8.40	8.40
SEm±	0.47	0.45	0.21	0.47	0.45	0.21	0.19	0.13	0.15	0.52	0.61	0.38	1.74	1.64	1.47	2.12	1.48	1.51	0.16	0.15	0.15	0.20	0.16	0.16	0.16
CD (P=0.05)	1.38	1.31	0.61	1.38	1.31	0.61	0.57	0.39	0.48	1.52	1.77	1.12	5.09	4.80	4.30	6.18	4.32	4.40	0.45	0.44	0.43	0.58	0.46	0.47	0.47
Interaction																									
CD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 2. Growth of coriander as influenced by different date of sowing and planting geometry

Treatment	Dry matter accumulation (g plant ⁻¹)											
	30 DAS			60 DAS			90 DAS			At harvest		
	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean
Date of sowing												
25 th October	0.33	0.37	0.35	3.40	3.69	3.54	11.68	12.18	11.93	13.98	15.19	14.58
10 th November	0.31	0.35	0.33	3.10	3.33	3.22	10.85	11.37	11.11	12.31	13.54	12.92
25 th November	0.29	0.32	0.31	2.74	2.92	2.83	9.98	10.57	10.28	10.81	11.70	11.25
10 th December	0.26	0.29	0.27	2.17	2.29	2.23	8.94	9.22	9.08	9.31	9.93	9.62
SEm±	0.01	0.01	0.01	0.06	0.05	0.05	0.37	0.30	0.32	0.28	0.33	0.23
CD (P=0.05)	0.03	0.02	0.02	0.22	0.17	0.16	1.28	0.92	0.98	0.97	1.14	0.79
Planting geometry												
30 cm x 5 cm	0.20	0.22	0.21	1.77	1.87	1.82	5.02	5.40	5.21	5.62	6.43	6.03
30 cm x 7.5 cm	0.25	0.28	0.27	2.41	2.59	2.50	8.52	9.17	8.85	9.55	10.66	10.10
30 cm x 10 cm	0.35	0.39	0.37	3.53	3.75	3.64	13.29	13.70	13.49	14.78	15.73	15.25
30 cm x 12.5 cm	0.39	0.43	0.41	3.70	4.02	3.86	14.63	15.08	14.85	16.46	17.54	17.00
SEm±	0.01	0.01	0.01	0.09	0.09	0.08	0.23	0.18	0.14	0.27	0.30	0.24
CD (P=0.05)	0.03	0.03	0.03	0.25	0.27	0.22	0.68	0.51	0.40	0.79	0.86	0.69
Interaction												
CD (P=0.05)	NS	NS	NS	S	S	S	S	S	S	S	S	S

Table 3. Interaction effects between different sowing dates and planting geometry on dry matter accumulation of coriander

Dry matter accumulation (g plant ⁻¹) at 60 DAS															
2020-21					2020-21					Mean					
Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	
Spacing					Spacing					Spacing					
S ₁	2.41	1.51	0.70	0.55	S ₁	2.61	1.60	0.95	0.65	S ₁	2.51	1.56	0.83	0.60	
S ₂	2.92	2.44	1.68	0.86	S ₂	3.16	2.59	2.00	0.96	S ₂	3.04	2.52	1.84	0.91	
S ₃	4.05	3.75	3.37	1.54	S ₃	4.36	3.98	3.10	1.88	S ₃	4.21	3.87	3.24	1.71	
S ₄	4.20	3.91	3.57	1.71	S ₄	4.63	4.11	3.62	2.05	S ₄	4.42	4.01	3.60	1.88	
SEm±	0.10				SEm±	0.09				SEm±	0.07				
CD (P=0.05)	0.28				CD (P=0.05)	0.25				CD (P=0.05)	0.22				
Dry matter accumulation (g plant ⁻¹) at 90 DAS															
Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	
Spacing					Spacing					Spacing					
S ₁	6.8	4.8	2.5	2.0	S ₁	6.5	4.7	3.7	2.2	S ₁	6.6	4.7	3.1	2.1	
S ₂	10.6	9.1	6.8	4.6	S ₂	9.9	9.1	7.9	5.1	S ₂	10.2	9.1	7.4	4.9	
S ₃	16.0	14.2	12.1	7.9	S ₃	15.5	14.1	12.4	8.2	S ₃	15.7	14.1	12.2	8.1	
S ₄	17.4	15.3	13.5	9.2	S ₄	16.8	15.3	13.5	10.1	S ₄	17.1	15.3	13.5	9.7	
SEm±	0.70				SEm±	0.44				SEm±	0.27				
CD (P=0.05)	2.05				CD (P=0.05)	1.29				CD (P=0.05)	0.79				
Dry matter accumulation (g plant ⁻¹) at harvest															
Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	
Spacing					Spacing					Spacing					
S ₁	7.4	4.9	2.7	2.1	S ₁	8.5	5.7	4.0	2.6	S ₁	8.0	5.3	3.4	2.3	
S ₂	11.5	9.5	7.3	5.0	S ₂	12.8	10.2	8.6	5.7	S ₂	12.2	9.9	8.0	5.3	
S ₃	17.8	15.0	12.3	8.3	S ₃	19.2	16.2	13.4	8.8	S ₃	18.5	15.6	12.9	8.5	
S ₄	19.9	16.5	14.0	9.7	S ₄	21.4	18.5	14.5	10.5	S ₄	20.7	17.5	14.2	10.1	
SEm±	0.463				SEm±	0.49				SEm±	0.33				
CD (P=0.05)	1.351				CD (P=0.05)	1.42				CD (P=0.05)	0.96				

D₁: 25th October, D₂: 10th November, D₃: 25th November, D₄: 10th December,
 S₁: 30 cm x 5 cm, S₂: 30 cm x 7.5 cm, S₃: 30 cm x 10 cm, S₄: 30 cm x 12.5 cm




Fig. 1. Crop growth rate of coriander as influenced by different date of sowing and planting geometry




Fig. 2. Relative growth rate of coriander as influenced by different date of sowing and planting geometry

from those recorded with the 10th November and 25th November sowing during both seasons. Sowing of coriander on 25th October produced significantly taller plants, higher number of branches plant⁻¹ (Table 1), dry matter plant⁻¹ (Table 2), crop growth rate (Fig. 1) and relative growth rate (Fig. 2) and minimum was observed when coriander was sown on 10th December at all the crop growth stages during both seasons (Table 1 and 2). In early growing season, i.e. 25th October sowing, coriander resulted in to healthy crop growth. The crop sown on 25th October enjoyed favourable environment particularly optimum cardinal temperatures might have helped in better physiological functions of the plant as manifested in increase in plant height, branches per plant increased accumulation of dry

matter per plant. Whereas, the plant did not have optimum opportunity for photosynthesis when it was sown lately. The results are in agreement with Khoja [8], Meena et al. [9], Nath et al. [10] in fennel and Moniruzzaaman et al. [11] in coriander.

Coriander sown with the planting geometry 30 x 5 cm recorded significantly higher plant population and taller plants followed by 30 x 7.5 cm, 30 x 10 cm and 30 x 12.5 cm during both seasons of experimentation. Sowing of coriander with 30 x 12.5 cm spacing, produced significantly higher number of branches plant⁻¹ (Table 1), dry matter plant⁻¹ (Table 2), crop growth rate (Fig. 1) and relative growth rate (Fig. 2). Tallest plants under closer spacing may be attributed to more

competition for light amongst the plants. Increased growth under 30 x 12.5 cm spacing might be attributed to more space plant⁻¹, available under wider spacing. The larger canopy development, associated with profuse branching, most likely increased interception, absorption and utilization of solar energy resulting in formation of higher photosynthates and finally dry matter plant⁻¹, which has resulted in better growth and development of coriander. The results are in line with Kiran et al. [12], Bhapkar et al. [13] and Nethravathi [14].

The plant population and growth characteristics of coriander, such as plant height, number of branches per plant, crop growth rate, and relative growth rate, did not show significant differences due to the combined influence of sowing date and planting arrangement. However, these parameters exhibited numerically superior values when sown on October 25th with a spacing of 30 x 12.5 cm, except plant height which was maximum under 25th October sowing with 30 x 5 cm spacing. Among growth parameters, dry matter accumulation of coriander varied significantly and was recorded significantly maximum under sowing date 25th October along with 30 x 12.5 cm, which was at par with 25th October sowing along with 30 x 10 cm spacing during both the years and on mean basis (Table 3). These results are in conformity with those reported by Meena et al. [15] and Haq et al. [16,17].

4. CONCLUSION

Based on the two-season experiments it has been concluded that the crop sown on 25th October and 30 x 12.5 cm spacing resulted in significantly better growth and development of coriander.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Spices board, India & Ministry of agriculture and farmers welfare, Govt. of India. (ON3019); 2021.
2. Szemplinski W, Nowak J. Nitrogen fertilization versus the yield and quality of coriander fruit (*Coriander sativum* L.). Acta Scientiarum Polonorum Hortorum Cultus. 2015;14(3):37-50.
3. Rangappa M, Bhardwaj HL, Showhda M, Hamama AA. Cilantro (*Coriandrum sativum* L.) response to nitrogen fertilizer rates. Journal of Herbs, Spices and Medicinal Plants. 1997;5(1): 63-68.
4. Gill BS, Randhawa GS, Saini SS. Effect of sowing dates and herb cutting management on growth and yield of fenugreek (*Trigonella foenum graecum*). Indian Journal of Agronomy. 2001;46(2): 364-367.
5. Kuri B, Jat N, Shivran A, Puniya M. Productivity and profitability of coriander varieties influence by sowing time and plant growth regulators. Annals of Agricultural Research, New Series. 2015;36(2):204-211.
6. Rassam GA, Naddaf M, Sefidkan F. Effect of sowing dates and plant density on seed yield and yield components of *Pimpinella anisum*. Res. Sci. J. Iran. Ministry of Agriculture. 2007;20: 127-133.
7. Gomez KA, Gomez AA. Statistical procedure for Agricultural Research., edn 2, John Wiley & Sons, New York. 1984;241-271.
8. Khoja JR. Effect of sowing time and sources of nitrogen on growth, thermal requirement, yield and quality of coriander (*Coriandrum sativum* L.). Ph.D. thesis. Rajashan Agricultural University, Bikaner; 2005.
9. Meena SS, Sen NL, Malhotra SK. Influence of sowing date, nitrogen and plant growth regulators on growth and yield of coriander (*Coriandrum sativum* L.). Journal of Spices and Aromatic crops. 2006;15(2):56-64.
10. Nath P, Jaiswal RC, Verma RB, Yadav GC. Effect of date of sowing, nitrogen levels and spacing on growth and yield of ajwain (*Trachyspermum ammi* L.) Sprague). Journal of Spices and Aromatic Crops. 2008;17(1):01-04.
11. Moniruzzaman M, Rahman MM, Hossain MM, Sirajul KAJM, Khaliq QA. Effect of sowing dates on year-round production of foliage of coriander (*Coriandrum sativum* L.). Bangladesh Journal of Agricultural Research. 2013;38(1):29-39.
12. Kiran MR, Naruka IS, Nayma S, Bepari AR. Effect of sowing time and plant geometry on growth, yield and quality of Black Cumin (*Nigella sativa* L.). International Journal of Current Microbiology and Applied Sciences. 2019;8(5):1915-1921.

13. Bhapkar PB, Sharma HD, Negi S, Pundir D, Sharma V, Kapil M, Reddy S. Effect of cutting and intra row spacing on yield and phenotypical attributes of coriander (*Coriandrum sativum* L.). International Journal of Current Microbiology and Applied Sciences. 2019;8(9):693-698.
14. Nethravathi R. Influence of planting geometry and fertilizer levels on the seed yield and quality of coriander (*Coriandrum sativum* L.). M.Sc. (Ag.) thesis; 2019.
15. Meena SS, Mehta RS, Lal G, Sharma YK, Meena RD, Kant K. Effect of sowing time and crop geometry on growth and seed yield of dill (*Anethum sowa* L.) International Journal of Seed Spices. 2015;5 (1): 79-82.
16. Haq MZ, Hossain MM, Haque MM, Das MR, Huda MS. Blossoming characteristics in black cumin genotypes in relation seed yield influenced by sowing time. American Journal of Plant Sciences. 2015;6(8): 1167-1183.
17. Kiran MR. Effect of sowing time and plant geometry on growth, yield and quality of black cumin (*Nigella sativa* L.) under Malwa plateau condition. M.Sc (Ag.) thesis; 2018.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113760>