



A Review on New Approaches in Dragon Fruit Production, Nutraceutical Insights and Morphological Dynamics

Sangeeta Lakshmeshwara ^{a++}, Shubham Jain ^{b++*},
S. Manasa ^c, Anshuman Singh ^{d#}, Alemla Imchen ^{e++},
Pasupuleti Venkata Siva Prasad ^{f†}, Osheen Raizada ^{g++}
and Vikas Yadav ^{h++}

^a Department of Fruit Science, University of Horticultural Sciences, Bagalkot, Karnataka, 587101, India.

^b Department of Fruit Science, ANDUAT, Ayodhya, India.

^c Krishi Vigyan Kendra, Near RTO Office, Rayachoty Road, Utukur, Kadapa, YSR District, Andhra Pradesh, 516003, India.

^d Department of Horticulture, KNIPSS Sultanpur, 228118, India.

^e Department of Horticulture, School of Agricultural Sciences (SAS), Nagaland University, Medziphema Campus, Nagaland, India.

^f Department of Fruit Science, College of Horticulture, Anantharajupeta, Dr Ysr Horticultural University, India.

^g Department of Fruit Science, Mahatma Gandhi University of Horticulture and Forestry, Sankra-Durg (C.G), India.

^h Department of Soil Science and Agricultural Chemistry, ANDUAT, Ayodhya, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2024/v27i5847

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116137>

Review Article

Received: 20/02/2024

Accepted: 24/04/2024

Published: 27/04/2024

⁺⁺ Ph.D Scholar;

[#] Assistant Professor;

[†] M.sc Horticulture;

*Corresponding author: E-mail: shubhu15296@gmail.com;

ABSTRACT

Dragon fruit (*Hylocereus spp.*) is recently introduced amazing fruit in India, its cultivation gaining popularity because of its high nutritional value and antioxidant property. Recently, it drawn the much attention of the Indian consumers owing to its pleasant flavour, colour, with their attractive appearance, tremendous health and nutritional benefits. Dragon fruit extracts from stems, flowers, peels and pulps have beneficial biological properties against pathogenic microbes like bacteria, fungi and viruses as well as diseases like diabetes, obesity, hyperlipidemia and cancer. Now a days, Indian growers are much interested in its cultivation due to its promising yield, easy cultivation, great commercial potential, which increases the economical status of the farmers. The area under cultivation is insufficient to fulfill the requirement of peoples hence, there is a need to increase dragon fruit production in India. At present, a very few information is available on production aspects of dragon fruit this review papers is dealt with the knowledge regarding the botany, nutritional composition, uses, health benefits, species, cultivation aspects of dragon fruit based on recent updates of some of the research findings.

Keywords: Dragon fruit; nutraceutical properties; botanical description; cultivation aspects.

1. INTRODUCTION

“Dragon fruit is a herbaceous perennial climbing cactus species belongs to the family cactaceae” (Jussieu [1], Patwary et al., [2]. “It is originated in tropical and sub-tropical forest regions of Mexico and South America” [3]. In early 1800, French brought dragon fruit from Guana to Vietnam as a ornamental plant but presently in Vietnam this dragon fruit become one of the foremost important commercial fruit crop [4]. In India it is introduced in the late 90’s as garden crop but currently in India, the area under dragon fruit cultivation also increasing gradually as commercial fruit crop. In the world, it is widely distributed in Vietnam, China, Mexico, Colombia, Thailand, Malaysia, Indonesia, Australia and United states (Mizrahi and Nerd, [3], Nobel and D la Barrera, [5]. In Indian states, the ideal fruit cultivation started at Karnataka, Tamil Nadu, Kerala, Maharastra, Gujarat, Orissa, Andra Pradesh, West Bengal and Andaman & Nicobar Island etc. The majority of dragon fruits in Indian markets are imported from Vietnam, Thailand, Malaysia and Sri Lanka [6]. “The dragon fruitcultivation is expanding in recent years due to its health and economic importance leading to its utilization as a source of functional materials to provide phytochemicals with strong antioxidant capability. The fruits are rich in vitamins, fiber, phosphorus, calcium, magnesium, antioxidan and phytochemicals. Dragon fruits become popular in Asian countries for its nutritional values, attractive feature and color” (Harivaindaran et al., [7], Hoa et al., [8]. Dragon fruit is recently introduced fruit crop in India, it has tremendous scope for its cultivation because of it's high nutritional property and returns per unit

area. At present, less information is available on cultivation aspects of dragon fruit. Research on different aspects of cultivation and health benefits of this fruit can help to maximize the benefits to worldwide growers and consumers and to expand the market of Dragon fruit.

2. COMPOSITION AND USES

Dragon fruit nutritional composition is varies with species to species of *Hylocer spp*, it is the abundant source of vitamins and mineral nutrients (Nurliyana et al. [9], Perween et al. 2018) which includes the vitamin A, vitamin B and richest source of vitamin C, carbohydrates, betacyanin's, antioxidants, flavonoids, carotenes, polyphenols, iron and Phyto albumins [10]. The 100 g fresh fruits biochemical analysis contain 83-88% moisture, proteins (0.5-0.16 g), fat (0.1-0.61g), calcium (6-8.8mg), phosphorus (19-36.1 mg), iron (0.4-0.6 mg), ascorbic acid (7-25 mg), poly phenols (7.23-1 μ g) and total flavonoids (26.8-46.29 mg). The fruit's titratable acidity ranged from 0.20 to 0.30 mg lactic acid, making it somewhat acidic [11]. Dragon fruit is recognized as the one of the tropical super fruit because of its abundant nutrients, has lot of various nutrients and less in calories. Reports suggested that it helps in reducing the diabetes and hypertension (Kumar et al., 2018), act as anti-cancerous as it is rich in antioxidant, it fights against free radicals causing cancer (Yusof et al., [12], anti-inflammatory by reducing the arthritis pain, the seeds are rich in polyunsaturated fats (omega-3 and omega-6 fatty acids), which lower triglycerides and reduce the risk of cardiovascular disorders Sonawane, [4]. It boosts the immune system by increasing platelet count

and lowering cholesterol levels and high in fiber, which aid constipation relief and contains low calories.

"Dragon fruit peel has high potential as a natural dye" [7]. "The red layer of the fruit is rich in vitamins like B1, B2, B3, C and minerals" [13]. "The higher ascorbic acid content was observed in premature stem than the dragon fruit flesh, which may help to prevent the risk of certain disease such as scurvy, weakness and anaemia" [14]. "The pulp and peel extract of fruit have phytochemical compounds, which have antimicrobial activity and can be used as a natural antioxidant" [15].

"The fruits can be eaten as raw or processed for ice cream, candies, cookies, shake, jam, jelly, wines, for special beverages or as flavor for all kinds of drinks and ingredients of various salad recipes. The dragon fruits flower have been cooked as soups, ingredient of filipino viands and lumpia. The pulp of skin is also processed as pickles, jam and be boiled as cleansing drinks. The pulp of skin and skin can also used for beauty soap preparation" [16]. Dragon fruit peel has a high potential to be used as a natural dye Harivaidaran et al., [7] the flesh of the pulp is juicy and contains many small black seeds. "It is also regarded as a possible source of antioxidants and micronutrients". (Ariffin et al., [17], Jaafar et al., [14], Lim et al., 2010)

3. AREA AND PRODUCTION

According to the recent data, the maximum area and production covered under Vietnam followed by China, Indonesia, Taiwan, Thailand, Malaysia and Phillipines etc [11] The maximum productivity is reported in Vietnam followed by Indonesia, China, Taiwan, Cambodia and India etc. Vietnam is the largest dragon fruit producer in the world, accounting the highest share in Asia, Europe and sometimes in the United States. The biggest importer and consumer of dragon fruit in this world is China and Vietnam, the main exporter of dragon fruit worldwide due to high global demand (Ratnala Thulaja and Abd Rahman, 2017). In India, Gujarat have the highest area and production followed by Karnataka, Maharashtra, West Bengal, Andhra Pradesh, Tamilnadu and the highest productivity is recorded in Maharashtra.

4. MORPHOLOGY AND PHYSIOLOGY OF DRAGON FRUIT

Dragon fruit belongs to cactus species, it is perennial, herbaceous, climbing vine. It is a CAM (Crassulacean acid metabolism) plant, during day time stomata closed which helps in reducing transpirational loss. It is a terrestrial or epiphytic cactus with succulent three-winged and green stems (Patel and Ishnava, 2019). It has thick, fleshy three angled stem and wax coating, aerial roots which helps the plant to climb and also helps in absorbing moisture from surrounding, areoles in stem from which spines arises, shallow rooted crop. It has beautiful hermaphrodite flower (length 25-30cm), which is white in colour, large and attractive in nature, flowering occurs in April-May month (Merten, [18]. They are scented and blooms only at night and last only one night. Young flower buds are edible in nature, which used as salad in Malaysian countries. It is cross pollinated crop because, stigma is present over the stamen which hinders self pollination, it is ornithophilous in nature, pollination is carried out by bats, hawk moths and honey bees. Flower starts open during evening (6:30 to 7 pm) it takes around 20 to 25 days from appearance of flower bud to full blossom (Karunakaran and M. Arivalagan, [19]. The fruits are spherical to oval in shape with bright red or yellow skin and green foliaceous scales, spineless with several broad leafy bracts and are larger resembling the skin of a dragon [8]. Seeds are black, small, elongate and kidney shaped. Britton and Rose, [20].

4.1 Four types of Dragon Fruits are Mainly Grown in India are

Hylocerus undatus: Red skin having white flesh fruits [20]

Hylocerus polychrous: Red skin with pink flesh fruits [20]

Hylocerus costaricensis : Red skin with purple flesh fruits [21]

Selenicereus megalanthus : Yellow skin with white flesh fruits [22]

Among these species, *Hylocerus spp* are diploid in and *Selenicereus app* is tetraploid in nature, the white and pink flesh fruits are widely cultivated types in India (Fig 1).



Fig. 1. Species of dragon fruit grown in India Wakchaure et al., 2020




Fig. 2. Time taken for flower bud to flower formation [23]

4.2 Dragon Fruits Fruiting Stages

Fruiting starts from June to November, it comes in 5-6 flushes, takes around 30 to 40 days for fruit set to maturity. Fruit is botanically known as berry, where endocarp and mascara are edible part and non-climacteric fruit.

By using a 3-digit numerical system dragon fruit phenological stages are classified based on the extended BBCH (Biologische Bundesanstalt, Bundessortenamt and Chemische Industrie)

scale, which assists to the standardization of its phenological stages (Fig 2). Bud development (0), shoot development (1), vegetatively propagated organ development (4), reproductive development (5), flowering (6), fruit development (7) and fruit maturation (8) are the seven major growth stages.

4.3 Soil and Climate

It is a hardy crop, it can be grow in wild range of soils but gives the best results when they grow

in well drained, sandy loamy soils which is rich in organic matter having pH of 5.5 to 6.5 is good for its cultivation. Water logged condition leads to rotting of root and to avoid this problem, soil may be mixed with tiny stone gravels or sand to provide the good drainage facility [11].

Dragon fruit is a long day and photosensitive crop (Morton 1987), that requires adequate sunlight and a dry tropical climate, while it is adaptable to humid, semi-arid, tropical and subtropical conditions. Even though sun-loving plant, it prefers temperatures between 20^oC and 29^oC. It can tolerate temperatures has high as 38^oC and low as 0^oC for short periods of time. Above 40^oC temperature will cause stems yellowing in the plants. It requires 500 – 1500 mm annual rainfall and can be grown at an elevation of about 1700 m above MSL. Excessive rain fall causes flower and fruit drop (Karunakaran and Arivalagan, [19].

4.4 Cultivars and Propagation

In India, currently there are no officially released varieties or clones. The development of dragon fruit varies and hybrids is currently progressing at ICAR-NIASM in Baramati. Almost all the varieties/clones being propagated and cultivated by farmers and nurseries are introduced from other countries and those remultiplied in India. The mainly grown cultivar of dragon fruits are *Hylocerus undatus*, *Hylocerus polyrhizus* and *Hylocerus costaricensis*. Among these types of dragon fruits, red fleshed red skin was popularized and high in area and production because it is particularly rich in betalains, which have the high antioxidant property and used as a natural food colorants and for its wider adaptability. Exotic varieties are Alice, American beauty, Bloody mary, Cosmic charlie, Costarican Sunset, Dark star, Delight, Thompson, Red Jaina, Zomorano etc are reported in other countries.

Sexual means (seeds) of propagation is very rare in dragon fruit because of the fact that even after a year, seedlings raised from seeds in dragon fruit remain to be smaller and have thin stem and takes long time to yield, less vigorous and are not true to its mother. Stem cuttings are the commercial method of propagation. The cutting should be prepared a couple of days before planting by removing the latex oozing from cutting by drying. The healthy and vigorous cutting are taken from mother plants after the fruiting season and are produce flowers within three years [24]. To prevent the diseases the cutting should be treated with fungicides. The better results were

observed, when the cuttings were taken from the middle portion of the main stem about 15-25 cm and gives the better shooting and rooting. Planting the cuttings in 12 x 30 cm size polyethylene bags, which is filled with mixture of soil, farmyard FYM and sand in the ratio of 1:1:2. The bags should be placed in a shady area to encourage better rooting. While too much moisture should be avoided to prevent cuttings from rotting. After 5-6 months, these cuttings develop strong roots and ready for planting [25]. Notably high survival percentage was observed in Nov, Dec and January [26].

It is commercially propagated by cuttings, but multiplication rates are extremely low, and obtaining sufficient planting material is difficult due to the large size of the cuttings required.

So, *in vitro* propagation was found to yield more planting material within short period of time. Explants should be selected based on their age and place of origin, whether they are from *in vitro* germinated seedlings or *in vivo* mature plants [27]. The *in vitro* response was observed using shoot tips 2cm in length supplemented with 2mgL⁻¹ of BAP. MS medium supplemented with 7mg L⁻¹ BAP and 1 mgL⁻¹ Kinetin were recorded the best combination in terms of time taken for multiple shoot induction, shoot length and number of multiple shoots [28]. Stem and leaf explants, regenerated the highest number of shoots on MS medium supplemented with 2.5mg/l BA and 0.01 NAA mg/l compare to that of the other hormone combinations. Rooting was observed in regenerated mature shoots after transferred onto MS basal medium with 0.01mg/l NAA.

4.5 Field Preparations and Planting

The field should be adequately cleaned by removing any trees, shrubs and their stubbles to allow for unrestricted movement of man and machines. A virgin land needs deep ploughing and harrowing. Plough the field until the soil achieves its fine tilth and is weed-free. After that, you should apply any type of organic compost in a proportionate ratio. Dragon fruit cultivation thrives well in areas with abundant sunlight and open spaces, making them ideal for planting, shaded regions are unsuitable for cultivating dragon fruit. Planting season for dragon fruit cuttings is rainy season i.e June – July. There are different planting systems like single pole system, continuous pyramid system. Generally single pole system consists of

concrete pole with ring structure is followed with spacing 3 x 3 m or 4 x 4 m, the vertical height of the pole ranges from 1.5 to 2 meters, allowing branching and hanging of the vines. By placing the dragon fruit trees next to the poles, facilitates the easy climbing. Per pole four plants are recommended, 300 to 400 poles and 1200 -1600 plants are required to establish 1 acre dragon fruit orchard (Wakchaure et al., 2021). To encourage the controlled growth, it is advised to limit lateral shoots, allowing 2-3 main stem to develop, with lateral removal of shoots as needed. The developed scale act as a useful tool for adoption of effective crop management practices such as pollination, nutrient management, plant propagation, timely harvest of fruits and pest management. Kundan Kishore, [23].

4.6 Irrigation and Nutrient Management

Even pithaya can survive with very low rainfall, it requires less amount of water (120 to 150 mm of irrigation/year). For good quality of fruit, there is a need of regular water supply to the plants because, it enables the plant to build sufficient reserves not only to flower at the most favourable time but also to ensure the development of the fruits. Drip irrigation method is adapted to avoid uneven and excess watering that can result in the flowers and the young fruits falling off. Critical stages of irrigation are flowering and fruit enlargement stages. Like other fruit crops dragon fruit also needs judicious application of nutrients for 1 to 2 year old plants need 10 to 15 Kg FYM with 300:200:200 g NPK/pillar. More than 3 year old plants need 540:720:300 g NPK/pillar applied in four split applications (April, July, October and January). The application of soluble chemical fertilizers at 500: 750: 300 g NPK/pillar with 4 split applications (in April, July, October and January) through fertigation, improved yield and yield attributing characters of dragon fruit. Thanki et al., [29].

4.7 Intercultural Operations

In the initial stages, plants grow into thick, dense vines that spread quickly. To grow in the direction of the stands, the lateral buds and branches must be pruned. Once the vines have reached the top of the stands, branches are allowed to grow. The tip of the main stem is removed to allow new shoots to grow laterally and climb at the ring that forms an umbrella-like structure of vines. After harvest, the plants are pruned to promote the growth of new, flower-bearing shoots.

4.8 Training and Pruning

The Indian institute of horticulture research, Bengaluru has developed the 4 different trellis system viz., (Wakchaure et al., 2021)

- Single pole with cement ring
- Single pole iron ring
- 'T' stands
- 'T' stands continuous pyramid.

5. SINGLE POLE WITH CEMENT AND IRON RING

Each trellis is made up of 6 feet tall height by 5-6 inch thick pole constructed at a depth of 2 feet. For support, single cement poles and rings are used. Rubber tires cut across are placed on top of poles in some regions. Plants can be planted near the poles to allow them to climb easily. 2-4 plants per pole, depending on climate condition. In comparison to others, this approach is less expensive. In India, this is the preferred method.

5.1 'T' Stands

'T' shape structure of the stand, Plants are planted between the two poles. Initial investment is high, around 6-6.5 lac/acre. After 4th year become clouded and chances of insect pest attack. Usually not prefer by the farmers.

5.2 'T' stands Continuous Pyramid Stands

Continuous type G.I. pipe stands and G.I. angles can also be used. This pyramid shape has a length of about 10-15 meters. A greater number of plants could be planted. Adequate aeration was provided to avoid disease and pests attack. Plants should be placed 2-3 feet apart on both sides of the structure. The distance between two structures could be between 5 and 6 meters. Compared to other trellis systems, the single pole system performed better in terms of growth and yield. Although cement poles are expensive, they are long-lasting and can be used.

5.3 Harvesting

Dragon fruits are harvested between 31 and 36 days after flowering and best for ideal maturity and quality. The two species of dragon fruit red and white pulp fruits harvested at 31 day after flowering showed better physicochemical and sensory quality. Harvesting is done when the fruit epicarp colour changes to red from green



Fig. 3. Different trellis systems namely (a) concrete pole and rings (b) T stand and iron wires (C) Wooden ladder for dragon fruit establishment

colour. Optimum time for harvesting is 7 days after colour transition. Flowering and fruit setting time significantly affect the quality of fruits, especially on total soluble solids contents [30]. Mature Dragon fruits have higher TSS, which is mainly higher in autumn fruits than in summer fruits [31]. Dragon fruit is non-climacteric and good for consumption, when it is ripe however, its quality deteriorates gradually after harvest and during storage (Nerd et al., [32], Chien et al., [33].

5.4 Yield and Storage

Dragon fruit's stem cuttings starts to yield after 9 months of planting, the economical yield starts after 2 years of planting, fruit starts producing in the month of July till August depends on the species and varieties. It fruits in 3-4 waves during one season. Each pole plants approximately produce 40-100 fruits per wave. Each fruit weight about 300-1000gm and 15-25 kg yield was recorded in each pole. It produces the 8-15 ton per acre of plant. The life span of dragon fruit is about 15-20 years. The fruits that are stored at 10°C and 93 percent RH had a longer storage life of 15 to 17 days. In normal room conditions, the post-storage life is 3 to 4 days. The average weight loss is 0.3 percent per day inside the cold room for 17 days storage and 2.6 percent per day in the regular room for 4 days after storage. After 3.5 days, fruits stored under normal room conditions exhibited 100 percent decay incidence and severity [34-36].

6. PHYSIOLOGICAL DISORDERS

6.1 Sun Burn Injury

Sun burn injuries have become prevalent in many places of India. In the month of March and April sun burn symptoms are appeared, when the

day and night temperatures varied greatly, particularly in the region where summer temperatures exceeded 40° C [37-39].

6.2 Diseases

In general, dragon fruit is tolerant to the major diseases and pests found in India. Anthracnose, brown spots and stem rots are the major diseases affecting the dragon fruit crop in India. Heavy rains, overwatering, or waterlogged circumstances make the crop prone to certain diseases.

7. ANTHRACNOSE

7.1 Causal Organism: *Colletotrichum siamense*

Symptoms: concentric lesions of reddish or orangish brown with ascervuli begins at the locations where the vine's spines extend from the edge, close to the ribs. It can be controlled by spraying with chlorothalonil 2gm per liter, mancozeb at 2gm per liter and carbendazim at 1gm per liter of water.

Stem canker (*Neoscytalidium dimidiatum*): Stem canker, the most destructive disease of dragon fruit, has been observed in several orchards with varying degrees of severity. Unmanaged orchards were severely affected by the disease.

Symptoms: The affected cladodes initially exhibit depressed, chlorotic spots with red flecks. As the disease advances, the spot centers rise and turn red to grey. The centers of spots expand and harden into brown scabs with

pycnidia embedded on the surface. As the disease progresses, affected cladodes begin to yellow and then rot. Over time, the necrotic tissues separate from the healthy tissues, leaving shot holes [40-42]. On fruits, chlorotic depressed spots are followed by rotting of the affected fruit (Kakade et., 2022)

Wil: *Fusarium* species. Symptoms: Drying of plant tissues

8. CONCLUSION

Dragon fruit has the great nutritional and health benefits, present market trade demand showed the importance of this fruit is more than the production in India. To meet this demand, there is a need to year round production of dragon fruit. There is a need to standardize the package of practice for cultivation. It is necessary to develop a training module on the set of methods for successful dragon fruit farming and business ventures. This crop as commercial crop for enhancing income of marginal and small scale farmers in dryland areas. Requirement of post harvest management practices is necessary to avoid the post harvest losses. Lack of information on the species and varieties availability in India there is essentiality of development of variety through conventional and biotechnology tools.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Jussieu de AL. Genera plantarum. Parisiis. 1789;15:312–317.
2. Patwary MMA, Rahman MH, Barua H, Sarkar S, Alam MS. Study on the growth and development of two dragon fruit (*Hylocereus undatus*) genotypes. The Agriculturist. 2013;11(2):52-57.
3. Mizrahi Y, Nerd A. Climbing and columnar cacti: New arid land fruit crops. In: Janick, J. (ed) Perspective on new crops and new uses. ASHS Press, American Society of Horticultural Science, Alexandria, Virginia. 1999;358- 366.
4. Sonawane MS. Nutritive and medicinal value of dragon fruit. The Asian Journal of Horticulture. 2017;12(2):267-271.
5. Nobel PS, De la Barrera E. Stem water relations and wet CO₂ uptake for a hemiepiphytic cactus during short term drought. Environmental and Experimental Botany. 2002;48:129-137.
6. Ahmad HB, Mohd MH, Nur SJ. Status and challenges of dragon fruit production in Malaysia. FFTC Agricultural Policy Platform (FFTC-AP). 2019;1-8.
7. Harivaindaran KV, Rebecca OPS, Chandran S. Study of optimal temperature, pH and stability of Dragon fruit (*Hylocereus polyrhizus*) peel for use as potential natural colorant. Pakistan Journal of Biological Sciences. 2008;11(18):2259-2263.
8. Hoa TT, Clark CJ, Waddell BC, Woolf AB. Postharvest quality of dragon fruit (*Hylocereus undatus*) following disinfecting hot air treatments. Postharvest Biology and Technology. 2006;41(1):62-69.
9. Nurliyana RD, Syed Zahir I, Mustapha Suleiman K, Aisyah MR, Kamarul Rahim K. Antioxidant study of pulps and peels of dragon fruits: A comparative study. International Food Research Journal. 2010;17:367–375.
10. Moshfeghi N, Mahdavi O, Shahhosseini F, Malekifar S, Taghizadeh SK. Introducing a new natural product from Dragon fruit into the market. International Journal of Research and Reviews in Applied Sciences. 2013;15(2):269-272.
11. Wakchaure GC, Satish K, Meena KK, Rane J, Pathak H. Dragon fruit cultivation in India: Scope, Marketing, Constraints and Policy Issues. Technical Bulletin No. 46. ICAR-National Institute of Abiotic Stress Management, Baramati, Pune, Maharashtra, India. 2021;54-1
12. Yusof YA, Salleh FSM, Chin NL, Talib RA. The drying and tabletting of pitaya powder. Journal of Food Process Engineering. 2012;35:763-771.
13. Le Bellec F, Vaillant F, Imbert E, Pitahaya (*Hylocereus spp.*): A new fruit crop, a market with a future. Fruits. 2006;61(04):237-250.
14. Jaafar RA, Rahman ARBA, Mahmud NZC, Vasudevan R. Proximate analysis of Dragon fruit (*Hylocereus polyrhizus*). American Journal of Applied Sciences. 2009;6(7):1341-1346.
15. Patel SK, Ishnava KB. *In-vitro* antioxidant and antimicrobial activity of fruit pulp and peel of *Hylocereus undatus* (Haworth) Britton and Rose. Asian Journal of

- Ethnopharmacology and Medicinal Foods. 2019;5(2):30-34.
16. Pascua LT, Pascua ME, Gabriel MLS. Dragon fruit production and marketing in the Philippines: Its Status, Constraints and Prospects. In: Jiang Y.L., P.C. Liu, P.H. Huang (eds). Improving Pitaya Production and Marketing. Food and Fertilizer Technology Center. Taipei, Taiwan. 2015;47-65.
 17. Ariffin AA, Bakar J, Tan CP, Rahman RA, Karim R, Loi CC. Essential fatty acids of Pitaya (Dragon fruit) seed oil. Food Chemistry. 2009;114:561-564.
 18. Merten S. A review of *Hylocereus* production in the United States. Journal of the Professional Association for Cactus Development. 2003;5:98- 105.
 19. Karunakaran G, Arivalagan M. Dragon Fruit - A New Introduction Crop with Promising market. Indian Horticulture. 2019;63(1):8-11.
 20. Britton NL, Rose JN. The cactaceae. Descriptions and Illustrations of Plants of the Cactus Family. USA, Carnegie Institution of Washington. 1920;2:183–195.
 21. Britton NL, Rose JN. The genus *Cereus* and its allies in North America. Contributions from the United States National Herbarium. 1909;12:413– 437.
Available:<http://www.jstor.org/stable/23491827> (accessed Feb 3, 2021).
 22. Moran RV. *Selenicereus megalanthus* (Schumann) Moran. Gentes Herbarum. 1953;8:325.
 23. Kundan Kishore. Phenological growth stages of dragon fruit (*Hylocereus undatus*) according to the extended BBCH-scale. Sci Hort. 2016;213:294–302.
 24. Rao CC, Sasanka VM. Dragon fruit 'The Wondrous Fruit' for the 21st century. Global Journal for Research Analysis. 2015;4(10):261-262.
 25. Tripathi PC, Karunakaran G, Sankar V, Senthil RK. Dragon fruit: Nutritive and Ruminative Fruit, Technical Bulletin No. 11/2014, pp1-9. Indian Institute of Horticultural Research, Bengaluru, India; 2014.
 26. Nandi P, Tarai RK, Ghosh SN. Study on rooting behaviour of different types of cutting of dragon fruit at different period of year. Int. J. Minor Fruits, Medicinal and Aromatic Plants. 2019;5(2): 45-49.
 27. Garcia-Rubio O, Malda-Barrera G. Micropropagation and reintroduction of the endemic mammillaria mathildae (Cactaceae) to its natural habitat. Hort. Sci. 2010;45(6):934-938.
 28. Arunkumar G, Kavino M, Auxilia J, Hemaprabha K. Efficient *In vitro* Propagation Protocol for Mass Multiplication in Dragon Fruit (*Hylocereus costaricensis*). Biological Forum. 2022;14(2a):20-25.
 29. Thanki DM, Kalyanjibhai DP, Suraj K. Influence of fertigation schedule on yield of dragon fruit (*Hylocereus polyrhizus* Britton & Rose), The Pharma Innovation Journal. 2022;11(9):518-521.
 30. Mallik B, Hossain M, Rahim A. Influences of variety and flowering time on some physiomorphological and chemical traits of Dragon fruit (*Hylocereus spp.*). Journal of Horticulture and Postharvest Research. 2018;1(2):115-130.
 31. Nomura K, Ide M, Yonemoto Y. Changes in sugars and acids in pitaya (*Hylocereus undatus*) fruit during development. The Journal of Horticultural Science and Biotechnology. 2005;80(6):711-715.
 32. Nerd A, Gutman F, Mizrahi Y. Ripening and postharvest behaviour of fruits of two *Hylocereus* species (Cactaceae). Postharvest Biology and Technology. 1999;17(1):39-45.
 33. Chien PJ, Sheu F, Lin HR. Quality assessment of low molecular weight chitosan coating on sliced red pitayas. Journal of Food Engineering. 2007;79(2):736-740.
 34. Jadhav PB. Extending the storage and Post-Storage life of dragon fruit using a cold room (Ecofrost). International Journal of Agriculture, Environment and Biotechnology. 2018;11(3):573-577.
 35. Britton NL. Flora of Bermuda. Charles Scribner's Sons, New York: 256; 1918.
Available:<https://www.biodiversitylibrary.org/item/16209#page/276/mode/1up> (accessed Feb 3, 2021).
 36. Dahanayake N, Ranawake AL. Regeneration of dragon fruit (*Hylocereus undatus*) plant-lets from leaf and stem explants. Tropical Agricultural Research and Extension. 2011;14(4).
 37. Kakade VD, Boraiah KM, Salunkhe S, Nangare DD, Sangram B Chavan, Wakchaure GC, Jadhav SD, Rajkumar, Ravi Kumar K, Taware PB, Tayade,

- Sammi KR. Technical Bulletin No. 56. ICAR-National Institute of Abiotic Stress Management, Baramati, Pune, Maharashtra, India; 2023.
38. Lata D, Narayana CK, Karunakaran G, Sudhakar Rao DV, Anuradha S. Fruit Maturity determination of red and white pulp dragon. *J. Hort. Sci.* 2022;17(1):157-165
39. Lim HK, CP, Tan CP, Bakar J, Ng SP. Effects of different wall materials on the physicochemical properties and oxidative stability of spray-dried microencapsulated red fleshed pitaya (*Hylocereus polyrhizus*) seed oil. *Food Bioprocess Technology.* 2012;5:1220-1227.
40. Mizrahi Y, Nerd A, Nobel PS. Cacti as crops. *Horticultural Review.* 1997;18:291-320.
41. Morton JF. Strawberry Pear. In: Morton J.F. (ed.): *Fruits of Warm Climates.* Miami, FL, USA, Morton. 1987;347–348.
42. Thulaja RN, Abd Rahman NA. Dragon Fruit. Singapore Infopedia; 2017. Available:https://eresources.nlb.gov.sg/infopedia/articles/sip_768_2005-01-11.html (Accessed mar 20, 2020).

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116137>