



Molluscicidal Activity of Aqueous Extracts of Selected Medicinal Plants on Fresh Water Snails and Fish

Reshmina Firoz Khan ^a, Ram P. Yadav ^{a*} and Ajay Singh ^a

^a Natural Product Laboratory, Department of Zoology, D.D.U Gorakhpur University, Gorakhpur (U.P), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/APRJ/2022/v10i4199

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95594>

Original Research Article

Received: 20/10/2022

Accepted: 29/12/2022

Published: 30/12/2022

ABSTRACT

Aim: To investigate the molluscicidal activity of aqueous extracts of *Phyllanthus niruri* and *Euphorbia tirucalli* on fresh water snails *Lymnaea (Radix) acuminata* and *Indoplanorbis exustus* (as target organisms) and on *Channa punctatus* fish (as non target organism), sharing the same habitat.

Methodology: The aqueous extracts of the stem bark and leaves of *Phyllanthus niruri* (a tropical small herb, commonly known as a Bhumi Amla), belonging to the family Phyllanthaceae and the aqueous extracts of stem bark and latex of *Euphorbia tirucalli* (a semi-arid tropical plant, commonly called milk bush) of the family Euphorbiaceae, were investigated as plant origin molluscicides, against freshwater snails, responsible for causing fascioliasis, *Lymnaea (Radix) acuminata* and *Indoplanorbis exustus*, as well as on *Channa punctatus*, using Singh and Agarwal method and POLO computer program.

Results: A significant time and dose-dependent effect of aqueous extracts of both the plants against the freshwater snails were observed. Thus increase in exposure time, the LC₅₀ of *Phyllanthus niruri* stem and leaf extracts were decreased from 267.89 mg DW/L (24h) > to

*Corresponding author: E-mail: rampratapy123@gmail.com, reshminakhan0891@gmail.com;

180.64mg DW/L (96h) against *Lymnaea acuminata* and 261.49 mg DW/L (24h) to 136.40 mg DW/L (96h) against *Indoplanorbis exustus*. Likewise, the LC₅₀ of aqueous extract of latex of *Euphorbia tirucalli* were decreased from 1.80 mg DW/L (24h) to 0.65 mg DW/L (96h) against *Lymnaea acuminata* and 0.90 mg DW/L (24h) to 0.28 mg DW/L (96h) against *Indoplanorbis exustus*, with the increase in exposure time. Its aqueous stem bark extract showed a similar pattern. These aqueous plant extracts at higher doses were also observed to be lethal to freshwater fish *Channa punctatus*, which share a common habitat with the freshwater snails, but the doses LC₉₀, (24h) of snails didn't cause any mortality to these fishes, thus indicating that these plant extracts can be safely used as molluscicides.

Keywords: Molluscicides; Euphorbiaceae; *Lymnaea acuminata*; *Indoplanorbis exustus*; *Channa punctatus*; Phyllanthaceae.

1. INTRODUCTION

The World Health Organization considers *Fascioliasis* and *Schistosomiasis* to be neglected zoonotic diseases. Recent research has validated fascioliasis inclusion on the list of major human parasitic illnesses. Fascioliasis is currently a vector-borne illness with the most extensive known latitudinal, longitudinal, and altitudinal distribution. Despite theoretical limitations connected to its biology, *Fasciola hepatica* has colonized five continents, despite being dependent on environmental and human activity. In hypo- to hyper endemic areas of Central and South America, Europe, Africa, and Asia, human fascioliasis displays a diversity of epidemiological characteristics that are linked to a wide range of environments. The presence of adequate aquatic snail intermediate hosts from the Lymnaeidae i.e. *Lymnaea acuminata* and Planorbidae *Indoplanorbis exustus* families is a significant aspect.

Over the last few decades, a variety of chemical molluscicides have been employed to treat Fascioliasis and other snail-borne infections. Copper sulphate and other copper salts are among the most notable, having previously served important roles but now being mostly abandoned due to low efficiency and inactivation by organic and inorganic materials in water. Despite the many advantages of utilizing synthetic molluscicides as mollusc control agents, they have the potential to disrupt entire ecosystems, inflicting harm to humans and non-target creatures. Due to their bioaccumulation and long-term persistence, synthetic pesticides have generated a high rate of toxicity in water body's organism. Plant molluscicides are receiving more attention from national and international agencies in the hopes of proving to be less expensive and more readily available than synthetic pesticides. Plant molluscicide research has become multidisciplinary as a result

plant products have been promoted in several countries due to their wide variety of desirable properties, including high toxicity, low cost, low mammalian toxicity, low water solubility, easy biodegradability, plentiful growth in endemic areas, and operator safety.

Thevetia peruviana, *Alstonia scholaris* (Family: Apocynaceae), *Euphorbia pulcherima*, and *Euphorbia hirta* (Family: Euphorbiaceae) are examples of common therapeutic plants with high molluscicidal action against freshwater snails. *Alstonia scholaris* has a variety of alkaloids (including pseudoakumigine, ursolic acid, and sitosterol), steroids, and triterpenoids, whereas *Euphorbia hirta* and *Euphorbia pulcherima* have the diterpenoids pulcherrol, sitosterol, hentriacontane, ellagic acid, and amyrin" [1]. "*Euphorbia hirta* latex has been found to be effective against the freshwater snails *Lymnaea acuminata* and *Indoplanorbis exustus*" [2].

The present study reports the molluscicidal effects of aqueous extracts of different parts of two plants *Phyllanthus niruri* and *Euphorbia tirucalli* against harmful freshwater snails, *Lymnaea acuminata* and *Indoplanorbis exustus*, which serve as intermediate hosts for *Fasciola hepatica* and *Fasciola gigantica*, in causing fascioliasis. Toxicity tests were also carried out on freshwater fish *Channa punctatus* (which shares common habitat with the snails) to see if there was any environmental toxicity.

2. MATERIALS AND METHODS

2.1 Test Plants

The plants under investigation were *Phyllanthus niruri* and *Euphorbia tirucalli*. Both the plants were collected easily, during the rainy season, from the Botanical Garden of Deen Dayal Upadhyay Gorakhpur University, Gorakhpur

(U.P) INDIA. *Phyllanthus niruri* (commonly called bhumi amla), belongs to the family Phyllanthaceae, while *Euphorbia tirucalli* (commonly called milk bush or pencil tree) belongs to the family Euphorbiaceae, the milky latex of which is extremely irritating to the skin and is toxic and therefore should be carefully handled.

2.2 Preparation of Aqueous Extracts of Stem Bark, and Latex

2.2.1 Stem bark

Fresh stem and leaves from *Phyllanthus niruri* and stem bark from *Euphorbia tirucalli* were minced with distilled water, homogenized for 5 minutes, and then centrifuged at 1000g for around 10 minutes. The molluscicidal activity of the obtained supernatant was tested.

2.2.2 Latex

By cutting the stem apex, the white latex from *Euphorbia tirucalli* was drained into glass tubes, and the latex was lyophilized at -40°C , and the lyophilized powder was employed for future use. To obtain the correct concentrations, the freeze-dried powder was combined with an appropriate volume of distilled water.

2.3 Test Animals

The target organisms for this research study, adult freshwater snails, *Lymnaea acuminata* ($2.5 \pm 0.9\text{cm}$ in shell height), and *Indoplanorbis exustus*, ($1.5 \pm 0.2\text{cm}$ in shell height) were collected from pool alongside the campus of Veer Abdul Hameed P.G. College, Medical Road, Gorakhpur district. The collected animals were kept in glass aquaria containing de-chlorinated tap water to acclimatize to laboratory conditions. The water in the aquaria was changed every 24 hours. To prevent the water fouling any dead animal were removed periodically. As the non-target organism, freshwater fish, *Channa punctatus* ($11.0 \pm 0.5\text{ cm}$ in total length), were captured from Ramgarh lake of Gorakhpur district.

2.4 Toxicity Experiments

The Singh and Agarwal [3] method was used to conduct the toxicity test for both snails, ten snails were housed in glass aquaria with 3L de-chlorinated tap water. The experimental snails were exposed for 96 h to four different concentrations of stem and leaves of *Phyllanthus*

niruri and stem bark and latex of *Euphorbia tirucalli*

For each concentration, six similar aquaria were set up. The snails were grown in the same way as the control group, but without any toxicity treatment.

Toxicity of aqueous extracts of stem bark and leaves and latex of both the plants under research, were also studied in a mixed population of fish and snails as the fish share the habitat with the snails. Accordingly, 10 snails *Lymnaea acuminata* and 10 fish *Channa punctatus*, were placed together in 6L of de-chlorinated tap water, and these mixed populations of fish and snails were checked for toxicity for previously determined LC_{90} (24h) of snails for 24h.

Snail behavioral reactions were detected for up to 2 hours after the treatment. Every 24h, 48h, 72h, and 96h, mortality was recorded LC_{10} , LC_{50} , and LC_{90} values, upper and lower confidence limits (UCL LCL), slope value 't' ratio, 'g' factor and heterogeneity were determined using the POLO computer program [4] utilizing the probit log analysis approach regression coefficient was also determined between exposure time and different values of LC_{50} [5].

3. RESULTS

Experimental conditions of water were calculated using APHA/WPCF method (1998) [6]. Accordingly, the parameters and their values determined were as follows:

Atmospheric temperature	35.0 -36.0 $^{\circ}\text{C}$
Water temperature	28.0- 29.0 $^{\circ}\text{C}$
pH of water	7.2-7.4
Dissolved Oxygen	6.9-7.4
Free carbon dioxide	4.6-6.7
Bicarbonate alkalinity	110.0- 111.0

3.1 Effects on Behavioural Changes and Poisoning Symptoms

Exposure to the aqueous extracts of root, latex, and stem bark of *Phyllanthus niruri* and *Euphorbia tirucalli*, caused significant behavioral changes in the freshwater snails *Lymnaea acuminata* and *Indoplanorbis exustus*, within 5-10 min of exposure to the toxicity test. Hyperactivity of the sluggish snails, during the first 30-40 min, was observed in the aquaria. With time, the snails got intoxicated and accordingly showed muscular twitching and

spiral twisting, which resulted ultimately in paralysis, and finally into the death of the snails.

3.2 Dose-mortality Response

LC values (LC_{10,50,90}) of aqueous extracts of stem bark and leaves of *Phyllanthus niruri* and stem bark and leaves and latex of *Euphorbia tirucalli* for period ranging from 24h to 96h for the snails, *Lymnaea acuminata* and *Indoplanorbis exustus* have been given in (Tables 1-6). In case of both the snail toxicity was time as well as dose dependent. There was a significant negative correlation between LC₅₀ values and exposure time (Tables 1-6). Thus increase in exposure time the LC₅₀ of *Phyllanthus niruri* aqueous stem bark and leaves extract, decreased from 267.89 mg DW/L (24h); > 225.13 mg DW/L (48h); > 205.10 mg DW/L (72h); > to 180.64 mg DW/L (96h) and 229.37 mg DW/L (24h); > 210.86 mg DW/L (48h) > 164.72 mg DW/L (72h); > to 136.40 mg DW/L (96h) in case of *Lymnaea acuminata* and *Indoplanorbis exustus*, respectively. Same trend of toxicity was observed in case of stem bark and latex of *Euphorbia tirucalli* against both the snails at all the exposure periods.

Additionally, laboratory tests show that both plant's latex and stem bark extracts were more toxic to *Indoplanorbis exustus* than *Lymnaea acuminata* across all exposure times. At higher doses, active moiety of plants, which were

effective against the snails, would also cause death amongst the fish. Consequently, a mixed population of 10 snails (*Lymnaea acuminata*) and 10 fish (*Channa punctatus*) were treated with the 24h, LC₉₀ of stem bark of *Phyllanthus niruri* and latex and stem bark of *Euphorbia tirucalli*, up to the LC₉₀ doses for snail *Lymnaea acuminata* there was no mortality amongst fish (Table 7).

The slope values given in toxicity Tables 1-6 were steep and heterogeneity factor was less than 1.0 indicates the result found to be within the 95% confidence limits of LC values. The regression test ('t' ratio) was greater than 1.96 and the potency estimation test ('g' value) was less than 0.5 at all probability levels.

4. DISCUSSION

"Several plant species having specific active chemical constituents in them, have been identified as natural molluscicides, and they are now being investigated. Among recent studies, some plant extracts have shown molluscicidal efficacy and saponins isolated from *Camellia oleifera* seeds have been used to create the novel chemical Luo-Wei, also known as tea-seed distilled saponin (TDS), which has been tested effectively against intermediate host snails *Oncomelania hupensis*, *Biomphalaria alexandrina*, and *Bulinus truncates* in China and Egypt" [7].

Table 1. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous stem and leaf extract of *Phyllanthus niruri* against *Lymnaea acuminata* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =173.98	160.40	187.40	9.11±4.05	3.39	0.012	0.24
	LC ₅₀ =267.89	230.98	402.17				
	LC ₉₀ =412.4	310.61	924.19				
48h	LC ₁₀ =101.03	59.10	120.93	3.00±1.35	3.16	0.173	0.19
	LC ₅₀ =225.13	197.22	326.02				
	LC ₉₀ =501.68	339.43	704.22				
72h	LC ₁₀ =80.46	30.24	105.29	2.73±1.23	2.83	0.249	0.24
	LC ₅₀ =205.10	181.77	294.36				
	LC ₉₀ =522.78	337.39	665.00				
96h	LC ₁₀ =93.23	62.73	110.67	2.75±1.24	3.99	0.238	0.22
	LC ₅₀ =180.64	168.01	203.17				
	LC ₉₀ =350.00	277.29	605.26				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of stem and leaf of *Phyllanthus niruri*
- Concentrations (Dry weight of stem and leaves) given are the final concentrations W/V in aquarium water
- Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: lower confidence limit UCL; upper confidence limit
- There was no mortality in control groups

Table 2. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous latex extract of *Euphorbia tirucalli* against *Lymnaea acuminata* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =0.65	0.53	0.77	0.13± 0.8	3.23	0.011	0.19
	LC ₅₀ =1.80	1.26	5.01				
	LC ₉₀ =4.93	2.50	38.6				
48h	LC ₁₀ =0.47	0.32	0.57	0.84±0.42	3.34	0.043	0.25
	LC ₅₀ =1.83	1.25	5.26				
	LC ₉₀ =7.07	3.15	73.78				
72h	LC ₁₀ =0.24	0.14	0.32	0.60 ±0.26	4.35	0.138	0.32
	LC ₅₀ =0.92	0.77	1.27				
	LC ₉₀ =3.45	2.09	10.22				
96h	LC ₁₀ =0.16	0.07	0.23	0.56 ±0.22	4.41	0.239	0.17
	LC ₅₀ =0.65	0.56	0.79				
	LC ₉₀ =2.63	1.70	6.70				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of latex extract of *Euphorbia tirucalli*
- Concentrations (Dry weight of latex) given are the final concentrations W/V in aquarium water
- Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: lower confidence limit. UCL; upper confidence limit
- There was no mortality in control groups

Table 3. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous bark extract of *Euphorbia tirucalli* against *Lymnaea acuminata* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =158.75	140.33	169.77	5.46±2.39	4.65	0.072	0.59
	LC ₅₀ =239.19	222.97	272.65				
	LC ₉₀ =360.38	304.67	509.15				
48h	LC ₁₀ =134.19	106.55	149.48	4.20±1.84	4.22	0.158	0.20
	LC ₅₀ =224.28	209.31	255.35				
	LC ₉₀ =374.84	307.66	582.97				
72h	LC ₁₀ =132.44	113.67	144.54	4.10±1.80	5.61	0.191	0.14
	LC ₅₀ =195.88	187.70	206.10				
	LC ₉₀ =289.72	260.70	349.98				
96h	LC ₁₀ =126.26	109.33	137.70	4.12±1.81	6.24	0.256	0.11
	LC ₅₀ =179.56	172.01	186.66				
	LC ₉₀ =255.37	236.63	289.39				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of bark extract of *Euphorbia tirucalli*
- Concentrations (Dry weight of latex) given are the final concentrations W/V in aquarium water
- Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: Lower confidence limit. UCL: Upper confidence limit
- There was no mortality in control groups

“Not until the mid-1960's was the first plant (*Phytolacca dodecandra*) used for control of schistosomiasis in an endemic focus, in Ethiopia [8]. The discovery of *Phytolacca dodecandra* berries as a naturally available potent molluscicide in Ethiopia has raised much awareness of the use of plants as effective

molluscicides, as they are effective in killing snail vectors while being safe to non-target organisms and easily degradable in the environment” [9,10].

“The molluscicidal properties of various leaf extracts prepared from tuba-tuba *Jatropha*

curcas L. were tested against the freshwater snail (*Oncomelania hupensis quadrasi*), an intermediate host of *Schistosoma* sp., the causative agent of schistosomiasis, with the crude petroleum ether extract at a concentration of 1.5 mg/mL¹ showing the highest mortality of

80%” [11]. “The molluscicidal efficacy of aqueous extract of *Achyranthes aspera* L. (Amaranthaceae) on adult *Biomphalaria pfeifferi* and *Lymnaea natalensis* snails, which are of major medicinal and veterinary value in Ethiopia, has been recently established” [12].

Table 4. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous stem and leaf extract of *Phyllanthus niruri* against *Indoplanorbis exustus* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =120.05	98.45	135.65	1.18±0.52	5.21	0.057	0.33
	LC ₅₀ =261.49	229.37	326.81				
	LC ₉₀ =569.58	422.65	999.79				
48h	LC ₁₀ =82.18	58.06	99.30	0.83±0.37	5.08	0.155	0.15
	LC ₅₀ =210.86	187.91	251.76				
	LC ₉₀ =540.99	397.20	977.31				
72h	LC ₁₀ =65.93	44.61	81.98	0.76±0.35	5.44	0.242	0.17
	LC ₅₀ =164.72	149.06	184.05				
	LC ₉₀ =411.50	323.24	636.52				
96h	LC ₁₀ =51.74	30.87	68.12	0.75±0.34	5.17	0.341	0.15
	LC ₅₀ =136.40	119.95	151.45				
	LC ₉₀ =359.57	286.41	547.97				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of stem and leaf of *Phyllanthus niruri*
- Concentrations (Dry weight of stem and leaves) given are the final concentrations W/V in aquarium water
- Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: Lower confidence limit UCL: Upper confidence limit.
- There was no mortality in control groups

Table 5. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous stem bark extract of *Euphorbia tirucalli* against *Indoplanorbis exustus* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =143.13	130.08	153.71	5.93±2.72	3.79	0.012	0.28
	LC ₅₀ =239.33	199.32	320.41				
	LC ₉₀ =367.44	278.86	731.45				
48h	LC ₁₀ =138.91	127.58	147.09	6.19±2.83	4.56	0.009	0.32
	LC ₅₀ =203.95	186.18	243.72				
	LC ₉₀ =299.44	248.80	440.99				
72h	LC ₁₀ =94.33	76.16	105.77	2.21±1.03	4.89	0.124	0.26
	LC ₅₀ =170.79	157.88	194.55				
	LC ₉₀ =309.22	251.03	466.56				
96h	LC ₁₀ =94.85	79.79	104.91	2.22±1.03	5.60	0.125	0.47
	LC ₅₀ =159.71	149.54	173.46				
	LC ₉₀ =266.66	228.09	352.40				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of stem bark extract of *Euphorbia tirucalli*
- Concentrations (Dry weight of stem and leaves) given are the final concentrations W/V in aquarium water. Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: lower confidence limit. UCL: upper confidence limit
- There was no mortality in control groups

Table 6. Toxicity (LC₁₀, LC₅₀ and LC₉₀) of aqueous latex extract of *Euphorbia tirucalli* against *Indoplanorbis exustus* at different time intervals

Exposure periods	Effective dose(W/V) (mg DW/L)	Limit (mg DW/L)		Slope value	't' ratio	'g' value	Heterogeneity
		LCL	UCL				
24h	LC ₁₀ =0.18	0.11	0.24	0.53±0.14	4.99	0.037	0.44
	LC ₅₀ =0.90	0.70	1.41				
	LC ₉₀ =4.39	2.39	14.27				
48h	LC ₁₀ = 0.08	0.12	0.44	0.38±0.86	4.24	0.124	0.28
	LC ₅₀ =0.77	0.56	1.33				
	LC ₉₀ =7.45	3.19	45.90				
72h	LC ₁₀ =0.04	0.01	0.07	0.34±0.73	4.24	0.206	0.45
	LC ₅₀ =0.51	0.39	0.79				
	LC ₉₀ =6.50	2.80	45.08				
96h	LC ₁₀ =0.02	0.006	0.05	0.34±0.70	4.58	0.292	0.70
	LC ₅₀ =0.28	0.21	0.36				
	LC ₉₀ =3.21	1.68	12.18				

- Batches of ten snails were exposed to four different concentrations of aqueous extracts of latex extract of *Euphorbia tirucalli*
- Concentrations (Dry weight of stem and leaves) given are the final concentrations W/V in aquarium water
- Regression coefficient showed that there was significant negative regression between exposure time and different LC values, LCL: Lower confidence limit UCL: Upper confidence limit
- There was no mortality in control groups

Table 7. Percent mortality (mean±SE) of *Lymnaea acuminata* and *Channa punctatus* caused by aqueous extract of stem, root and latex extracts (i.e. 24h LC₉₀ of snail) of *Phyllanthus niruri* and *Euphorbia tirucalli* after 24h exposure period

Plants	Plant parts	Experimental animals	Concentration (mgDW/L)(w/v)	% Mortality
<i>Phyllanthus niruri</i>	Stem and leaf	<i>L.acuminata</i>	412.48
		<i>C.punctatus</i>	-	zero
<i>Euphorbia tirucalli</i>	Stem bark	<i>L.acuminata</i>	498.84
		<i>C.punctatus</i>	-	zero
<i>Euphorbia tirucalli</i>	Latex	<i>L.acuminata</i>	4.93
		<i>C.punctatus</i>	-	zero

- Each aquarium contained ten fish (*Channa punctatus*) and ten snails (*Lymnaea acuminata*) in 6L de-chlorinated tap water
There was no mortality in control group

“Among the bioactive plant compounds, the most prevalent terpene compounds are responsible for molluscicidal activity. Thymol and α-pinene both have been investigated to be active against *B. glabrata*, causing mortality in concentration dependent patterns and having a fatal effect at doses commensurate with WHO recommendations (LC₉₀ of 7.11 and 10.34 g mL⁻¹, respectively), and are also reported to be inhibitors of acetyl cholinesterase of *B. glabrata* snails” [13]. In a study, the molluscicidal action of *Moringa oleifera* flowers was tested in *Biomphalaria glabrata* using an aqueous extract. In adult molluscs, this extract was fatal, with a lethal concentration of LC₅₀=2.37 mg/mL Also, fruits of the species *Randia nilotica* were

employed on molluscs of the species *Biomphalaria pfeifferi* and *Bulinus truncatus* in the second aqueous extract. On both molluscs employed in the trials, these extracts showed molluscicidal action. The molluscicidal effect of aqueous extract of leaves from the species *Anagallis arvensis* on molluscs, *Biomphalaria alexandrina*, was also feasible to reach lethal concentrations, with 50 mg/L being the lethal concentration and 90 mg/L being the lethal concentration [14].

The results show that *Phyllanthus niruri* aqueous stem and leaf extract and *Euphorbia tirucalli* aqueous stem, leaf, and latex extract are poisonous to the freshwater snails *Lymnaea*

acuminata and *Indoplanorbis exustus*. Toxicity data from this investigation indicates that the plant extracts mentioned above have significant molluscicidal efficacy. In *Lymnaea acuminata* and *Indoplanorbis exustus*, the extracts generated considerable behavioural changes, with the most visible symptom of discomfort being muscular and spiral twisting of the body, followed by crawling on one another. The type and early commencement of these behavioural reactions indicate that the extracts most likely include neurotoxins, which may be active in the exposed animals' neuromuscular system, among other places. Singh and Agarwal [3] noticed similar behaviour reactions in their laboratory investigation "on the acute toxicity of *Euphorbia royleana*, *Euphorbia antisyphiliatica*, and *Euphorbia tirucalli* lattices on snail *Lymnaea acuminata*".

No behavioural symptoms or death were seen in control animals that were not given treated water, showing that the changed behaviour and mortality were caused by something other than plant moieties. The animal's concentration-dependent response might be caused by a number of elements, including penetration rate, slope, variability, and maximum impact.

The LC₅₀ values of the studied aqueous extracts of *Phyllanthus niruri* aqueous stem bark and leaves extract dropped from 267.89 mg DW/L (24h); > to 180.64 mg DW/L (96h) and 229.37 mg DW/L (24h); > to 136.40 mg DW/L (96h) in the case of *Lymnaea acuminata* and *Indoplanorbis exustus*. At all exposure times, the stem bark and latex of *Euphorbia tirucalli* were shown to be harmful to both snails in the same way.

Reduced toxicity might be attributed to soil particle adsorption or temperature-induced acceleration of the toxicant breakdown process. Perschbacher and Sarkar [15] found that "the toxicity of *Masea ramentacea* and tea seed cake was short-lived, and fish could be stocked into ponds four days after spraying the plant pesticides".

Increased mortality with longer exposure durations might be caused by a number of reasons, which may work independently or in concert. For example, the updating of the active moiety is time dependent, resulting in a gradual rise in the drug's entry and effects in the snail body. The rate of detoxification in the animal body, as well as the stability (life span) of the active moiety of pesticide in the environment,

affect the mortality and exposure periods connection.

The data on toxicity is subjected to a statistical analysis, which reveals numerous key facts. The χ^2 test for goodness of fit (heterogeneity) revealed that the mortality counts were not statistically heterogeneous, and that other factors, such as resistance, did not have a significant impact on the LC₅₀ values, which were determined to be within the 95 % confidence limits. The slope is therefore a measure of the target animal's vulnerability to plant molluscicides. "Rapid absorption and start of effects are also indicated by a steep slope. Despite the fact that the slope alone is not a highly accurate predictor of toxicological mechanism, it is a valuable metric for this type of research. Since LC₅₀ of the aqueous extracts was within the 95% confidence levels, it is self-evident that the concentration response lines would fall within the same range in a replication test of random samples" [16].

5. CONCLUSION

It is believed that the aqueous extract of above medicinal plant extracts used for the aquatic snail control and safe for the doses against non-target organism which share the same habitat.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh SK, Yadav RP, Singh A. The toxicity of *Thevetia peruviana* and *Alstonia scholaris* plants to target and non-target aquatic organism. Proceedings of the Symposium on Functional Biodynamics and Ecophysiology of Animals. 2009:301–307.
2. Yadav RP, Singh A. efficacy of *Euphorbia hirta* latex as plant derived molluscicides against freshwater snails. Rev. Inst. Med. Trop. Sao Paulo. 2011;53(2):101-106.
3. Singh A, Agarwal RA. Possibility of using latex of euphorbiales for snail control. The Sci. Total Environment. 1988;77:231-236.
4. Robertson JL, Russell RM, Preisler HK, Savin NE. Bioassay with arthropods: POLO computer programme for analysis of bioassay data. 2nd ed. Boca Raton: CRC Press. 2007:1-224.

5. Sokal RR, Rohlf FJ. Introduction of biostatistics. W. H. Freeman and Company, San Francisco. 1973;368.
6. APHA /AWWA/ WEF. Standard methods for the examination of water and wastewater. 20th edition, American Public Health Association, New York, USA; 1998.
7. Jia TW, Wang W, Sun LP. et al. Molluscicidal effectiveness of Luo-Wei, a novel plant-derived molluscicide, against *Oncomelania hupensis*, *Biomphalaria alexandrina* and *Bulinus truncatus*. Infect Dis Poverty. 2019;8:27.
8. Lemma AP, Goll J. Duncan and Mazengia B. Control of schistosomiasis with use of endod in Adwa, Ethiopia: Results of a five year study. Proceedings of the International Conference on Schistosomiasis, Cairo 5.0. P. Press. 1978:415-436.
9. Lemma A. Preliminary report on the molluscicidal property of endod (*Phytolacca dodecandra*), Ethiopian Medical Journal. 1965;3:187.
10. Kloos H, McCullough FS. Plant molluscicides. Journal of Plant Medicinal. 1982;46:195-209.
11. Ludevese F, Ludevese Pascua G, Portugaliza H. Molluscicidal activity of *Jatropha curcas* L. crude leaf extracts against *Oncomelania hypensis quadrasi* mollendorf. Journal of Science, Engineering and Technology. 2018;6:177-186.
12. Mandefro B, Mereta ST, Tariku Y, et al. Molluscicidal effect of *Achyranthes aspera* L. (amaranthaceae) aqueous extract on adult snails of *Biomphalaria pfeifferi* and *Lymnaea natalensis*. Infect Dis Poverty. 2017;6:133.
13. Edilene Carvalho Gomes Ribeiro, José Antonio Costa Leite, Tássio Rômulo Silva Araújo Luz, Daniella Patrícia Brandão Silveira, Samara Araújo Bezerra, Gleycka Cristine Carvalho Gomes Frazão, Luciana Patrícia Lima Alves Pereira, Everton Gomes Guimarães dos Santos, Paulo Roberto Campos Flexa Ribeiro Filho, Alexandra Martins Santos Soares, Ricardo Luvizotto Santos, Denise Fernandes Coutinho. Molluscicidal activity of monoterpenes and their effects on inhibition of acetylcholinesterase activity on *Biomphalaria glabrata*, an intermediate host of *Schistosoma mansoni*, *Acta tropica*. 2021;223:106089.
14. Keyla Nunes Farias, José AAdosS, Robson Xavier Faria. Aqueous extracts: An alternative in the control of the intermediate host of schistosomiasis. 2021;4(3). OAJBS.ID.000302
15. Perschbacher PW, Sarkar J. Toxicity of selected pesticides to the snakehead, *channa punctata*. Asian Fish Sci. 1992;(2):249-254.
16. Rand GM, Petrocelli SR. Fundamentals of aquatic toxicology. Rand GM, Petrocelli SR. (Eds.) Hemisphere Publishing, New York. 1988:415.

© 2022 Khan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/95594>