



Effect of Phosphorus in Conjoint with Organics and Bioinoculant on Growth, Yield and Quality Parameters of Blackgram in *Typic Chromustert*

V. Suthakar ^{a*}, R. Indirani ^b, P. Kannan ^a and E. Murugan ^c

^a Department of Soils and Environment, Agricultural College and Research Institute, Madurai - 625104, India.

^b Department of Soils and Environment, Agricultural College and Research Institute, Chettinad - 630102, Sivagangai District, India.

^c Department of Plant Breeding and Genetics, Agricultural College and Research Institute, Madurai - 625104, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2131332

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89672>

Original Research Article

Received 16 May 2022
Accepted 21 July 2022
Published 27 July 2022

ABSTRACT

Aims: Use of phosphorus fertilizer along with farmyard manure, vermicompost and phosphorus solubilizing bacteria (*Bacillus megaterium*) in black calcareous soil was conducted to assess the impact on the growth, yield and quality parameters of blackgram variety VBN 8 during the year 2022.

Study Design: Randomized Block Design (RBD) with twelve treatments and three replications.

Place and Duration of Study: Kalligudi block of Madurai district, representing the Southern part of Tamil Nadu during the summer season of 2022.

Methodology: Twelve treatments were followed with various combinations of organics viz, vermicompost and farmyard manure, bioinoculant, and inorganic chemical fertilizer at different ratios. The combinations are 100 per cent and 75 per cent of P₂O₅ on STCR (Soil Test Crop Response) basis mixed with farmyard manure (1:10 ratio) and vermicompost (1:5 ratio). The impact of different treatments were recorded on shoot length, root volume, SPAD value, Leaf Area Index, grain yield, protein, cystine, and methionine content.

*Corresponding author: E-mail: saisuthakar007@gmail.com;

Results: Among the treatments, application of 100% of P₂O₅ on Soil Test Crop Response (STCR) basis incubated for 30 days with vermicompost at 1:5 ratio + phosphorus solubilizing bacteria @ 2 kg ha⁻¹ T₁₁ has improved the black gram shoot length (27.6 cm), root volume (2.43 cm³), SPAD value (45.91), Leaf Area Index at 20 DAS (1.09) and 40 DAS (3.7), grain yield (936.1 kg ha⁻¹) protein content (23.21%), cystine content (1.97%) and methionine content (2.29%).

Conclusion: The integration of vermicompost and bioinoculant along with phosphorus fertilizer in black calcareous soil has improved the blackgram yield by 35.8% over the use of recommended dose of fertilizer alone. This result highlights the need of integrated application of phosphorus fertilizer along with organics and phosphorus solubilizing bacteria in black calcareous soil for sustainable production of blackgram.

Keywords: Calcareous soil; bioinoculant; phosphorus; vermicompost.

1. INTRODUCTION

Phosphorus is a critical plant nutrient that cannot be replaced by another element to sustain plant life. The use of phosphorus fertilizer is a crucial component for high yielding food crops, and the majority of the mineral P used annually at the global level is for food production. It plays a vital role in the molecular structure of crops and aids in the transformation of energy as well as the regulation of several enzymatic activities. Phosphorus is also required for energy production, nucleic acid synthesis, photosynthesis, glycolysis, respiration, redox reactions, membrane synthesis and integrity, enzymatic activation or inactivation, carbohydrate metabolism and signaling. Therefore, insufficiency of phosphorus levels in the soil will greatly affect the growth and development of crops [1].

The bioavailability of phosphorus in soil is affected due to its complexation with soil mineral particles viz., calcium (Ca), magnesium (Mg), aluminum (Al), and iron (Fe) which will causes significant yield losses of crops [2]. In calcareous soils, calcium will react with the phosphorus which reduces its bioavailability due to the formation of calcium phosphate minerals which are less soluble at about pH 8. Surface adsorption and precipitation are key aspects of P retention in calcareous soil, which reduces its availability. As a result of the low P solubility, fertilizer phosphorus efficiency is relatively low in alkaline calcareous soil. Adding P fertilizer at normal rates with conventional methods may not give optimal crop yield and quality.

Several P fertilizer management strategies have been found to improve the P nutrition for plants grown in alkaline calcareous soil. Phosphorus application, combined with organic manures and phosphorus-solubilizing bacteria, effectively supplies P to growing plants in alkaline

calcareous soil [3]. Long-term manure application to alkaline calcareous soil had revealed that there is a significant amount of P accumulation, with 50 to 66 percent in plant available forms [4]. The affinity constants and adsorption capacity of soils for P are reduced by organic amendments, especially manure. This could be due to the production of organic acids by manure application competing for P fixation sites and the complexation of exchangeable cations by manure components [5]. Phosphorus solubilizing bacteria (PSB) can be beneficial because these bacteria will secrete some phenolic compounds, protons, organic and mineral acids into the soil, causing soil acidification and resulting in P release from calcium phosphate. PSB secreted organic acids chelate with some of the cations such as Ca²⁺, Al³⁺, and Fe³⁺, which may increase bioavailable P [6]. However, the potential of PSB in soils of a calcareous nature and with an alkaline reaction has not been well documented. Considering the crop phosphorus requirements and its availability in calcareous soil, this study was conducted to manage phosphorus availability by the integration of organic, phosphorus solubilizing bacteria and inorganic sources for optimum blackgram production in calcareous soils.

2. MATERIALS AND METHODS

The current experimental study was conducted on black gram, variety Vamban 8, during the summer season 2022 at Kalligudi block of Madurai district, which represents the southern part of Tamil Nadu. Climatic conditions of experimental site were 37.7°C and 20.3°C average maximum and minimum temperature respectively with a mean rainfall of 840 mm [7]. The soil texture was clay loam and classified as Vertisol soil order. The experiment was laid out in Randomized Block Design (RBD) with twelve treatments and three replications, as mentioned in Table 1.

Table 1. Treatment schedule for field experiment

Treatments	Details
T ₁	Absolute control
T ₂	Recommended N, P ₂ O ₅ , K ₂ O @ 25:50:25 kg ha ⁻¹
T ₃	100% of P ₂ O ₅ on STCR basis alone
T ₄	100% of P ₂ O ₅ on STCR basis incubated for 30 days with farm yard manure at 1:10 ratio
T ₅	75% of P ₂ O ₅ on STCR basis incubated for 30 days with farm yard manure at 1:10 ratio
T ₆	100% of P ₂ O ₅ on STCR basis incubated for 30 days with vermicompost at 1:5 ratio
T ₇	75% of P ₂ O ₅ on STCR basis incubated with vermicompost for 30 days at 1:5 ratio
T ₈	100% of P ₂ O ₅ on STCR basis + PSB @ 2 kg ha ⁻¹
T ₉	T ₄ + PSB @ 2 kg ha ⁻¹
T ₁₀	T ₅ + PSB @ 2 kg ha ⁻¹
T ₁₁	T ₆ + PSB @ 2 kg ha ⁻¹
T ₁₂	T ₇ + PSB @ 2 kg ha ⁻¹

Table 2. Initial physico-chemical properties of the experimental soil

Particulars	Values	Methods
A. Mechanical composition		
Clay (%)	34.76	International pipette method [12]
Silt (%)	20.41	
Fine sand (%)	17.01	
Coarse sand (%)	25.68	
Texture class	Clay loam	
B. Physical properties		
		Core sampler method [13]
Bulk density (Mg m ⁻³)	2.50	
Particle density (Mg m ⁻³)	1.34	
Porosity (%)	41	
C. Physico-chemical properties		
Ph	8.32	Potentiometry [14]
EC (dSm ⁻¹)	0.18	Conductometry [14]
D. Chemical properties		
Free CaCO ₃ (%)	7.63	Rapid titration method [12]
Organic carbon (g kg ⁻¹)	3.80	Dichromate wet digestion method [15]
Alkaline KMnO ₄ – Nitrogen (kg ha ⁻¹)	234	Alkaline permanganate method [16]
Olsen -Phosphorus (kg ha ⁻¹)	10.34	Olsen method [17]
NH ₄ OAc – Potassium (kg ha ⁻¹)	214.24	Neutral normal NH ₄ OAc method [18]

The enriched phosphorus fertilizer was prepared by mixing two different sources of organic manures with inorganic phosphorus fertilizer (single super phosphate) at different combinations. The combinations are 100 per cent and 75 per cent of P₂O₅ on STCR (Soil Test Crop Response) basis mixed with farmyard manure (1:10 ratio) and vermicompost (1:5 ratio). These combination were prepared and kept for incubation at 60 per cent moisture content in grow bag over a period of one month. Nitrogen and potassium were applied as urea and muriate of potash on soil test crop response basis in all the treatments except T₁, T₂, and T₈. T₁ was the

absolute control, T₂ received recommended dose of fertilizer (N, P₂O₅, K₂O @ 25:50:25 kg ha⁻¹) whereas T₈ was an application of phosphorus on soil test crop response basis with phosphorus solubilizing bacteria.

The observations viz, shoot length, root volume, soil plant analysis development (SPAD) chlorophyll meter value and seed yield were measured at the harvest stage in each plot. Leaf Area Index was measured at 20 and 40 days after sowing (DAS). SPAD value was measured by using MC - 100 chlorophyll concentration meter. Root volume was measured by average

root length and breadth. The quality parameters were estimated by standard procedure viz, protein content by Lowry's method [8], cystine content by Goa procedure [9], and methionine content by sodium nitroprusside method [10]. The data collected were analyzed by using AGRES software [11].

3. RESULTS AND DISCUSSION

3.1 Effect of Phosphorus Fertilizer Integrated with Organics and PSB on Growth Attributes of Blackgram

3.1.1 Shoot length

The perusal of the data (Table 3) shows that the growth parameters were significantly affected by the application of inorganic phosphorus fertilizer along with organics and PSB. A higher shoot length of 27.6 cm at the harvest stage was recorded in T₁₁ (100% of P₂O₅ on STCR basis incubated for 30 days with vermicompost at 1:5 ratio + PSB @ 2 kg ha⁻¹). T₁₁ was followed by T₁₂ (75% of P₂O₅ on STCR basis incubated with vermicompost for 30 days at 1:5 ratio + PSB @ 2 kg ha⁻¹) of 25.9 cm. The least shoot length was reported in absolute control T₁ (13.07 cm) which was on par with T₃ (13.57 cm). The treatment which received the recommended dose of fertilizer T₂ alone recorded comparatively lower shoot length (17.07 cm) than the conjoint application of inorganic fertilizer with organics and PSB. A positive response from the combined application of vermicompost, PSB and inorganic fertilizer may be due to their specific role in shoot morphology, cell division, cell wall plasticity, energy metabolism, and improved root development, which may promote better shoot growth. According to [19], phosphate solubilizing bacteria and organics application along with phosphorus fertilizer might increase the bioavailability of insoluble soil phosphorus for plant use by producing organic acids. [20] reported that, because of the mineralization, solubilization, and translocation of phosphorus through the production of organic acid and proton extrusion by the use of biofertilizer and organics have improved the phosphorus availability to plants, which in turn improves the plant shoot length.

3.1.2 Root volume

The root volume was also enhanced by the application of enriched phosphorus with vermicompost and PSB. Among the treatment,

T₁₁ achieved a higher root volume at the harvest stage (2.43 cm³) which was followed by T₁₂ (2.29 cm³). The treatment which received 100% of P₂O₅ on STCR basis incubated for 30 days with vermicompost at 1:5 ratio without PSB T₆ recorded comparatively lower root volume (1.93 cm³). The lowest root volume was recorded in absolute control T₁ (1.22 cm³). The positive influence of phosphorus, organics and bioinoculant on the formation, multiplication, and elongation of root cells could have resulted in the growth of the voluminous root system in blackgram. Combined application of PSB and organics along with inorganic fertilizer leads to higher Indole Acetic Acid production, a hormone produced by plants that regulate the growth and elongation of roots, which was likely the cause of improvements in seedlings' root morphological features. This is consistent with the study of [21].

3.1.3 SPAD value

The data on the SPAD value in various treatments revealed that the higher SPAD value of blackgram at the harvest stage was significantly affected by the different combinations of inorganic phosphorus fertilizer, farmyard manure, vermicompost, and PSB. The SPAD value was higher in the treatment where phosphorus fertilizer was integrated with vermicompost and PSB T₁₁ (45.91) as compared to the application of organics, PSB, and phosphorus fertilizer alone. The treatment which received organic and inorganic fertilizer without PSB T₆ recorded comparatively lowest SPAD value of 39.01. The lowest SPAD value was observed in absolute control T₁ (29.16) which was on par with T₃ (29.76). This occurred mostly as a result of PSB having the ability to enhance nitrogen fixation, which can encourage plant uptake of N and leaf chlorophyll content, this agrees with the observations of [22]. One of the most important parameters measured as an indicator of leaf nitrogen content, chloroplast development, and general plant health is the chlorophyll content of leaf tissue [23].

3.1.4 Leaf area index

The Leaf Area Index at 20 and 40 DAS was significantly higher in the T₁₁ treatment with 100% of P₂O₅ on STCR basis incubated for 30 days with vermicompost at 1:5 ratio + PSB @ 2 kg ha⁻¹ (1.09 and 3.70) followed by T₁₂ (1.01 and 3.43). The lowest Leaf Area Index at 20 and 40 DAS was noticed in absolute control T₁ (0.4 and 1.83) which was on par with the application of inorganic phosphorus fertilizer alone T₃ (0.42 and

1.85). The higher LAI might be due to the greater supply of nitrogen by this treatment which in turn increased the leaf size. Same observation was found by [24].

3.2 Effect of Phosphorus Fertilizer Integrated with Organics and PSB on Yield of Blackgram

The grain yield of blackgram recorded in the treatment T₁₁ (936.1 kg ha⁻¹) was significantly higher (Fig.1). The grain yield of absolute control T₁ (431.8 kg ha⁻¹) was on par with the treatment

received single super phosphate alone T₈ (487.06 kg ha⁻¹) without organic manure and PSB. Likewise, there was no significant difference in the grain yield of the treatments T₄ (728.66 kg ha⁻¹) and T₇ (739.28 kg ha⁻¹). However, yield increase was more prominent in PSB inoculated and organics applied treatments compared to un-inoculated. This could be attributed to the release of unavailable P into the mobile pool by rapid mineralization of organic P and solubilization of Ca-P through acidifying and chelating mechanisms [25].

Table 3. Integrated effect of phosphorus in conjoint with organics and bioinoculant on growth parameters of blackgram

Treatments	Shoot length (cm)	Root volume (cm ³)	SPAD value	Leaf Area Index	
				20 DAS	40 DAS
T ₁	13.07	1.22	29.16	0.40	1.83
T ₂	17.07	1.51	33.37	0.56	2.39
T ₃	13.57	1.27	29.76	0.42	1.85
T ₄	19.63	1.76	36.69	0.70	2.73
T ₅	18.40	1.65	35.07	0.64	2.57
T ₆	21.60	1.93	39.01	0.82	3.01
T ₇	20.23	1.80	37.45	0.75	2.74
T ₈	14.87	1.38	31.15	0.47	2.06
T ₉	25.30	2.23	43.03	0.97	3.39
T ₁₀	23.70	2.10	41.21	0.90	3.21
T ₁₁	27.60	2.43	45.91	1.09	3.70
T ₁₂	25.90	2.29	43.92	1.01	3.43
S.Ed	0.40	0.04	0.66	0.01	0.05
C.D (p=0.05)	0.84	0.09	1.38	0.02	0.12




Fig. 1. Integrated effect of phosphorus in conjoint with organics and bioinoculant on yield of blackgram

Table 4. Integrated effect of phosphorus in conjoint with organics and bioinoculant on quality parameters of blackgram

Treatment	Protein content (%)	Cystine content (g/16g N)	Methionine content (g/16g N)
T ₁	14.23	0.89	0.68
T ₂	16.51	1.22	1.08
T ₃	14.50	0.94	0.75
T ₄	18.64	1.41	1.37
T ₅	17.75	1.32	1.24
T ₆	19.87	1.58	1.71
T ₇	18.88	1.45	1.44
T ₈	15.41	1.06	0.90
T ₉	21.97	1.79	2.02
T ₁₀	21.08	1.70	1.88
T ₁₁	23.21	1.97	2.29
T ₁₂	22.28	1.84	2.00
S.Ed	0.30	0.03	0.03
C.D (p=0.05)	0.63	0.07	0.07

3.3 Effect of Phosphorus Fertilizer Integrated with Organics and PSB on Quality Parameters of Blackgram

The data depicted in Table 4 revealed that the integrated application of chemical fertilizer with organics and bioinoculant has a significant effect on protein, cystine, and methionine content. Addition of T₁₁ recorded higher protein (23.21%), cystine (1.97 g/16g N), and methionine content (2.29 g/16g N) which was on par with T₁₂ (22.28% and 1.84 and 2 g/16g N). The lowest protein, cystine and methionine content were recorded with absolute control T₁ which is statistically on par with T₈. As PSB plays a important role in N fixation and the fixed nitrogen might have enhanced the synthesis of amino acids. Besides, phosphate would have replaced the sulphate from the colloidal complex and increased the absorption of sulphate by the black gram. According to [26], the protein content is dependent on plant growth and nutritional composition. Increased protein content in grain may be due to increased nitrate uptake and translocation, which provides nitrogen for amino acid synthesis. The highest percentage of crude protein content in seeds of blackgram with vermicompost and PSB applied treatment might be due to more availability of nitrogen. PSB may enhance both the rhizosphere's nutrient environment and its use in the plant system.

4. CONCLUSION

In black calcareous soil, it is possible to attain higher growth, yield and quality of blackgram by

integrated use of phosphorus fertilizer along with vermicompost and phosphorus solubilizing bacteria. Our results confirmed that application of 100% of P₂O₅ on STCR basis incubated for 30 days with vermicompost at 1:5 ratio + PSB @ 2 kg ha⁻¹ with nitrogen and potassium on STCR basis performs well in black calcareous soil. It reduced the phosphorus fixation and also solubilizes the fixed phosphorus as calcium phosphate by the production of various organics acid, which in turn increases the phosphorus availability, growth, yield, and quality of crops.

ACKNOWLEDGEMENTS

Authors thank the Tamil Nadu Agricultural University, AC & RI, Madurai for providing laboratory facilities for conducting the experiment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hussain RM. The effect of phosphorus in nitrogen fixation in legumes. *J Plant Nutr Soil Sci.* 2017;167(2):125-37.
2. Lun F, Liu J, Ciais P, Nesme T, Chang J, Wang R et al. Global and regional phosphorus budgets in agricultural systems and their implications for phosphorus-use efficiency. *Earth Syst Sci Data.* 2018;10(1):1-18. DOI: 10.5194/essd-10-1-2018.

3. Khalil AA. Significance of some soil amendments and phosphate dissolving bacteria to enhance the availability of phosphate in calcareous soil. *Int Sch Res Not.* 2013;2013:1-7. DOI: 10.1155/2013/438949.
4. Zhang L, Chen J, Chu G. Legacy phosphorus in calcareous soil under 33 years of P fertilizer application: Implications for efficient P management in agriculture. *Soil Use Manag.* 2022;38(3):1380-93. DOI: 10.1111/sum.12792.
5. Ahmad M, Ishaq M, Shah WA, Adnan M, Fahad S, Saleem MH et al. Managing phosphorus availability from organic and inorganic sources for optimum wheat production in calcareous soils. *Sustainability.* 2022;14(13):7669. DOI: 10.3390/su14137669.
6. Adnan M, Fahad S, Zamin M, Shah S, Mian IA, Danish S et al. Coupling phosphate-solubilizing bacteria with phosphorus supplements improve maize phosphorus acquisition and growth under lime induced salinity stress. *Plants (Basel).* 2020;9(7):900. DOI: 10.3390/plants9070900, PMID 32708749.
7. Karpagam S, Mary PCN, Kannan S, Gurusamy S, Shanmugasundaram R, Ramamoorthy P. Effect of climate change on morphological characteristics of the soils of Vaigai River Basin, Alluvial Tract, Madurai District, Tamil Nadu. *J Pharmacogn Phytochem.* 2020;9(6):312-6.
8. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951;193(1):265-75. PMID 14907713.
9. Goa JOHN, Larsen B, Sjöberg B, Dam H, Sjöberg B, Toft J. Micro determination of cysteine plus cystine in proteins. *Acta Chem Scand.* 1961;15:853-5. DOI: 10.3891/acta.chem.scand.15-0853.
10. Horn MJ, Jones DB, Blum AE. Colorimetric determination of methionine in proteins and foods. *J Biol Chem.* 1946;166(1):313-20. DOI: 10.1016/S0021-9258(17)35008-1, PMID 20273698.
11. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & sons; 1984.
12. Piper CS. Soil and plant analysis. Bombay, India: Hans Publishers; 1966.
13. Gupta RP, Dakshinamoorthy C. Procedures for physical analysis of soil and collection of agrometeorological data. New Delhi: Indian Agricultural Research Institute. 1980;293.
14. Jackson ML. Soil chemical analysis. New Delhi: Prentice Hall of Indian Private Limited; 1973.
15. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci.* 1934;37(1):29-38. DOI: 10.1097/00010694-193401000-00003.
16. Subbiah BV, Asija GL. A rapid method for the estimation of nitrogen in soil. *Curr Sci.* 1956;26:259-60.
17. Olsen. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. Washington: United States Department of Agriculture; 1954.
18. Standford S, English L. Use of flame photometer in rapid soil tests for K and Ca. *Agron J.* 1949;41:446-7.
19. Alori ET, Glick BR, Babalola OO. Microbial phosphorus solubilization and its potential for use in sustainable agriculture. *Front Microbiol.* 2017;8:971. DOI: 10.3389/fmicb.2017.00971, PMID 28626450.
20. Singh J, Bhatt R, S DS, Dhillon BS, Al-Huqail AA, Alfagham A et al. Integrated use of phosphorus, farmyard manure and biofertilizer improves the yield and phosphorus uptake of black gram in silt loam soil. *Plos One.* 2022;17(4):e0266753. DOI: 10.1371/journal.pone.0266753, PMID 35421141.
21. Elhaissofi W, Khourchi S, Ibnyasser A, Ghoulam C, Rchiad Z, Zeroual Y et al. Phosphate solubilizing rhizobacteria could have a stronger influence on wheat root traits and aboveground physiology than rhizosphere P solubilization. *Front Plant Sci.* 2020;11:979. DOI: 10.3389/fpls.2020.00979, PMID 32765544.
22. Israr D, Mustafa G, Khan KS, Shahzad M, Ahmad N, Masood S. Interactive effects of phosphorus and *Pseudomonas putida* on chickpea (*Cicer arietinum* L.) growth, nutrient uptake, antioxidant enzymes and organic acids exudation. *Plant Physiol Biochem.* 2016;108:304-12. DOI: 10.1016/j.plaphy.2016.07.023, PMID 27485620.

23. Ling B, Goodin DG, Mohler RL, Laws AN, Joern A. Estimating canopy nitrogen content in a heterogeneous grassland with varying fire and grazing treatments: konza Prairie, Kansas, USA. *Remote Sens.* 2014;6(5):4430-53. DOI: 10.3390/rs6054430. phosphate-solubilizing bacteria with phosphorus supplements improve maize phosphorus acquisition and growth under lime induced salinity stress. *Plants (Basel)*. 2020;9(7):900. DOI:10.3390/plants9070900, PMID 32708749.
24. Joshi D, Gediya KM, Patel JS, Birari MM, Gupta S. Effect of organic manures on growth and yield of summer cowpea [*Vigna unguiculata* (L.) Walp.] under middle Gujarat conditions. *Agric Sci Dig.* 2016;36(2):134-7.
25. Adnan M, Fahad S, Zamin M, Shah S, Mian IA, Danish S et al. Coupling
26. Rekha K, Neha RPP C, Patel S. Effect of FYM, phosphorus and PSB on growth, yield and quality of Greengram [*Vigna radiata* (L.) Wilczek] on loamy sand. *Int J Bio-Resource Stress Manag.* 2018;9(2): 220-3. DOI:10.23910/IJBSM/2018.9.2.3C0375b.

© 2022 Suthakar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89672>