



## Association of Endothelial Nitric Oxide Synthase Gene Polymorphisms T-786c & 27bp (4b/4a) with Obesity in Egypt

Rizk Elbaz<sup>1</sup>, Lamiaa A. A. Barakat<sup>2</sup>, Ahmed Nageb<sup>2\*</sup> and Mohamed Sherif<sup>3</sup>

<sup>1</sup>Genetics Unit, Faculty of Medicine, Mansoura University, Egypt.

<sup>2</sup>Department of Chemistry, Faculty of Science, Port Said University, Egypt.

<sup>3</sup>Internal Medicine Hospital, Faculty of Medicine, Mansoura University, Egypt.

### Authors' contributions

*This work was carried out in collaboration among all authors. Author AN designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors RE and LAAB managed the analyses of the study. Author MS managed the literature searches. All authors read and approved the final manuscript.*

### Article Information

DOI: 10.9734/IJBCRR/2019/v25i230073

#### Editor(s):

(1) Dr. Cheorl-Ho Kim, Molecular and Cellular Glycobiology Unit, Department of Biological Science, Sungkyunkwan University, Chunchun-Dong 300, Jangan-Gu, Suwon City, Kyunggi-Do 440-746, South Korea.

#### Reviewers:

(1) Naguibou M. Djibril, University of Abomey Calavi (UAC), Cotonou- Benin.

(2) Samah Mohamed Elattar, Cairo University, Egypt.

(3) Yaya Kassogue, University of Sciences, Techniques and Technologies of Bamako, Mali.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47557>

Received 18 December 2018

Accepted 09 March 2019

Published 18 March 2019

Original Research Article

### ABSTRACT

**Background and Objective:** Endothelial nitric oxide synthase gene polymorphism (eNOS) is one of three isoforms that synthesize nitric oxide (NO), that participates in several biological processes have been associated with obesity. This study was undertaken to determine if eNOS gene (T786C) and 27bp (4b/4a) were associated with susceptibility of obesity.

**Materials and Methods:** The study was carried out on 200 cases divided into 100 obese patient and 100 healthy as control. The mean age cases was (27.02 ± 10.90) they include 79 female and 21 males. All participants were subjected to an estimation of their body mass index (BMI), weight hip ratio (WHR), in addition to random blood sugar (RBS), total cholesterol, triglyceride (TG), and lactate dehydrogenase enzyme (LDH). DNA was amplified using PCR-SSP for detection of relation between polymorphism and endothelial nitric oxide synthase gene in two parts T786C and 27bp (4b/4a).

\*Corresponding author: E-mail: [drahmednageb@gmail.com](mailto:drahmednageb@gmail.com);

**Results:** All cases showed that there were significant difference between cases and controls regarding to their chemical lab's analysis (TG, Cholesterol, LDL and HDL).All cases showed significant frequency of T786C TT, CC, TC vs. controls ( $p < 0.001$ ) these was considered risk factor for disease. On the other hand there no significant difference between 27bp aa, bb, and ab ( $p = 0.618$ ) vs. controls.

**Conclusion:** The polymorphism T786C not the 27bp in eNOS was associated with obesity.

*Keywords: Endothelial nitric oxide; gene; polymorphism; obesity.*

## ABBREVIATIONS

*Endothelial nitric oxide (eNOS),  
Polymerase chain reaction with sequence  
specific primers PCR-SSP.  
Nitric oxide NO.*

## 1. INTRODUCTION

Obesity is a medical condition in which surplus body fat accumulated to the range that it might had a negative effect on health [1], people are generally considered obese when their body mass index (BMI), a measurement obtained by dividing a person's weight by the square of the person's height, is over  $30 \text{ kg/m}^2$ , with the range  $25\text{--}30 \text{ kg/m}^2$  defined as overweight [1], some East Asian countries use lower values [2], obesity increases the incidence of various diseases and conditions, specially cardiovascular diseases, type 2 diabetes, obstructive sleep apnea, definite types of cancer, osteoarthritis and depression [3,4].

Obesity is most commonly caused by a mixing of excessive food intake, lack of physical activity, and genetic susceptibility [1,5], a few cases are caused firstly by genes, endocrine disorders, medications, or mental disorder [6], on the other hand obese people eat little next to gain weight because of a slow metabolism is not medically supported [7], on average, obese people have a greater energy usage than their normal people because of the energy required to maintain an increased body mass [7,8].

Obesity might be a cause of death which can be preventable worldwide, with increasing rates in adults and children [1]. In 2015, 600 million adults (12%) and 100 million children were obese in 195 countries [9] Obesity is more common in women than men [1]. Several studies viewed that obesity is one of the most dangerous public health problems of the 21st century [10]. In 2013, obesity is classified as a disease by the American Medical Association [11,12].

Impaired nitric oxide production is involved in the pathogenesis of several diseases such as hypertension, diabetes mellitus, obesity, erectile dysfunction, and migraine [13], a large number of studies showed that polymorphisms in  $\text{NOS}_3$  gene affect the susceptibility to these diseases [13], although  $\text{NOS}_3$  is a highly polymorphic gene, three genetic polymorphisms in this gene have been widely studied: the single nucleotide polymorphisms (SNPs) g.-786T>C located in  $\text{NOS}_3$  promoter and in exon 7, respectively, and the variable number of tandem repeats 4b/4a (VNTR) characterized by 27 bp repeat in intron 4 [13]. The C allele for the T786C polymorphism, which results in reduced eNOS expression and nitric oxide production was associated with increased risk for hypertension [14]. The VNTR in intron 4 affects eNOS expression [15]. And the susceptibility to hypertension [14]. Obesity [16].

## 2. MATERIALS AND METHODS

### 2.1 Study Group

This study includes 200 cases 100 obese patients they were recruited from the Department of Diabetes and Endocrine Unit in Specialized Medical Hospital Mansoura University, Egypt as well as Ministry of Health Hospitals of Dakahlia, Egypt during the period September 2016 to May 2018, and 100 healthy control. The mean age of cases were  $27.02 \pm 10.90$  years they were in the form of 21 male and 79 female. According to the definition of metabolic syndrome given by WHO, ATP and IDF (75%) of patient were classified as having metabolic syndrome while the rest, (25%) were not complicated and were characterized as just having simple obesity.

### 2.2 Control Group

For comparison 100 healthy controls were selected.

### 2.3 Biochemical Analysis

After 12 h of fasting, a blood was collected from each case and control in an empty tube blood sample for biochemical analysis. If the sample were not analyzed immediately, they will frozen and stored at -70 C. Total cholesterol, triglyceride (TG), LDL and HDL were measured by enzymatic methods on automatic biochemistry analyzer.

### 2.4 Capture Column Kit Extraction and Purification

The generation DNA purification capture column kit (Gentra System, USA) is based on a proprietary system that uses two reagents, a DNA purification solution and a DNA elution solution, along with a specially formulated purification matrix. In this kit , a sample is applied directly to the purification matrix contained a spin column .the cells contained in sample lyse upon contact with the matrix .once the cells were lysed, DNA was captured by the matrix material which make it possible to efficiently wash away contaminants ,leaving the DNA bound to the matrix. Contaminants, including protein heme and RNA were removed from the matrix by washing with DNA purification solution.

Following removal contaminants, the DNA released from the matrix using DNA elution solution and heat. Samples of purified DNA were ready for analysis and not require precipitation.

### 2.5 Pcr Amplifications of Each Enos Studied

Single nucleotide polymorphism (SNPs) for nitric oxide synthase gene (eNOS) were genotyped in this case-control study C786T and 27bp polymorphism using polymerase chain reaction PCR. Amplification were performed in sequence-specific primer polymerase chain reaction (SSP-PCR) employing a forward and reverse primer for each part. The region containing one (Restriction Fragment Length Polymorphisms) RFLPs within the eNOS gene was amplified with tag DNA polymerase, PCR buffer, Mgcl2 and dNTPs.

The entire reaction volume plus 5  $\mu$  L of bromophenol blue track dye were loaded into 2% agarose gel (Bohringer Mannheim) containing ethidium bromide. And for 30 minutes at 100V Gels were electrophoresed, then photographed

under UV light (320 nm) and then detect the presence or absence of an allele specific bands.

### 2.6 Primer Sequences and PCR Condition of eNOS Gene Polymorphism

The T786C genotype was performed using PCR amplification, the amplified product was digested with NgoMIV enzyme. Briefly primer sequences were forward primer: 5'-ATG CTC CCA CCA GGG CAT CA-3' and reverse primer: 5'-GTC CTT GAG TCT GAC ATT AGG G-3'.

The 27 bp (4b/4a) was determined using PCR amplification, not followed by restriction enzyme digestion of the amplified product. Briefly primer sequences were forward primer: 5'AGG CCC TAT GGT AGT GGC CTT T-3' and reverse primer: 5'TGC TCC TGC TAC TGA CAG CA-3'

### 2.7 Statistical Analysis

Statistical analysis of data was done using the software statistical package (SPSS program version 17). The student t-test was used to compare the numerical values related to cholesterol, other chemical parameter and body mass index whereas CHI square test used to compare frequencies of different genotypes and alleles between cases and controls.

## 3. RESULTS

Cases and controls showed a non-significant difference regarding to their age ( $p = 0.74$ ). However, cases showed a significant levels of BMI, cholesterol, TG, HDL-C and LDL-C ( $p < 0.001$ ) (Table 1).

Regarding to descriptive data of studied cases of obesity, cases showed a significant difference vs. control (normal, no disease) with  $p < 0.001$  (Table 2).

Regarding to distribution of eNOS gene polymorphism (T786C) (table 3): all genotypes (TT), (TC), and (CC) were highly significant ( $p < 0.001$ ) vs. controls. While on alleles analysis both (T) and (C) were significantly ( $p < 0.001$ ).

Comparing all cases with obesity and healthy controls regarding their genotype distribution of eNOS gene polymorphism (27 bp), (Table 4): all genotypes (aa), (ab), and (bb) were non-significantly ( $p = 0.618$ ) vs. controls. While on alleles analysis (a) and (b) did not show any significant difference. ( $p = 0.482$ ).

**Table 1. Descriptive data of studied cases of obesity and healthy controls**

	Patients (N=100)	Control (N=100)	t	P
Hip	122.69 ± 12.96	89.26 ± 17.18	15.536	<0.001*
Weight	106.03 ± 16.95	68.66 ± 17.77	15.216	<0.001*
Height	162.47 ± 8.26	166.38 ± 7.55	3.495	0.001*
BMI	40.13 ± 6.40	25.02 ± 7.67	15.132	<0.001*
WHR	0.95 ± 0.14	0.82 ± 0.12	7.268	<0.001*
waist	116.16 ± 15.47	74.57 ± 24.76	14.245	<0.001*
Age	27.02 ± 10.90	27.51 ± 10.26	0.327	0.744
Cholesterol	246.32 ± 60.23	181.16 ± 44.48	8.703	<0.001*
TG	140.76 ± 95.91	101.74 ± 47.85	3.640	<0.001*
HDL-C	49.94 ± 15.60	37.54 ± 13.48	6.014	<0.001*
LDL-C	168.85 ± 64.86	124.10 ± 40.89	5.835	<0.001*

N: number of cases, t: Student t-test, TG: Triglyceride, HDL: High-density lipoprotein, LDL: Low-density lipoprotein, \*p =0.001 (significant).

**Table 2. Descriptive data of studied cases of obesity**

Disease	Patients		Control		χ <sup>2</sup>	p
	N	%	N	%		
obesity	53	53.0%	0	0.0%	200.000	<0.001*
obesity+D.M	21	21.0%	0	0.0%		
obesity+HTN	12	12.0%	0	0.0%		
obesity+D.M+ HTN	14	14.0%	0	0.0%		
normal, no disease	0	0.0%	100	100.0%		

N: Number of cases, %: percentage of cases, χ<sup>2</sup>: Chi-square test  
D.M.: Diabetes Mellitus, HTN. : Hypertension

**Table 3. Comparison between all cases with obesity and healthy controls regarding their genotype distribution of eNOS gene polymorphism in (T786C)**

T786C		Patients		Control		χ <sup>2</sup>	p
		N	%	N	%		
Genotype	TT	33	33.0%	84	84.0%	53.736	<0.001*
	TC	55	55.0%	14	14.0%		
	CC	12	12.0%	2	2.0%		
Alleles	(T)	121	60.5%	182	91%	50.641	<0.001*
	(C)	79	39.5%	18	9%		

N= number of cases, % = percentage of cases, TT = thymine thymine, TC =thymine cytosine, CC= cytosine cytosine, T =thymine, C=cytosine. Significance using χ<sup>2</sup>: Chi-square test:  
\*p<0.001 (significant)

**Table 4. Comparison between all cases with obesity and healthy controls regarding their genotype distribution of eNOS gene polymorphism in (27 bp) repetition**

27bp		Patients		Control		χ <sup>2</sup>	p
		N	%	N	%		
Genotype	aa	15	15.0%	14	14.0%	0.961	0.618
	ab	63	63.0%	58	58.0%		
	bb	22	22.0%	28	28.0%		
Alleles	(a)	93	46.5%	86	43%	0.495	0.482
	(b)	107	53.5%	114	57%		

N= number of cases, % = percentage of cases, a=allele a, b= allele b  
Significance using χ<sup>2</sup>: Chi-square test.

### 3.1 Electrophoresis Result of PCR Showing Enzymatic Digestion of T786C Polymorphism of eNOS Gene

Wild type TT is found which appear at 236bp in lanes 1, 2, 4 and 5, digestion of PCR product of T786C polymorphism of eNOS gene using NgoMIV enzyme. Which digest the 236-bp fragment into 203 and 33-bp fragments (heterozygous mutated genotype TC which has 236, 203, 33 bp fragments lanes 6 only) but (homozygous mutated genotype CC is found which has 203, 33 bp fragments lanes 3, 7) by using DNA size marker 50bp.

### 3.2 Electrophoresis Result of PCR Showing PCR Amplification of Intron 4b/a (27bp) Polymorphism of eNOS Gene

PCR product of intron 4b/a polymorphism have ban size (220) bp in bb carrier homozygous lanes 2, 3, 6 and 7 and have ban size (193) in aa homozygous lanes 4 and ba carrier heterozygous which has (220,193 bp fragments lanes 1 and 5) by using DNA marker 50 bp.




Fig. 1. Enzymatic digestion of T786C polymorphism of eNOS gene




Fig. 2. PCR amplification of intron 4b/a polymorphism of eNOS gene

#### 4. DISCUSSION

Overweight and obesity are major risk factors for a number of chronic diseases, including diabetes, cardiovascular diseases and cancer. Once considered a problem only in high income countries, overweight and obesity are now dramatically on the rise in low- and middle-income countries [1].

Obesity is one of the leading preventable causes of death worldwide [17,18]. Growing evidence supports the association of diseases with NOS<sub>3</sub> haplotypes (combination of alleles in close proximity, within a DNA block). This approach may be more informative than the analysis of genetic polymorphisms one by one [19]. Haplotypes including the SNPs g.-786T>C and Glu298Asp, g- G894T and the VNTR in intron 4 affected the susceptibility to hypertension [20]. And there is association between NOS<sub>3</sub> and the susceptibility to obesity [16]. And diabetes mellitus [21].

The present study aims mainly to investigate the association of the eNOS gene polymorphism (T786 C, and 27bp) with the possibility of occurrence obesity, the study results showed that homozygous mutated TT and homozygous mutated CC genotypes, mutant T and C allele of T786C polymorphism had significant frequency among cases of obesity compared with controls. On the contrary, homozygous mutated bb and homozygous mutated aa genotypes, mutant b and a allele of 27 bp polymorphism had no significant frequency among cases of obesity compared with controls.

Souza-Costa DC et al. [16] A Brazilian study suggested that the eNOS gene polymorphism is associated with hypertension in obese children and adolescents. Further studies examining the possible interactions of eNOS haplotypes with environmental factors and other genetic markers might cause the development of obesity and its complications are warranted.

The present research exhibited a significant association of T786C with occurrence of obesity and these results in harmony with results of Josiane A. Miranda et al. [22] reported a similar association of the T786C polymorphism predispose to MetS in both obese children and adolescents.

Bressler J. et al. [23] in the United States in a study carried in four communities suggested that

interaction between incidence of obesity and NOS<sub>3</sub>.

In partial agreement with our result Baráth A et al. [24] have reported that no significant differences were seen in the case of the eNOS 4th intron 27-bp repeat polymorphism and the eNOS T-786C promoter polymorphism.

On other hand to our result, Roberta Fernanda da Silva et al.[25] a study on Brazilian patients did not demonstrate a significant difference in plasma NO<sub>2</sub> concentration blood pressure and obesity taking into account the haplotype results (-786T/C, 4b/4a, and 894G/T). In general, different levels of Training status promote different results in these variables; however, these relationships need to be studied further.

On the contrary to the present research Hela Ben Nasr et al. [26] suggested that among Tunisian patients, eNOS gene polymorphism 27pb (4b/a) was significantly associated with obesity.

Our study reported that endothelial nitric oxide gene polymorphism (T786C) is a risk factor for development of obesity

#### 5. CONCLUSIONS

The C786T polymorphism of eNOS gene was found to be significantly associated with development of obesity and T, C alleles, (CC and TT genotypes of C786T) might significantly considered genetic risk factor for development of obesity.

#### CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

#### ACKNOWLEDGEMENTS

We would like to thank the staff members of the department of diabetes and endocrine unit in

specialized medical hospital Mansoura University, Mansoura, Egypt for their sincere help and cooperation.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. WHO. World Health Organization 2015"Obesity and overweight Fact sheet N°311. WHO. January 2015. Retrieved 2 February 2016.
2. Kanazawa M, Yoshiike N, Osaka T, Numba Y, Zimmet P, Inoue S. Criteria and classification of obesity in Japan and Asia-Oceania. *Nutrition and Fitness: Obesity, the Metabolic Syndrome, Cardiovascular Disease, and Cancer. World Review of Nutrition and Dietetics.* 2005;94:1–12. DOI:10.1159/000088200. ISBN 978-3-8055-7944-5. PMID 16145245.
3. Haslam DW, James WP (October 2005). "Obesity". *Lancet* (Review). 366 (9492): 1197–209. DOI:10.1016/S0140-6736(05)67483-1. PMID 16198769.
4. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, Zitman FG. Overweight, obesity, and depression: A systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry.* 2010;67(3):220–9. DOI:10.1001/archgenpsychiatry.2010.2. PMID 20194822.
5. Yazdi FT, Clee SM, Meyre D. Obesity genetics in mouse and human: Back and forth, and back again. *Peer J.* 2015;3: e856. DOI:10.7717/peerj.856. PMC 4375971. PMID 25825681
6. Bleich S, Cutler D, Murray C, Adams A. Why is the developed world obese? *Annual Review of Public Health (Research Support).* 2008;29:273–95. DOI:10.1146/annurev.publhealth.29.020907.090954. PMID 18173389.
7. Oxford Handbook of Medical Sciences (2<sup>nd</sup> ed.). Oxford: OUP Oxford. 2011. P. 180. ISBN 9780191652295.
8. Kushner R. Treatment of the obese patient (Contemporary Endocrinology). Totowa, NJ: Humana Press. 2007;158. ISBN 978-1-59745-400-1. Retrieved 5 April 2009.
9. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, et al. Health effects of overweight and obesity in 195 Countries over 25 Years. *The New England Journal of Medicine.* 2017;377(1): 13–27. DOI:10.1056/NEJMoa1614362. PMC 5477817. PMID 28604169.
10. DiBaise JK, Zhang H, Crowell MD, Krajmalnik-Brown R, Decker GA, Rittmann BE. Gut microbiota and its possible relationship with obesity. *Mayo Clinic Proceedings (Review).* 2008;83 (4):460–9. DOI:10.4065/83.4.460. PMID 18380992.
11. Pollack A. A.M.A. Recognizes Obesity as a Disease. *New York Times.* Archived from the original on 23 June 2013.
12. Weinstock, Matthew (21 June 2013). The facts about obesity. H&HN. American Hospital Association. Retrieved 24 June 2013.
13. Lacchini R, Silva PS, Tanus-Santos JE (May). A pharmacogenetics-based approach to reduce cardiovascular mortality with the prophylactic use of statins. *Basic & Clinical Pharmacology & Toxicology.* 2010 ;106(5):357–61. DOI:10.1111/j.1742-7843.2010.00551.x. PMID 20210789.
14. Niu W, Qi Y. An updated meta-analysis of endothelial nitric oxide synthase gene: Three well-characterized polymorphisms with hypertension. *PLOS ONE.* 2011;6 (9): e24266. DOI:10.1371/journal.pone.0024266. PMC 3166328. PMID 21912683.
15. Zhang MX, Zhang C, Shen YH, Wang J, Li XN, Chen L, Zhang Y, Coselli JS, Wang XL. Effect of 27nt small RNA on endothelial nitric-oxide synthase expression. *Molecular Biology of the Cell.* 2008;19(9):3997–4005. DOI:10.1091/mbc.E07-11-1186. PMC 2526692. PMID 18614799.
16. Souza-Costa DC, Belo VA, Silva PS, Metzger IF, Lanna CM, Machado MA, Tanus-Santos JE. *International Journal of Obesity.* 2011;35:387–392.
17. Barness LA, Opitz JM, Gilbert-Barness E. Obesity: Genetic molecular, and environmental aspects. *Am J Med Genet A.* 2007;143A:3016-3034.
18. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA.* 2004;291(10):1238–45.

- DOI:10.1001/jama.291.10.1238. PMID 15010446.
19. Crawford DC, Nickerson DA. Definition and clinical importance of haplotypes. Annual Review of Medicine. 2005;56:303–20. DOI:10.1146/annurev.med.56.082103.104540. PMID 15660514.
20. Sandrim VC, Coelho EB, Nobre F, Arado GM, Lanchote VL, Tanus-Santos JE (Jun). Susceptible and protective eNOS haplotypes in hypertensive black and white subjects. Atherosclerosis. 2006;186(2): 428–32. DOI:10.1016/j.atherosclerosis.2005.08.003 PMID 16168996.
21. Jia Z, Zhang X, Kang S, Wu Y. Association of endothelial nitric oxide synthase gene polymorphisms with type 2 diabetes mellitus: A meta-analysis. Endocrine Journal. 2013;60(7):893–901. DOI:10.1507/endocrj.ej12-0463. PMID 23563728.
22. Josiane A. Miranda, Vanessa A. Belo, De'bora C. Souza-Costa, Carla MM. Lanna, Jose E. Tanus-Santos., 2012 Mol Cell Biochem DOI:10.1007/s11010-012-1456-y.
23. Andrade VL, Sertorio JTC, Eleuterio NM, Tanus-Santos JE, Fernandes KS, Sandrim VC. Elsevier J., Nitric Oxide. 2013;33:83–87.
24. Baráth A, Endreffy E, Bereczki C, Gellén B, Szücs B, Németh I, Túri S., Acta Physiol Hung. 2007;94(1-2):49-66.
25. Roberta Fernanda da Silva ID, A' tila Alexandre Trape', Thai's Amanda Reia, Riccardo Lacchini, Gustavo Henrique Oliveira-Paula, Lucas Cezar Pinheiro, Jose'Eduardo Tanus-Santos, Andre' Mourão Jacomini, Carlos Roberto Bueno Ju' nior, Anderson Saranz Zago., PloS ONE. 2018;13(10):e0206254.
26. Hela Ben Nasr, Saloua Dimassi, Refka M'hadhbi, Haithem Debbabi, Mondher Kortas, Zouhair Tabka, Karim Chahe. Functional G894T polymorphism and intron-4 VNTR variant of nitric oxide synthase (NOS3) gene are susceptibility biomarkers of obesity among Tunisians. 2016;10(4):465-75. DOI:10.1016/j.orcp.2015.04.008. Epub 2015 May 5.

© 2019 Elbaz et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/47557>*