



Super-sech Soliton Dynamics in Optical Metamaterials with Generally Parabolic Law of Nonlinearity Using Lagrangian Variational Method

Amour Ayela¹, Gaston Edah^{2*}, Camille Elloh³
and Gaétan Djossou³

¹Institut de Mathématiques et de Sciences Physiques (IMSP), Bénin.

²Département de Physique, Faculté des Sciences et Techniques (FAST), Bénin.

³Chaire Internationale de Physique Mathématique et Applications (CIPMA), Bénin.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/PSIJ/2019/v21i330109

Editor(s):

(1) Pratima Parashar Pandey, Professor, Polymer Composites, Nanotechnology Applied Sciences, College of Engineering & Technology, Integrated Institutes of Learning, Dr. APJ Abdul Kalam Technical University, Greater Noida, India.

(2) Junjie Chen, Department of Electrical Engineering, University of Texas at Arlington, USA.

(3) Christian Brosseau, Distinguished Professor, Department of Physics, Université de Bretagne Occidentale, Brest, France.

Reviewers:

(1) Marin Marin, Transilvania University of Brasov, Romania.

(2) Yahaya Shagaiya Daniel, Kaduna State University, Nigeria.

(3) Olumide Adesina, Olabisi Onabanjo University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47389>

Received: 09 December 2018

Accepted: 21 February 2019

Published: 22 March 2019

Original Research Article

Abstract

Aims/ Objectives: This paper studies the impact of the generally parabolic law of nonlinearity on the evolution of the energy of super-sech soliton dynamics.

Study Design: generally parabolic law of nonlinearity terms study.

Place and Duration of Study: Department of Physics, Faculty of Sciences and Technology (FAST), University of Abomey Calavi, Bénin. between February 2018 and January 2019.

*Corresponding author: E-mail: ggastonedah@gmail.com

Methodology: Variational approach, namely, the Lagrangian Variational Method (LVM) is presented. The different results are obtained using standard fourth order Runge-Kutta method for integration of the system of ordinary differential equation systems.

Results: Dynamics of the different parameters (amplitude, center position, pulse width, chirp, frequency and phase) has been presented with respect to propagating distance.

Conclusion: This study reveals that the generally parabolic law of nonlinearity terms don't affect the energy of the system but influence the pulse phase.

Keywords: Lagrangian approach; generally parabolic law; super-sech soliton; metamaterial.

2010 Mathematics Subject Classification: 53C25; 83C05; 57N16

1 INTRODUCTION

The metamaterial is a new type of microstructured material which has been extensively used and studied during the recent years. Metamaterials are artificial composite structures with both negative permittivity and negative permeability. They also have fascinating physical properties and spectacular uses [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. Metamaterials are an emerging technology with applications in a range of diverse areas. Metamaterials are artificially engineered materials with properties not available in natural systems such as negative permeability and permittivity, display anomalous behaviour, such as negative refraction, superlensing, backward wave propagation and reverse Doppler shifting. Consequently they are

many applications including energy harvesting, object cloaking, high data rate communications, sensors and detectors, imaging, anti-vibration, noise reduction, seismic protection and antennae [12]. Metamaterials can either be used to improve the performance of existing applications. Nowadays, it is possible to use this material as waveguide in order to optimize the data transmission. This is precisely the framework of the present research. This research aims to study the dynamics of a soliton pulse, super-sech soliton which is propagated in a metamaterial, in order to assess the impact of the generally parabolic law of nonlinearity on the pulse profile along its path in the metamaterial. The dynamics of solitons in optical metamaterials is governed by the model [13, 9, 14, 15, 16, 17, 5, 6]:

$$iq_z + aq_{tt} + b|q|^2q = i\alpha q_t + i\lambda(|q|^2q)_t + i\nu(|q|^2)_tq + \theta_1(|q|^2q)_{tt} + \theta_2|q|^2q_{tt} + \theta_3q^2q_{tt}^* \quad (1.1)$$

This equation was recently used by Douvagai et al. where they introduced an additional nonlinear term (1.2). These last ones used the complex envelope ansatz method and the F-expansion method to solve the generally nonlinear Schrödinger equation (GNLSE) with the additional parabolic law nonlinearity. Bright and dark soliton solutions are obtained [14].

$$iq_z + aq_{tt} + b(|q|^2 + \sigma|q|^4)q = i\alpha q_t + i\lambda(|q|^2q)_t + i\nu(|q|^2)_tq + \theta_1(|q|^2q)_{tt} + \theta_2|q|^2q_{tt} + \theta_3q^2q_{tt}^* \quad (1.2)$$

Recent work by Biswas et al. has taken this additional term into account [18]. Similarly, Foroutan et al. studied disturbances of the optical soliton in a metamaterial with an additional anti-cubic nonlinear term using two approaches: the extended trial equation method and the improved G'/G-expansion method. The Faroutan equation used is:

$$iq_z + aq_{tt} + (b_1|q|^{-4} + b_2|q|^2 + b_3|q|^4)q = i\alpha q_t + i\lambda(|q|^2q)_t + i\nu(|q|^2)_tq + \theta_1(|q|^2q)_{tt} + \theta_2|q|^2q_{tt} + \theta_3q^2q_{tt}^* \quad (1.3)$$

The bright, dark and singular soliton are retrieved in this research [15]. Indeed, apart from the case of the basic nonlinear Schrödinger equation containing only the terms of second dispersion and self phase modulation whose exact solution are known, the (GNLSE) family is not completely integrable and can not be solved exactly. In literature, diverse numerical and direct methods have been proposed to construct exact solutions nonlinear partial differential equations modeling the wave

propagation in various media. The partial differential equations are converted into dimensionless ordinary equations by employing suitable transformations. Among them, there are the collective variables method, the method of moments, the Lagrangian variational method (LVM), the G/G'-expansion method, the trial solution method, the extended tanh function method, the Riccati approach, the soliton ansatz method, the collocation method, the homotopy analysis method, the Keller box method, the geometric approach and so on [7, 19, 20, 21, 22, 23, 24, 25, 16, 26, 27]. In this work, we extend the model used in [14, 28] by considering the effect of higher-order parabolic law nonlinearity (1.4) and propose to solve it by Lagrangian variational method. The LVM developed by Anderson [29] is based on minimization of action. This approach consists in derive a set of ordinary differential equations of some important quantities of solitary wave. The objective of such a study would be to exhibit the contribution of these terms on the dynamics of the optical soliton. These terms appear in the metamaterial context when considered as centrosymmetric materials and high order polarization vectors are taken into account in the Maxwell equation [28]. The new equation is therefore given by (1.4) and we called it the general parabolic law nonlinearity Schrödinger equation.

$$iq_z + aq_{tt} + \sum_{k=1}^n b_k |q|^{2k} q = i\alpha q_t + i\lambda(|q|^2 q)_t + i\nu(|q|^2)_t q + \theta_1(|q|^2 q)_{tt} + \theta_2 |q|^2 q_{tt} + \theta_3 q^2 q_{tt}^* \quad (1.4)$$

In equation (1.4), the unknown or dependent variable $q = q(z, t)$ represents the wave profile, while z and t are the spatial and temporal variables respectively. The first and the second terms are the linear spatial evolution terms and the group velocity dispersion, while the third term introduces the generally parabolic law of nonlinearity, the fourth, fifth and sixth terms represent inter-modal dispersion, self steepening and the nonlinear dispersion respectively. Finally, the last three terms with θ_k for $k = 1, 2, 3$ appear in the context of metamaterials [5].

2 LAGRANGIAN VARIATIONAL METHOD

The main idea of LVM is based on extending Euler-Lagrange least-action principles to dissipative systems. LVM is used to express the generalized GNLSE in terms of fundamental parameters (collective variables). This consists in finding the Lagrangian of GNLSE, then choosing any convenient trial function f (ansatz) assumed to best approximate the behaviour of the pulse in order to derive the set of variational equations [30, 18, 31, 32, 33, 34, 35]. Let's write the (1.4) in the form:

$$iq_z + aq_{tt} + \sum_{k=1}^n b_k |q|^{2k} q = \zeta, \quad (2.1)$$

where

$$\zeta = i\alpha q_t + i\lambda(|q|^2 q)_t + i\nu(|q|^2)_t q + \theta_1(|q|^2 q)_{tt} + \theta_2 |q|^2 q_{tt} + \theta_3 q^2 q_{tt}^* \quad (2.2)$$

is considered as a perturbation term. Let's consider the equation (2.1) without perturbation term ($\zeta = 0$) and look for the solution q on the form:

$$q(z, t) = u(z, t) + iv(z, t), \quad (2.3)$$

where u and v are real functions. Substituting (2.3) in (2.1), we obtain:

$$u_z + av_{tt} + \sum_{k=1}^n b_k (u^2 + v^2)^k v = 0, \quad (2.4)$$

$$-v_z + au_{tt} + \sum_{k=1}^n b_k (u^2 + v^2)^k u = 0. \quad (2.5)$$

The equations (2.4) and (2.5) can be deduced respectively from Euler-Lagrange equations given by:

$$\frac{\partial L_0}{\partial v} - \frac{\partial}{\partial z} \left(\frac{\partial L_0}{\partial v_z} \right) - \frac{\partial}{\partial t} \left(\frac{\partial L_0}{\partial v_t} \right) = 0 \quad (2.6)$$

$$\frac{\partial L_0}{\partial u} - \frac{\partial}{\partial z} \left(\frac{\partial L_0}{\partial u_z} \right) - \frac{\partial}{\partial t} \left(\frac{\partial L_0}{\partial u_t} \right) = 0, \quad (2.7)$$

where the Lagrangian L_0 is given by:

$$L_0 = \frac{1}{2} (u_z v - v_z u) + \sum_{k=2}^n \frac{b_{k-1}}{2k} (u^2 + v^2)^k - \frac{a}{2} (u_t^2 + v_t^2). \quad (2.8)$$

When we express respectively u and v as follows: $u = \frac{1}{2}(q + q^*)$; $v = \frac{i}{2}(q^* - q)$, the Lagrangian L_0 can be rewritten as follows:

$$L_0 = \frac{i}{4} (q_z q^* - q_z^* q) + \sum_{k=2}^n \frac{b_{k-1}}{2k} |q|^{2k} - \frac{a}{2} |q_t|^2. \quad (2.9)$$

The averaged Lagrangian of equation the without right hand side is defined as:

$$L = \int_{-\infty}^{+\infty} L_0 dt. \quad (2.10)$$

Then

$$L = \int_{-\infty}^{+\infty} \left[\frac{i}{4} (q_z q^* - q_z^* q) + \sum_{k=2}^n \frac{b_{k-1}}{2k} |q|^{2k} - \frac{a}{2} |q_t|^2 \right] dt. \quad (2.11)$$

3 SUPER-SECH PARAMETER DYNAMICS

The ansatz function f that we assume in this paper is the super sech soliton [5]:

$$f = X_1 \operatorname{sech}^m \left[\frac{t - X_2}{X_3} \right] \exp \left[i \left(\frac{X_4}{2} (t - X_2)^2 + X_5 (t - X_2) + X_6 \right) \right]; \quad (3.1)$$

where X_1 represents the amplitude of the pulse, X_2 the temporal position, X_3 the width, X_4 the chirp, X_5 the frequency and X_6 the phase. m is the parameter of the super-sech. In this paper, m is set equal to 2. Substituting $q = f$ in (2.11), we obtain:

$$L = L_1 + \sum_{k=2}^n \frac{b_{k-1}}{2k} \int_{-\infty}^{+\infty} \operatorname{sech}^{2k} \left[\frac{t - X_2}{X_3} \right] dt, \quad (3.2)$$

where

$$\begin{aligned} L_1 = & \frac{2}{3} X_1^2 X_3 X_5 \dot{X}_2 + \frac{6 - \pi^2}{36} X_1^2 X_3^3 \dot{X}_4 - \frac{2}{3} X_1^2 X_3 \dot{X}_6 + \frac{8}{35} b_1 X_1^4 X_3 \\ & - \frac{a}{90} \frac{X_1^2}{X_3} (48 + 60 X_3^2 X_5^2 - (30 - 5\pi^2) X_3^4 X_4^2). \end{aligned} \quad (3.3)$$

So for $n = 6$, the average Lagrangian is:

$$\begin{aligned} L = & \frac{2}{3} X_1^2 X_3 X_5 \dot{X}_2 + \frac{6 - \pi^2}{36} X_1^2 X_3^3 \dot{X}_4 - \frac{2}{3} X_1^2 X_3 \dot{X}_6 + \frac{8}{35} b_1 X_1^4 X_3 \\ & - \frac{a}{90} \frac{X_1^2}{X_3} (48 + 60 X_3^2 X_5^2 - (30 - 5\pi^2) X_3^4 X_4^2) + \frac{256}{2079} b_2 X_1^6 X_3 \\ & + \frac{512}{6435} b_3 X_1^8 X_3 + \frac{71}{1251} b_4 X_1^{10} X_3 + \frac{127}{2948} b_5 X_1^{12} X_3; \end{aligned} \quad (3.4)$$

\dot{X}_j , ($j = 1, 2, 3, 4, 5, 6$) stands for derivative of X_j with respect to z . Now, let's come back to the full equation (2.1) where the term of right-hand side ζ is non zero. When one applies the Euler-Lagrange equations to (1.4), the variational equations are written as:

$$\frac{\partial L}{\partial X_j(z)} - \frac{d}{dz} \frac{\partial L}{\partial \dot{X}_j(z)} = \int_{-\infty}^{+\infty} \zeta f_{X_j}^* dt + c.c. \quad (3.5)$$

Substituting the expression of the average Lagrangian given in equation (2.11) and the ansatz function f in ζ , then performing the integration of the right-hand side of (3.5), we obtain the following set of variational equations:

$$\begin{aligned} \dot{X}_1 &= -aX_1X_4 + \frac{2X_1^3X_4}{35(\pi^2-6)} ((24\pi^2-235)\theta_1 + (24\pi^2-157)(\theta_2-\theta_3)), \\ \dot{X}_2 &= 2aX_5-2\alpha - \frac{24}{35} ((3\lambda+2\nu)X_1^2 + (6\theta_1+2\theta_2-2\theta_3)X_1^2X_5), \\ \dot{X}_3 &= 2aX_3X_4 - \frac{4}{35(-6+\pi^2)} ((-307+36\pi^2)\theta_1 + (-85+12\pi^2)(\theta_2-\theta_3))X_1^2X_3X_4, \\ \dot{X}_4 &= -2aX_4^2 + \frac{672a}{35(\pi^2-6)X_3^4} - \frac{1}{(\pi^2-6)X_3^2} (\frac{144b_1}{35}X_1^2 + \frac{1024b_2}{231}X_1^4 + \frac{3072b_3}{715}X_1^6 \\ &+ \frac{568b_4}{139}X_1^8 + \frac{1648b_5}{425}X_1^8 + \frac{288}{35}\lambda X_1^2X_5) + \frac{4}{175} \frac{X_1^2}{X_3^4} ((30\pi^2-245)X_3^4X_4^2 - 360X_3^2X_5^2 - 3168)\theta_1 \\ &+ \frac{4}{175} \frac{X_1^2}{X_3^4} ((30\pi^2-245)X_3^4X_4^2 - 360X_3^2X_5^2 - 864)(\theta_2+\theta_3), \\ \dot{X}_5 &= -\frac{2(X_1X_4+2X_5)aX_5}{X_1} + \frac{4\alpha X_5}{X_1} - \frac{4}{35} \frac{X_1X_5(24\pi^2X_5-13X_1X_4-144X_5)}{\pi^2-6} \\ &+ \frac{4}{35} \frac{X_1X_5(72\pi^2X_5-91X_1X_4-432X_5)\theta_1}{\pi^2-6} - \frac{48}{35} (X_1X_4-3X_5)\lambda X_1 \\ &+ \frac{4}{35} \frac{X_1X_5(24\pi^2X_1X_4+24\pi^2X_5-157X_1X_4-144X_5)\theta_2}{\pi^2-6} - \frac{48}{35} (X_1X_4-2X_5)\nu X_1, \\ \dot{X}_6 &= \frac{1}{35} \frac{35aX_3^2X_5^2-56a}{X_3^2} + \frac{30}{35}b_1X_1^2 + \frac{512}{693}b_2X_1^4 + \frac{896}{1365}b_3X_1^6 + \frac{497}{834}b_4X_1^8 \\ &+ \frac{309}{500}b_5X_1^{10} + \frac{1}{350} \frac{X_1^2}{X_3^2} ((30\pi^2-245)X_3^4X_4^2 - 840X_3^2X_5^2 + 928)\theta_1 \\ &- \frac{2}{35} (6\lambda+24\nu)X_1^2X_5 + \frac{1}{350} \frac{X_1^2}{X_3^2} ((30\pi^2-245)X_3^4X_4^2 - 120X_3^2X_5^2 + 928)(\theta_2+\theta_3). \end{aligned} \quad (3.6)$$

4 RESULTS AND DISCUSSION

The numerical study of the evolution of the different parameters of the super-sech soliton momentum has been made in order to appreciate the impact of the generally parabolic law nonlinearity terms on the dynamics of such an pulse in a metamaterials. The different results are obtained using standard fourth order Runge-Kutta method for integration of the system of ordinary differential equation systems. The dynamics of the system have been presented in Figure2 for the following parameter values: $a = 0.1$, $b_1 = -20$, $\alpha = -0.25$, $\lambda = 0.1$, $\nu = 0.1$, $\theta_1 = -0.01$, $\theta_2 = -0.02$, $\theta_3 = -0.3$, $b_2 = 0.001$, $b_3 = 0.1$, $b_4 = 0.1$, $b_5 = 2$.

The analysis of this curve shows that the amplitude, the pulse width, the chirp and the frequency slip vary periodically as a function of z . Indeed, it should be noted that the choice of the initial condition is

of paramount importance for such a study. These parameters have been chosen so that the super-sech soliton propagates itself without attenuation. The variational equations $\dot{X}_1, \dot{X}_2, \dot{X}_3$ obtained are identical to those of Veljkovic et al. [5]. This explains the resemblance of the representative curves of the amplitude, the center position and the pulse width. The terms of high order added to the equation don't influence the evolution of these parameters ($\dot{X}_1, \dot{X}_2, \dot{X}_3$). On the other hand the variational equations: $\dot{X}_4, \dot{X}_5, \dot{X}_6$ are functions of the terms of high order introduced and show dissimilarities. The different terms $b_k, k = 1, \dots, 6$, rather influence the parameters of the pulse phase. This confirms the absence of these terms in the expression that describes the variation of the energy (4.3). A comparison of our results to those obtained in literature [5, 10] gave excellent agreement. As the pulse width propagates, the amplitude X_1 , the pulse width X_3 , the frequency X_5 and the chirp X_4 vary periodically.

A particular attention has been drawn on the energy of the system. The energy is defined as:

$$L = \int_{-\infty}^{+\infty} |q|^2 dt. \quad (4.1)$$

In the case of the super-sech soliton, one has:

$$E = \frac{4X_1^2 X_3}{3}. \quad (4.2)$$

The evolution of the energy is given by:

$$\frac{dE}{dz} = \left[\theta_1 \left(\frac{192 - 32\pi^2}{35(-6 + \pi^2)} \right) + (\theta_2 - \theta_3) \left(\frac{-976 + 128\pi^2}{35(-6 + \pi^2)} \right) \right] X_1^4 X_3 X_4. \quad (4.3)$$




Figure 1: Variation of energy




Figure 2: Variation of normalized pulse parameters(X_1 -soliton amplitude, X_2 -center position of the soliton, X_3 -pulse width, X_4 -soliton chirp, X_5 -soliton frequency, X_6 -soliton phase) with propagation distance

5 CONCLUSION

This paper presents lagrangian variational approach for super sech soliton dynamics in optical metamaterials. The optical soliton dynamics is governed by the generalized nonlinear Schrödinger equation including generally parabolic law of nonlinearity. This equation is solved by lagrangian approach where a six paramater (amplitude, center position, pulse width, chirp, frequency and phase) super-sech soliton test function has been used to approximate the exact solution. Numerical simulations have made it to represent these parameters graphically as a function of the propagation distance. This study reveals that the generally parabolic law of nonlinearity terms don't affect the energy of the system, but affect the pulse phase. Finally, the analysis of these results revealed that the choice of the initial condition is crucial for such a study. A comparison with other results gave excellent agreement. This work could be proposed in telecommunication

to optimize the transmission of information. The results with those additional laws of nonlinearity will be reported in future.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

References

- [1] Veselago VG. The electrodynamics of substances with simultaneously negative values of ϵ and μ . Sov. Phys. Usp. 1968;10:509-514.
- [2] Pendry JB. Negative refraction makes a perfect lens. Physics Review Letter. 2000;18:3966-3969.
- [3] Shalaev, et al. Optical negative-index metamaterials. Nat. Photon. 2007;13:41-48.

- [4] Zharova, et al. Nonlinear transmission and spatio-temporal solitons in metamaterials with negative refraction. *Optics Express*. 2005;13:1291-1298.
- [5] Veljković, et al. Super-sech soliton dynamics in optical metamaterials using collective variables. *Electronics and Energetics, Sov. Phys. Usp.* 2017; 1:39-48.
- [6] Veljković, et al. Super-gaussian soliton in optical metamaterials using collective variables. *Journal of Computational and Theoretical Nanoscience*. 2015;12:5119-5124.
- [7] Biswas, et al. Cubic-quartic optical solitons in kerr and power law media. *Optik*. 2017;144:357-362.
- [8] Biswas, et al. Singular soliton in optical metamaterials by ansatz method and simplest equation approach. *Optik*. 2014;61:1550-1555.
- [9] Biswas, et al. Bright and dark solitons in optical metamaterials. *Optik*. 2014;125:3299-3302.
- [10] Green, et al. Dynamics of gaussian optical solitons by collective variables method. *Applied Mathematics and Information Sciences*. 2008;2:259-273.
- [11] Solymar E. Shamonina Waves in metamaterials. Oxford University Press; 2009.
- [12] Steve M, et al. A state of the art review of smart materials. *Review of metamaterials in the UK*; 2015.
- [13] Agrawal GP. Nonlinear fiber optics. Academic Press, Boston; 1989.
- [14] Douvagai, et al. Electromagnetic wave solitons in optical metamaterials. *Optik*. 2017;140:735-742. Edah, et al. Pulse propagation in a non linear medium. *Open Phys*. 2017;13:151-156.
- [15] Faroutan, et al. Solitons in optical metamaterials with anti-cubic law of nonlinearity by ETEM and IGEM. *Journal of the European Optical Society*. 2018;14:16.
- [16] Zhou, et al. Analytical study of thirring optical solitons with parabolic law nonlinearity with spatio-temporal dispersion *Optik*. 2017;142:73-76.
- [17] Zhou, et al. Soliton in optical metamaterials with parabolic law nonlinearity and spatio-temporal dispersion. *Journal of Optoelectronics and Advanced Materials*. 2014;16:1221-1225.
- [18] Biswas, et al. Conservation laws for perturbed solitons in optical metamaterials. *Physics*. 2018;8: 898-902.
- [19] Cai W. Shalaev Optical metamaterials: fundamentals and application. Springer, New York; 2010.
- [20] Daniel Y, et al. Impact of thermal radiation on electrical MHD flow of nanofluid over nonlinear stretching sheet with variable thickness. *Alexandria Engineering Journal*. 2017;57(3): 2187-2197.
- [21] Daniel Y, et al. Entropy analysis in electrical magnetohydrodynamic (MHD) flow of nanofluid with effects of thermal radiation, viscous dissipation, and chemical radiation. *Theoretical and Applied Mechanics Letters*. 2017;7(4):235-24.
- [22] Daniel Y, et al. Double stratification effects on unsteady electrical MHD mixed convection flow of nanofluid with viscous dissipation and joule heating. *Journal of Applied Research and Technology*. 2017;15(5):464-476.
- [23] Daniel Y, et al. Effects of buoyancy and thermal radiation on MHD flow over a stretching porous sheet using homotopy analysis method. *Alexandria Engineering Journal*. 2015;54(3):705-712.
- [24] Fujioka, et al. Chaotic soliton in the quadratic-cubic nonlinear under nonlinearity management Schrödinger equation. *Chaos*. 2011;21:033120.
- [25] Saha M, Sarma AK. Modulation instability in nonlinear metamaterials induced by cubic-quintic nonlinearities and higher order dispersive effects. *Optik*. 2013;291:321-325.
- [26] Zhou, et al. Perturbation theory and optical soliton cooling with anti-cubic nonlinearity. *Optik*. 2017;142:73:76.
- [27] Zhou, et al. Optical solitons in birefringent fibers with parabolic law nonlinearity. *Optics*. 2014;44:399-409.

- [28] Houria, et al. Chirped soliton solutions for the generalized nonlinear Schrödinger equation with polynomial nonlinearity and non-kerr terms of arbitrary order. Journal of Optics. 2016;18:1-9.
- [29] Anderson D. Variational approach to nonlinear pulse propagation in optical fibers. Physics Review. 1983;27:3135-3145.
- [30] Adrian A, Nail A. Comparison of Lagrangian approach and method of moments for reducing dimensionality of soliton dynamical systems. Chaos. 2008;18:0331291-0331298.
- [31] Biswas, et al. Optical soliton perturbation by semi-inverse variational principle. Optik. 2017;143: 131-134.
- [32] Cheng, et al. Dark soliton solutions to the Schrödinger equation for ultra-short propagation in metamaterials. Journal Nonlinear Optics. 2009;18:271-284.
- [33] Faroutan, et al. Solitons in optical metamaterials with anti-cubic law of nonlinearity by ETEM and IGEM. Journal of the European Optical Society. 2018;14:16.
- [34] Moubissi, et al. Non lagrangian collective variable approach for optical soliton in fibers. Journal of Physics A. 2001;34:129-136.
- [35] Nakkeeran, et al. Generalized projection operator method to derive the pulse parameters equations for the nonlinear Schrödinger equation. Optics Communications. 2005;244:377382.

©2019 Ayela et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47407>