



Morphological Exploration of Lignicolous Macrofungi: *Xylaria*

**Priya John ^{a*}, Kotramma Addangadi ^a, Anjali Suansia ^a
and Megha Vyas ^a**

^a Department of Plant Pathology, N. M. College of Agriculture, Navsari Agricultural University, Navsari, 396450, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/mrji/2024/v34i81467>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119478>

Received: 05/05/2024

Accepted: 08/07/2024

Published: 12/07/2024

Original Research Article

ABSTRACT

Xylaria is a large and first described genus of *Xylariaceae* which is generally recognized as one of the most diverse and relatively well-known largest family of Ascomycota. The fruiting bodies of *Xylaria* species were collected from Navsari, Gujarat during rainy season. All the specimens were examined for their morphological characters based on shape, size and colour of stromata followed by detailed microscopic examination of perithecia and ascospores with light microscope and were identified. From the total collection seven species were designated based on detailed macro- and micro-morphological characteristics. Among the different collected samples five were identified upto species level and two upto genus level. These were *Xylariahypoxylon*, *X. longipes*, *X. polymorpha*, *X. filiformis* and *X. carpophila*. The diversity of *Xylaria* species in this region invokes and unveils opportunities for exploring rich source of bioactive compounds such as alkaloids, terpenoids, cytochalasins, cell-degrading enzymes, exopolysaccharides, xylaramide, xylarinxyloketal, xylaranic

*Corresponding author: E-mail: pjohn@nau.in;

acids and helvonic acid which exhibited antifungal and antibacterial activities. So, exploitation of *Xylaria* fungi creates new opportunities to develop bio-based commercial products to combat global crop and human pathogens.

Keywords: *Xylaria*; stromata; morphology; pileus; ascospore; stipe.

1. INTRODUCTION

"*Xylaria* Hill ex Schrank (1789) is the largest and first described genus of *Xylariaceae* which is generally identified as one of the most diverse and comparatively well-known largest family of ascomycota and widely distributed throughout the globe evidenced by reports from different regions with broad ecological diversity" [1]. *Xylaria* differs from other genera in *Xylariaceae* by producing upright, barrel shaped to clavate, multi-peritheciate stromata [2]. At present, about 300 estimated species are thought to have a place in *Xylaria* [3]. They are mostly saprophytic, may be pathogenic Proffer [4], Hartman et al. [5] and endophytic in nature and appeared to exist on decayed wood, dung, litter and soil Davis et al. [6], Okane et al. [7]. Kshirsagar et al. [8], Ramesh et al. [9] and Hande & Hiwarale [10] reported "more than 10 *Xylaria* species in western and northern states of India". "However, existing awareness and research on these fungi in Gujarat state (India) is at their primary stage with only few reports" [11-13].

"Gujarat shares unique diversity of various flora and fauna because Gujarat is known to have varying climatic conditions like moist deciduous (ending part of Western Ghat) to extreme desert conditions (Rann of Kutchh), Savanna type grassland, arid to semiarid conditions in northern part of the state and also has the longest seacoast in country. Angiosperm flora of Gujarat documented and studied thoroughly but mycoflora has always been neglected by the botanists" [13]. "The reports on fungal diversity of Gujarat could be found in the literature but they are only restricted to pathogenic fungi associated to agricultural crops or human beings" Bhavsar et al. [14], Assudani [15], Dhingani et al. [16], Nagadesi and Arya [12]. In the present study we attempted to document the diversity of *Xylaria* which is a rich source of various secondary metabolites due to endophytic nature of many species Hacıoglu et al. [17], Nilza [18], Fan et al. [19]. The diversity of *Xylaria* species in this region invokes and unveils the opportunities for exploring rich source of bioactive compounds such as alkaloids, terpenoids, cytochalasins, exopolysaccharides, cell-degrading enzymes, xylamide, xylarin,

xyloketals, xylaric acids and helvonic acid which exhibited antifungal, antibacterial and also antiviral activities. Thus, it will be a new opportunity to develop bio-based commercial products to combat global crop and human pathogens. "It is also valued for various lignocellulosic enzymes that have a wide application in pharmaceutical, in textile industry, paper and pulp industry etc". Soto et al. [20].

2. MATERIALS AND METHODS

The survey was conducted at the different locations of Navsari, Gujarat for the collection of naturally growing *Xylaria* during the rainy seasons of 2017 and 2018 when the temperature was around 28- 30°C and relative humidity was more than 85%. The sites of collection were soil, open lands, farm lands, roadside, research field etc. The fruiting bodies were kept in the sterile polyethylene bags labeled with the location, name of the host substrate and date of collection. These were carried to the laboratory. All the fruiting bodies were micro-photographed. All the specimens were examined for their morphological characters based on shape, size and colours of pileus and length and diameter of stipe followed by detailed microscopic examination of perithecia and ascus with light microscope and were identified by referring published reports [21,22,8].

3. RESULTS

From the total collection seven species were identified. Among which five were identified upto species level and two upto genus level. These were *X. hypoxylon*, *X. longipes*, *X. polymorpha*, *X. filiformis* and *X. carpophila*.

***Xylariacarpophila*:** The pileus was long, slender, rough texture, black, 1.6-3.8mm width, unbranched and the tip was pointed. Stipe was black in colour with 15.67cm long and diameter was 0.5cm. Perithecia were barrel shaped with papillate ostioles. Asci 8 spored. Ascospores pale yellow to dull white and 8-10 × 4-5µm in size.

Host: on humus rich soil and leaf litter.

Table 1. Characteristics of *Xylaria* species

Sr. No.	<i>Xylaria</i> sp.	Pileus Colour	Pileus Diameter (mm)	Pileus Shape	Stipe Length (mm)	Stipe Diameter (mm)	Stipe Colour	Size of Spore (µm)	Host
1	<i>X. carpophila</i>	Black	1.6-3.8	pointed	15.67	0.5	black	8-10 X 4-5	Soil
2	<i>X. filiformis</i>	White	1-1.2	Hairy to slender	38	1.9	black	9.14 X 4-6	Royal palm seed
3	<i>X. hypoxylon</i>	White or ash white	1-1.9	Finger like (2-5 pointed)	19.67 - 53	0.8- 3.82	black	3.5-7.5 X 2.5 – 5.5	Wood chips, soil
4	<i>X. longipes</i>	Smoky white to greyish white	7-11	Culb shaped	22-62	1.9 – 2.87	black	8-11 X 3-5	Soil
5	<i>X. polymorpha</i>	Black	1.8–2.9	Finger like (single)	12.67 - 13	1 – 1.3	black	8-11 X 4-6	Wood chips

***Xylariafiliformis*:** Pileus was simple, long, slender, 2 mm width, filiform and white smooth surfaced, which become wrinkled at maturity with very thin apical tip and distinct. The stipe was 38cm long, smooth, black in colour and diameter was 1.9cm. Perithecia were solitary with papillate ostioles. Asci are cylindrical. Ascospores are black and 9-14 x 4-6µm in size,

Host: seeds of royal palm tree.

***Xylariahypoxylon*:** Pileus was slender, finger like (2-5 pointed) shaped, white or ashy white in colour with smooth surface, 1-1.9mm in diameter. Stipe was 19.67-53cm length x 0.8-3.82cm width, smooth and black in colour. Perithecia were barrel shaped with papillate ostioles. Asci cylindrical and 8 spored. Ascospores black and 3.5-7.5 x 2.5-5.5µm in size.

Host: decaying wood log/sticks/chips, soil, dry fruits shells of *Cassia fistula*

***Xylarialongipes*:** Fruiting bodies were found in single or in-group. Pileus was club shaped, tough, 7-19mm in diameter and lengthened to round apex. Surface was smoky white to greyish white and turns black, cracky and scaly with age. Stipe was 22-62cm length x1.9-2.87cm width and black in colour. Perithecia with papillate ostiole. Asci 8 spored. Ascospores brown to black and 8-11 x 3-5µm in size.

Host: soil, decaying woodlog /sticks.

***Xylaria polymorpha*:** Fruiting body commonly grows on the decaying wood. Pileus was finger shaped. The tip of the fruiting body is rounded and white in colour when young and turns to black eventually with the age. Stipe was short (12.67-13cm), cylindrical and black in colour. Surface was rough and wrinkled at maturity. Asci cylindrical with 8 spores. Ascospores are 8-11 x 4-6µm in size.

Host: on unknown wood logs, chips

4. DISCUSSION

Few studies on fungal diversity of Gujarat have been carried out earlier [12,23] studied 14 species of genus *Xylaria* which were collected from different districts of Gujarat. Species mentioned in this paper have already been documented in the previous studies while existence of these five species of genera *Xylaria* in Navsari is being reported for the first time. Current study is preliminary but very essential contribution to fungal diversity of Gujarat state and further studies are needed. More species are expected from this genus and therefore; further studies are warranted to investigate the diversity of macro-fungi occurring within the state by using morphological and microscopic observations.



Fig. 1. A-*Xylariacarpophila*, B-*Xylaria filiformis*, C to E-*Xylaria ypoxylon*, F&G-*Xylaria longipes*, H&I - *Xylaria polymorpha*, J to L-Perithecia, M&N-Ascus with ascospores

5. CONCLUSION

Total ten species were collected and morphological and microscopic examinations of the specimens enabled the identification two samples upto species level and five upto genus level. These were *Xylaria hypoxylon*, *X. longipes*, *X. polymorpha*, *X. filiformis* and *X. carpophila*. The diversity of *Xylaria* species in this region invokes and unveils opportunities for exploring rich source of bioactive compounds such as alkaloids, terpenoids, cytochalasins, cell-degrading enzymes, exopolysaccharides, xylaramide, xylarinxyloketals, xylaranic acids and helvonic acid which exhibited antifungal and antibacterial activities. So, exploitation of *Xylaria* fungi creates new opportunities to develop bio-based commercial products to combat global crop and human pathogens.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lee Y, Han SS, Jeong IS. Taxonomical characteristics of *Xylaria* spp. collected from Malaysia. *Mycobiol.* 2002;30(4):193-196.
2. Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EHC et al. Families of Sordariomycetes. *Fungal Divers.* 2016; 79:1-317.
3. Kirk PF, Cannon PF, Minter DW. And Stalpers JA. *Dictionary of the fungi*, 10th edn. CABI, Egham CABI Bioscience, CBS, Landcare Research; 2008.
4. Proffer TJ. *Xylaria* root rot of urban trees caused by *Xylaria polymorpha*, *PI Dis.* 1988;72:79.
5. Hartman J, Beale J, Bachi P. Root and collar rots of tree fruits. uk cooperative extension service, plant pathology fact sheet. Agriculture College, University Kentucky (PPFS-FR-T-10); 2008.
6. Davis EC, Franklin JB, Shaw AJ. And Vilgalys R. Endophytic *Xylaria* (Xylariaceae) among liverworts and angiosperms: Phylogenetics, evolution and symbiosis. *American J Bot.* 2003;90(11): 1661-1667.
7. Okane I, Srikitikulchai P, Toyama K, Laessoe T, Sivichai S, Nigel H, Nakagiri A, Potacharoen W, Suzuki K. Study of endophytic Xylariaceae in Thailand: diversity and taxonomy inferred from rDNA sequence analyses with saprobes forming fruit bodies in the field. *Mycosci.* 2008;49: 359-372.
8. Kshirsagar AS, Rhatwal SM, Gandhe RV. The genus *Xylaria* from Maharashtra, India. *Indian Phytopath.* 2009;62(1):54-63.
9. Ramesh V, Thalavaipandian A, Karunakaran C, Rajendran A. Identification and comparison of *Xylaria curta* and *Xylaria* sp. from Western Ghats-Courtallum Hills, India. *Mycosphere* 2012;3:607-615.
10. Hande DV, Hiwarale SV. Diversity of *Xylaria* species from Amravati region, Amravati, MS, India. *Inter Res J Biol Sci.* 2003;2(1):67-69.
11. Korat C, Chopada G, John P. Studies on biodiversity of fleshy fungi in Navsari (South Gujarat), India. *Inter J Biodiversity Conservation.* 2013;5(8):508-514.
12. Nagadesi PK, Arya A. New records of lignicolous fungi from Ratanmahal Wildlife Sanctuary, Gujarat, India. *Inter Letters Natural Sci.* 2014;8:23-29.
13. Rajput KS, Koyani RD, Patel HR, Vasava AM, Patel RS, Patel AD, Singh AP. A preliminary checklist of fungi of Gujarat State, India. *CurRes Environ Appl Mycol.* 2015;5(4):285-306.
14. Bhavsar HK, Modi DJ, Sood NK, Shah H. A study of superficial mycoses with clinical mycological profile in tertiary care hospital in Ahmedabad, Gujarat. *NJ Med Res.* 2012;2(2):160-164.
15. Assudani HJ, Pandya JM, Sarvan RR, Sapre AM, Gupta AR, Mehta SJ. Etiological diagnosis of microbial keratitis in a tertiary care hospital in Gujarat. *NJ Med Res.* 2013;3:60-62.
16. Dhingani JC, Solanki KU, Kansara SS. Management of root rot disease [*Macrophomina phaseolina* (Tassi.) Goid] of chickpea through botanicals and oil cakes. *The Bioscan.* 2013;8(3): 739-742.
17. Hacıoglu N, Akata I, Dulger B. Antimicrobial potential of *Xylaria polymorpha* (Pers.) Grev. *African J Microbiol Res.* 2011;5(6):728-730.

18. Nilza 2013. Screening of endophytic fungi Xylaria for antibacterial and antifungal activity. PhD Thesis, Patiala, Punjab, India: Thapar University; 2013.
19. Fan NW, Chang HS, Cheng MJ, Hsieh SY, Liu TW, Chen IS. Secondary metabolites from the endophytic fungus *Xylariacubensis*. Helvetica Chemica Acta. 2014;97(12):1689-1699.
20. Soto GG, Medina-Gonzalez GE, Trevino-Ramirez JE, Hernandez-Luna CE. Native macrofungi that produce lignin-modifying enzymes, cellulases, and xylanases with potential biotechnological applications. Bioresources. 2015;10(4):6676-6689.
21. Rogers JD. The xylariaceae: Systematic, biological and evolutionary aspects. Mycologia. 1979;71:1-41.
22. Rogers JD, Ju YM, Lehmann J. Some *Xylaria* species on termite nests. Mycologia. 2005;97:914-923.
23. Koyani RD, Patel HR, Vasava AM, Rajput, KS. Xylariaceae: Overview and addition to fungal diversity of Gujarat state. Studies Fungi. 2016;1(1):69-79.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/119478>