



# Influence of Colored Shade Nets and Media Types on the Development of *Rhoeo discolor* (Boat Lily)

Udaya T. V. <sup>a\*</sup>, Sashikala Beura <sup>a</sup> and Pratheeksha, C. T. <sup>b</sup>

<sup>a</sup> Department of Floriculture and Landscape Architecture, Odisha University of Agriculture and Technology, Bhubaneswar, India.

<sup>b</sup> Department of Floriculture and Landscape Architecture, University of Horticultural Sciences, Bagalkot, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81229>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121218>

Original Research Article

Received: 01/06/2024

Accepted: 03/08/2024

Published: 06/08/2024

## ABSTRACT

Study was carried out Department of Floriculture and Landscaping, College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar, Odisha during 2018-19. The experiment was designed to determine the optimal conditions for enhancing plant growth, focusing on the effects of different colored shade nets (green, white, red and black) with open light and various media mixture combinations (soil, coco peat, and FYM). Key growth parameters such as plant height, plant spread, leaf number, leaf area, chlorophyll content, and number of suckers were measured over a specified growth period. Results indicated that boat lily grown under red shade net and pots filled with soil, FYM and cocopeat media mixture in 1:1:1 (v/v) proportion performed best in all the characters. It has significantly enhanced plant height (22.11 cm), number of leaves per plant (15.89) and leaf area (110.42 cm<sup>2</sup>), plant spread in E-W direction (28.78 cm), plant spread in N-S direction (28.78 cm) and leaf thickness (1.32 mm).

\*Corresponding author: E-mail: [udayflori@gmail.com](mailto:udayflori@gmail.com);

**Cite as:** T. V., Udaya, Sashikala Beura, and Pratheeksha, C. T. 2024. "Influence of Colored Shade Nets and Media Types on the Development of *Rhoeo Discolor* (Boat Lily)". *Journal of Advances in Biology & Biotechnology* 27 (8):1084-92. <https://doi.org/10.9734/jabb/2024/v27i81229>.

**Keywords:** Photo selective; shade nets; *Rhoeo discolor*; plant height; cocopeat; FYM; chlorophyll.

## 1. INTRODUCTION

Ornamental plants are grown primarily for their aesthetic appeal, which is influenced by various environmental factors such as light, temperature, humidity, and growing media. Shade nets are commonly used in ornamental plant production to provide physical protection and modify the light environment [1,2]. Most pot plants have natural habitats that differ significantly from the conditions under which they are commercially grown [3].

The use of colored shade nets has emerged as a valuable tool for horticulturists to manipulate the microclimate and light conditions [2,1]. These differently colored nets selectively filter specific light wavelengths, which can impact plant growth and development in various ways. For instance, the selective filtering of light by colored shade nets can have differing effects on plant growth and morphology - red and yellow nets have been observed to promote vegetative growth, while blue nets may inhibit it, potentially influencing leaf characteristics and nutrient accumulation [3].

In addition to altering light quality, the use of colored shade nets can also impact temperature and humidity levels within the growing environment. For example, black nets can increase humidity and lower canopy temperature, which may benefit crops but also potentially encourage pest infestations [3]. Overall, the use of colored shade nets can contribute to more sustainable horticultural practices by enhancing soil moisture retention and improving water use efficiency [3].

The coloured shade cloths are manufactured in the following colours: blue, grey, pearl, red, white, and yellow. The blue shade cloth is designed to absorb ultraviolet (UV), red, and far-red; enriching the blue spectral region. The red shade cloth absorbs UV, blue, and green and enriches the red and farred spectral region. The yellow shade cloth is designed to reduce the UV and blue; enriching the green, yellow, red, and far-red wavelengths. The white shade cloth absorbs UV and enriches the wavelengths blue, green, yellow, red, and far-red. The pearl and grey shade cloths do not enrich or absorb the different wavelengths, but the pearl is designed to scatter the light to a greater extent than the other shade cloth colours mentioned Traditional black shade cloth does not scatter light [4]. By

selectively filtering light wavelengths, these nets alter the intensity and quality of light reaching the plants, which in turn influences crucial physiological processes such as photosynthesis and nutrient accumulation. Additionally, shade nets play a role in regulating temperature and humidity levels within the growing environment, with potential effects on pest dynamics and water use efficiency. Understanding the specific impacts of each colored shade net is essential for optimizing growing conditions and promoting sustainable horticultural practices [5].

Growing media are vital components of plants in containers as they determine the physical support, accessibility to water and nutrients for the plants. Potting mix is the most common form of growing media available from nurseries, and it usually contains a mixture of inorganic and organic substrates. Inorganic materials may be directly mined, such as sand or originate from processed materials such as perlite, vermiculite and polystyrene. Organic material added to the potting mix because of its low bulk density and its ability to increase the fine structural component of media. Thus, water available to plant is enhanced. Currently, organic materials used in nurseries include cocopeat, farmyard manure, peat, sawdust, wood bark, Vermicompost and sphagnum moss, etc.

Cocopeat is a versatile natural fiber extracted from monocarp tissue or husk of the coconut. The husk contains 20% to 30% fiber of varying length and holds 8-9 times its weight in water. Can be reused for up to 4 years. The properties of Cocopeat make it resistant to bacterial and fungal growth, easy to handle and excellent oxygenation properties, which is vital for healthy root development. Environment-friendly [6].

Farmyard manure (FYM) is composed of dung and urine of farm animals along with litter and leftover materials from roughages or greens fed to the farm animals. Usually, cow dung and urine of animals along with their litter and waste fed or collected and placed in trench daily and when filled in, it is covered with the field. It decomposes in two-three months when it is considered useable [6]. So readily available and economically feasible potting mixture may efficient in the vertical garden establishment.

Hence, the present experiment has been planned out to investigate the effect of different

coloured shade net and potting media mixture contributing to growth and development of *Rhoeo discolor*.

are dark to metallic green with white streaks above, with glossy purple underneath and also grown as ornamental houseplants.

## 2. MATERIALS AND METHODS

The experiment was laid out at Department of Floriculture and Landscaping, College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar, Odisha during the year 2018-2019 in a completely randomized block design (CRD) with three replications.

### 2.1 Plant Species

***Tradescantia spathacea***: synonym *Rhoeo discolor*, boat lily or Moses-in-the-cradle, is a herb in the Commelinaceae family. Native to Belize, Guatemala, and southern Mexico but widely cultivated as an ornamental and naturalized in parts of Florida, Texas, Hawaii, and various oceanic islands. It has fleshy rhizomes and rosettes of waxy lance-shaped leaves. Leaves

**Net characteristics**: The present experiment deals with the microenvironment created by the four different coloured shading nets (green, white, black, and red) with shading intensity of 75% and plant response under these shade nets compared with the open field microenvironment.

**Growing media**: Three types of potting media used for this experiment consisting of 100% soil used as a control, Mixture of soil and farmyard manure 1:1 (v/v) proportion and the third is a mixture of soil, farmyard manure and cocopeat 1:1:1 (v/v) proportion. These filled in standard 6" sized earthen pots.

**Preparation of planting material**: In *Rhoeo discolor*, divisions were collected from matured plants with one or two roots per division. Later the divisions were planted in each pot according to treatments.

**Chart 1. Characteristics of plant used in the experiment**

| Boat lily | Venation | Shape  | Arrangement | Margins | Apex      | Texture | Leaf type | Persistence |
|-----------|----------|--------|-------------|---------|-----------|---------|-----------|-------------|
|           | Parallel | Linear | Alternate   | Entire  | Acuminate | Medium  | Simple    | Evergreen   |

**List 1. Experimental layout**

|                           |   |
|---------------------------|---|
| <b>Design</b>             | CRD   |
| <b>Replications</b>       | 3   |
| <b>Treatments</b>         | 15  |
| <b>Growing conditions</b> | control (without shade net) and Colored shade net (green, white, black and red) |
| <b>Container</b>          | The dimension of pot Height: 6" Top width 15 cm and Bottom width 8 cm           |
| <b>Media</b>              | Soil, FYM and Cocopeat  |
| <b>Period of study</b>    | Five months   |

**List 2. Treatment details**

|   |  |
|---|--|
| <b>T1</b> : Control {Open + Soil}                           | <b>T9</b> : White colour shade net + Soil + FYM + Coco peat  |
| <b>T2</b> : Open + Soil + FYM                               | <b>T10</b> : Black colour shade net + Soil                   |
| <b>T3</b> : Open + Soil + FYM + Coco peat                   | <b>T11</b> : Black colour shade net + Soil + FYM             |
| <b>T4</b> : Green colour shade net + Soil                   | <b>T12</b> : Black colour shade net + Soil + FYM + Coco peat |
| <b>T5</b> : Green colour shade net + Soil + FYM             | <b>T13</b> : Red colour shade net + Soil                     |
| <b>T6</b> : Green colour shade net + Soil + FYM + Coco peat | <b>T14</b> : Red colour shade net + Soil + FYM               |
| <b>T7</b> : White colour shade net + Soil                   | <b>T15</b> : Red colour shade net + Soil + FYM + Cocopeat    |
| <b>T8</b> : White colour shade net + Soil + FYM             | * Open- control, Soil- control.                              |

### 3. RESULTS AND DISCUSSION

The results of the experiment entitled the influence of coloured shade net and media mixture on growth and development of *Rhoeo discolor*, at 30 DAP revealed that treatment T15 with red shade-net and soil + FYM + cocopeat mixture (1:1:1 v/v) significantly enhanced the plant height (11.74 cm) and that stood at par with treatments T12 (black shade net + soil + FYM+ cocopeat), T9 (white shade net + soil + FYM + cocopeat) and T14 (red shade net + soil + FYM), the minimum plant height (3.67 cm) recorded in T4 (green shade net + soil) (Table 1). The number of leaves per plant was significantly higher (8.12) in T12 (black shade net + soil + FYM + cocopeat) and the data remained at par with T3 (open + soil + FYM+ cocopeat), T5 (green shade net + soil + FYM), T6 (green shade net + soil + FYM+ cocopeat), T8 (white shade net + soil + FYM), T9 (white shade net + soil + FYM + cocopeat), T11 (black shade net + soil + FYM), T14 (red shade net+ soil + FYM)and T15 (red shade net + soil + FYM + cocopeat), minimum (4.23) in T4 (green shade net + soil).

Hence at 30 DAP, T15 (red shade net + soil + FYM + cocopeat), significantly increased the plant height (11.74 cm) and number of leaves per plant (8.12) per plant along treatments T9 (white shade net + soil + FYM+ cocopeat), T12 (black shade net + soil + FYM+ cocopeat) and T14 (red shade net + soil + FYM) were also showing better result for both the character. (Table 1)

At 90 DAP, T15 (red shade net + soil + FYM+ cocopeat), increased the plant height (19.35 cm) over other treatments being lowest (4.33 cm) in T4 (green shade net + soil). The T15 (red shade net + soil + FYM+ cocopeat) also increased the leaf number, and the data stood at par with T14 (red shade net + soil + FYM). Hence after (13.67 cm) 90 DAP the plant height as well as the number of leaves per plant significant in T15 (red shade net + soil + FYM+ cocopeat) over other treatments presented in the Table 1.

Results at 150 DAP given in Table 2 reveals that among treatments plant height recorded significantly maximum (22.11 cm) in T15 (red shade net + soil + FYM+ cocopeat) and that stood at par with T14 (red shade net + soil + FYM), T13 (red shade net + soil), T12 (black shade net + soil + FYM+ cocopeat) and T9

(white shade net + soil + FYM + cocopeat) and lowest (5.11 cm) in T4 (green shade net + soil).

Plant spread for East-West direction was significantly maximum (30.56 cm) in T5 (green shade net + soil + FYM) and it was at par with T12 (black shade net + soil + FYM + cocopeat), T14 (red shade net + soil + FYM) and T15 (red shade net + soil + FYM + cocopeat), the least (10.44 cm) in T1 (open + soil) (Table 2). Plant spread from North-South direction significantly wider (29.33 cm) in T14 (red shade net + soil + FYM) and data abide at par with T15 (red shade net + soil + FYM + cocopeat) and T13 (red shade net + soil), the spread narrow (11.44 cm) in T1 (open + soil).

There was a significant difference between growing conditions for the number of suckers per plant (Table 2). The maximum number of suckers per plant (6.22) was recorded in T3 (open + soil + FYM + cocopeat) and data stood at par with T9 (white shade net + soil + FYM + cocopeat) and T15 (red shade net + soil + FYM + cocopeat), whereas plants under T4 (green shade net + soil) and T10 (black shade net + soil) did not produce any suckers.

The number of leaves per plant was found to be significantly higher (15.89) in T15 (red shade net + soil + FYM + cocopeat) and minimum (7.22) in T10 (black shade net + soil) (Table 2).

The leaf area recorded maximum (110.42 cm<sup>2</sup>) in T15 (red shade net + soil + FYM + cocopeat) and the data stood at par with T12 (black shade net + soil + FYM + cocopeat) and T14 (red shade net + soil + FYM) (Table 2), it was minimum (38.32 cm<sup>2</sup>) in T4 (green shade net + soil).

Leaf thickness significantly augmented (1.56 mm) in T5 (green shade net + soil + FYM) and the data stood at par with T6 (green shade net + soil + FYM + cocopeat), T11 (black shade net + soil + FYM), T12 (black shade net + soil + FYM + cocopeat), T14 (red shade net + soil + FYM) and T15 (red shade net + soil + FYM + cocopeat) being lowest (0.79 mm) in T4 (green shade net + soil) (Table 2).


The SPAD reading for chlorophyll content (Table 2) found to be significantly higher (22.43) in T4 (green shade net + soil) and data stood at par with T5 (green shade net + soil + FYM) and T6 (green shade net + soil + FYM + cocopeat), the lowest (5.33) was in T12 (black shade net + soil + FYM + cocopeat).

**Table 1. Influence of coloured shade nets and potting media on Plant Height and Number of Leaves per plant of *Rhoeo discolor* at 30 and 90 days after planting**


| Treatments  | 30 DAP            |                            | 90 DAP            |                            |
|---|-------------------|----------------------------|-------------------|----------------------------|
|   | Plant Height (cm) | Number of Leaves per plant | Plant Height (cm) | Number of Leaves per plant |
| T <sub>1</sub> - Soil                                   | 5.48              | 5.33                       | 7.72              | 7.00                       |
| T <sub>2</sub> - Soil + FYM                             | 5.91              | 5.80                       | 7.95              | 6.96                       |
| T <sub>3</sub> - Soil + FYM+ Cocopeat                   | 7.26              | 7.42                       | 9.33              | 9.20                       |
| T <sub>4</sub> - Green Shade Net+ Soil                  | 3.67              | 4.23                       | 4.33              | 6.44                       |
| T <sub>5</sub> - Green Shade Net+ Soil + FYM            | 7.50              | 7.00                       | 9.53              | 9.60                       |
| T <sub>6</sub> - Green Shade Net+ Soil + FYM+ Cocopeat  | 7.40              | 8.00                       | 9.57              | 10.20                      |
| T <sub>7</sub> - White Shade Net+ Soil                  | 6.80              | 4.44                       | 9.33              | 7.22                       |
| T <sub>8</sub> - White Shade Net+ Soil + FYM            | 9.56              | 6.78                       | 13.78             | 8.46                       |
| T <sub>9</sub> - White Shade Net+ Soil + FYM+ Cocopeat  | 10.60             | 7.98                       | 14.22             | 9.20                       |
| T <sub>10</sub> - Black Shade Net+ Soil                 | 6.58              | 4.56                       | 7.78              | 5.80                       |
| T <sub>11</sub> - Black Shade Net+ Soil + FYM           | 9.83              | 6.78                       | 13.41             | 8.82                       |
| T <sub>12</sub> - Black Shade Net+ Soil + FYM+ Cocopeat | 11.32             | 8.12                       | 17.21             | 9.82                       |
| T <sub>13</sub> - Red Shade Net+ Soil                   | 7.19              | 5.22                       | 10.12             | 7.67                       |
| T <sub>14</sub> - Red Shade Net+ Soil + FYM             | 10.37             | 6.89                       | 17.22             | 12.33                      |
| T <sub>15</sub> - Red Shade Net+ Soil + FYM+ Cocopeat   | 11.74             | 7.78                       | 19.35             | 13.67                      |
| <b>S.Em ±</b>   | 0.47              | 0.51                       | 0.62              | 0.70                       |
| <b>CD @ 5%</b>  | 1.37              | 1.47                       | 1.78              | 2.03                       |

**Table 2. Influence of coloured shade nets and potting media on different growth parameters**

| Treatments  | Plant Height (cm) | Plant Spread E-W (cm) | Plant Spread N-S (cm) | Number of Suckers per plant | Number of Leaves per plant | Leaf Area (cm <sup>2</sup> ) | Leaf Thickness (mm) | Chlorophyll Content (SPAD) |
|---|-------------------|-----------------------|-----------------------|-----------------------------|----------------------------|------------------------------|---------------------|----------------------------|
| T <sub>1</sub> – Soil                                   | 8.89              | 10.44                 | 11.44                 | 1.44                        | 8.33                       | 44.26                        | 1.20                | 6.28                       |
| T <sub>2</sub> - Soil + FYM                             | 9.22              | 13.78                 | 13.78                 | 1.78                        | 8.22                       | 52.60                        | 1.16                | 9.72                       |
| T <sub>3</sub> - Soil + FYM+ Cocopeat                   | 10.78             | 16.56                 | 15.22                 | 6.22                        | 10.67                      | 63.20                        | 1.23                | 9.69                       |
| T <sub>4</sub> - Green Shade Net+ Soil                  | 5.11              | 18.11                 | 16.22                 | 0.00                        | 9.22                       | 38.32                        | 0.79                | 22.43                      |
| T <sub>5</sub> - Green Shade Net+ Soil + FYM            | 10.67             | 30.56                 | 28.22                 | 1.22                        | 11.22                      | 65.23                        | 1.56                | 21.29                      |
| T <sub>6</sub> - Green Shade Net+ Soil + FYM+ Cocopeat  | 10.89             | 25.33                 | 25.11                 | 1.22                        | 11.33                      | 73.62                        | 1.33                | 18.69                      |
| T <sub>7</sub> - White Shade Net+ Soil                  | 13.56             | 17.78                 | 19.56                 | 0.78                        | 9.67                       | 44.28                        | 0.92                | 12.70                      |
| T <sub>8</sub> - White Shade Net+ Soil + FYM            | 17.00             | 22.67                 | 21.11                 | 3.67                        | 10.11                      | 74.26                        | 0.86                | 15.26                      |
| T <sub>9</sub> - White Shade Net+ Soil + FYM+ Cocopeat  | 19.44             | 23.11                 | 22.11                 | 5.78                        | 10.78                      | 75.21                        | 1.19                | 10.71                      |
| T <sub>10</sub> - Black Shade Net+ Soil                 | 11.00             | 15.89                 | 16.78                 | 0.00                        | 7.22                       | 46.28                        | 1.07                | 7.33                       |
| T <sub>11</sub> - Black Shade Net+ Soil + FYM           | 16.89             | 24.33                 | 24.89                 | 1.78                        | 10.56                      | 78.36                        | 1.32                | 6.52                       |
| T <sub>12</sub> - Black Shade Net+ Soil + FYM+ Cocopeat | 19.67             | 26.78                 | 24.22                 | 2.33                        | 11.22                      | 96.43                        | 1.40                | 5.33                       |
| T <sub>13</sub> - Red Shade Net+ Soil                   | 20.67             | 25.67                 | 26.56                 | 0.11                        | 11.78                      | 72.68                        | 1.19                | 8.34                       |
| T <sub>14</sub> - Red Shade Net+ Soil + FYM             | 21.89             | 29.33                 | 29.33                 | 3.11                        | 12.22                      | 108.20                       | 1.30                | 12.33                      |
| T <sub>15</sub> - Red Shade Net+ Soil + FYM+ Cocopeat   | 22.11             | 28.78                 | 28.78                 | 4.89                        | 15.89                      | 110.42                       | 1.32                | 14.82                      |
| <b>S.Em ±</b>   | <b>1.05</b>       | <b>1.59</b>           | <b>1.28</b>           | <b>0.48</b>                 | <b>0.94</b>                | <b>5.66</b>                  | <b>0.09</b>         | <b>1.31</b>                |
| <b>CD @ 5%</b>  | <b>3.05</b>       | <b>4.60</b>           | <b>3.69</b>           | <b>1.37</b>                 | <b>2.72</b>                | <b>16.36</b>                 | <b>0.27</b>         | <b>3.79</b>                |



**Fig. 1. General view of different treatment effects on *Rhoeo discolor***



**Fig. 2. Best treatment T<sub>15</sub> (red shade net with soil + FYM + cocopeat media)**

Thus, at 150 DAP the T5 (green shade net + soil + FYM) increased Plant spread East-West (30.56 cm) and leaf thickness (1.56 mm) significantly, while T15 (red shade net + soil + FYM + cocopeat) performed best in all the characters and it had significantly enhanced the plant height (22.11 cm), the number of leaves per plant (15.89) and leaf area (110.42 cm<sup>2</sup>). Whereas T14 (red shade net + soil + FYM) increased the plant spread North-South (29.33 cm) and also showed good results for all the characters. The number of suckers per plant was maximum (6.22) in T3 (soil + FYM + cocopeat), and T4 (green shade net + soil) significantly increased the SPAD readings for chlorophyll content of the leaf.

The results of the present investigation revealed that Boat lily plants grown under red shade net and pots filled with soil, FYM and cocopeat media mixture in 1:1:1 (v/v) proportion significantly enhanced the various growth parameters viz., plant height (22.11 cm), number of leaves per plant (8.12), leaf area (110.42 cm<sup>2</sup>), plant spread EW (28.78 cm), plant spread N-S (28.78 cm) and leaf thickness (1.32 mm). The luxuriant vegetative growth with high plant height under red shade net confirms the study

conducted by Crowley and Kessler [7], where *Ageratum* was tallest after 4 weeks and 8 weeks under red shade net and shortest under control.

More number of leaves under red shade net similar observation was also represented by Kawabata et al. [8] in *Dracaena marginata* under red shade net, has influenced more new cane growth and produced more new leaves compared to other treatments, this may be attributed to the enhanced metabolic activities like high photosynthesis and low transpiration during the plant growth due to the quality of light received by the plants. The result corroborated with the findings of Cummings et al. [9] in peas, Leite et al. [10] in *Phalaenopsis* and Oren-Shamir et al [11] in *Pittosporium variegatum*.

Splendid growth in the media mixture comprised soil, FYM and cocopeat; wider plant spread under this media is supported by the study conducted on pot mums by Nair et al. [12] found that Cocopeat only gave wider plant spread N-S, plant spread E-W. Khelikuzzaman [13] conducted study on *Tradescantia* spp. and proved that soil mixture cocopeat, topsoil and sand (1:1:1 v/v) was significantly increasing

vegetative growth and better than other soil mixtures. More number of suckers per plant under this media was supported by the studies of Smitha [14] reported more number of suckers per plant in Anthurium with cocopeat applied alone or in combinations with farmyard manure. The reason might be due to the ability to provide proper aeration to plants in order to sustain the development and growth of roots and shoots. The media improved water relationship, free air movement, retains moisture and nutrients for the growth of the plants perhaps due to optimum bulk density, the potentiality of adequate water absorption of this growing media [15].

#### 4. CONCLUSION

The experiment demonstrated that the combination of a red shade net and a media mixture of soil, farmyard manure (FYM), and cocopeat (1:1:1 v/v) significantly enhances the growth and development of *Rhoeo discolor*. At 150 days after planting (DAP), treatment T15 yielded the highest plant height (22.11 cm), leaf number (15.89), and leaf area (110.42 cm<sup>2</sup>), indicating superior vegetative growth. Additionally, the results showed that the red shade net positively influenced various growth parameters, corroborating findings from previous studies on other plant species. The study underscores the importance of optimal growing conditions, including shade and media composition, in maximizing plant performance. Overall, the findings provide valuable insights for improving the cultivation practices of *Rhoeo discolor* and similar ornamental plants.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Wang Y, Wang J, Chen D, Hui Z, Hu X. Shade increased seed yield and quality of *incarvillea sinensis* var. *przewalskii*. *Plants*. 2023;12(16):934.
2. Kuniga T. Modification of the light environment influences the production of horticultural crops. *Japan Agricultural Research Quarterly: JARQ*. 2020;54(4):285-291.
3. Jokar A, Zare H, Zakerin A, Aboutalebi Jahromi A. Effects of shade net colors on mineral elements and postharvest shelf life and quality of fresh fig (*Ficus carica* L.) under rain-fed condition. *Horticulturae*. 2021;7(5):93.
4. Shahak Y, Elazar G, Gal E and Ganelevin R. Colornets: Crop protection and light-quality manipulation in one technology. *Acta Horticulturae*. 2004;659:143-151.
5. Rajapakse NC, Shahak Y. Light quality manipulation by horticulture industry. In: G. Whitlam and K. Halliday (eds.), *Light and Plant Development*, Blackwell Publishing. 2007;290-312.
6. Sandeep K, Fatmi U, Talang D, Priyatham K. Effect of potting media on growth and development in different species of *Nephrolepis* fern under shade net conditions (*N. falcata*, *N. cardifolia* duffii, *N. multifolia*). *Journal of Pharmacognosy and Phytochemistry*. 2019;7(5):3006-3009.
7. Crowley K, Kessler JR. Effects of colored plastic film on several field grown and greenhouse grown cut flower species, MS Thesis, Auburn University, Auburn, AL; 2007.
8. Kawabata A, Lichty J, Kobayashi K, Sahai W. Effects of photo selective shade cloths on potted *Dracaena deremensis* 'Janet Craig' and *Dracaena marginata* 'Colourama'. *Journal of Hawaiian and Pacific Agriculture*. 2007;14:49-54.
9. Cummings IG, Foo E, Weller JL, Reid JB, Koutoulis A. Blue and red photo selective shade cloths modify pea height through altered blue irradiance perceived by the cry1 photoreceptor, *Journal of Horticultural Science & Biotechnology*. 2008;83(5):663-667.
10. Leite CA, Ito RM, Lee GTS, Ganelevin R, Fagnani MA. Light spectrum management using colored nets to control the growth and blooming of *Phalaenopsis*. *Acta Horticulturae*. 2008;770:177-184.
11. Oren-Shamir M, Gussakovsky E, Shpiegel E, Nissim-Levi A, Ratner K, Ovadia R, Giller Y, Shahak Y. Coloured shade nets can improve the yield and quality of green decorative branches of *Pittosporum variegatum*. *The Journal of Horticultural*



- Science and Biotechnology. 2001;76: 353-361.
12. Nair AS, Usha-Bharathi T. Influence of potting media composition on pot mum production. International Journal of Life Sciences. 2015;10(1):73-76.
  13. Khelikuzzaman MH. Effect of different potting media on growth of a hanging ornamental plant (*Tradescantia* sp.). Journal of Tropical Agriculture and Food Science. 2007;35(1):41–48.
  14. Smitha. Effect of different growing media and pH on growth and development of *Anthurium andreanum* Lind, M.Sc. (Hort.) Thesis submitted to University of Agricultural Sciences, Bangalore. 1999;53-54.
  15. Padhiyar BM, Bhatt DS, Desai KD, Patel VH, Chavda JR. Influence of different potting media on growth and flowering of pot chrysanthemem var. Ajina purple. International Journal of Chemical Studies. 2017;5(4):1667-1669.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/121218>