



Evaluation of Novel Insecticides for Controlling Mango Leaf Hopper (*Amritodes atkinsoni*) in Mango (*Mangifera indica*) Cultivation

Sunil Kumar Mandal ^{a+++*}, Surendra Prasad ^{b++} and Manoj Kumar ^{b#}

^a RRS, Jhanjharpur, Madhubani, Dr. Rajendra Prasad Central Agricultural University, Pusa (Samastipur)- 848125, Bihar, India.

^b Department of Entomology, Dr. Rajendra Prasad Central Agricultural University, Pusa (Samastipur)- 848125, Bihar, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i82797>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121333>

Original Research Article

Received: 13/06/2024

Accepted: 17/08/2024

Published: 23/08/2024

ABSTRACT

The study was carried out to the evaluation of newer insecticides molecules against mango leaf hopper, *Amritodes atkinsoni* Leth. infesting in Maldah. The treatments Thiomethoxam 25 WG (0.3 gm/L) and Imidacloprid 17.8 SL (0.3 ml/L) were found highest per cent reduction (87.74% and 86.08%) against mango leaf hopper. Moreover, Clothianidin 50 WDG (0.25 gm/L), Difentheuron 50 WP and Acetamiprid 20 SP (0.5 gm/L) were next effective treatments against mango leaf hopper

⁺⁺ Assistant Professor-cum Scientist;

[#] Associate Professor-cum Senior Scientist;

*Corresponding author: E-mail: skmandal6464@gmail.com;

Cite as: Mandal, Sunil Kumar, Surendra Prasad, and Manoj Kumar. 2024. "Evaluation of Novel Insecticides for Controlling Mango Leaf Hopper (*Amritodes Atkinsoni*) in Mango (*Mangifera Indica*) Cultivation". *Journal of Experimental Agriculture International* 46 (8):1103-9. <https://doi.org/10.9734/jeai/2024/v46i82797>.

i.e. 81.22%, 79.98% and 77.73%, respectively. The other treatments viz, Cypermethrin 10 EC (1.5 ml/L), Buprofezin 25 EC (1.0 ml/L) and Azadirachtin 3000 PPM (3 ml/L) were found least effective against mango leaf hopper over control. The significantly highest fruit yield and benefit cost ratio was also found in treatment Thiomethoxam 25 WG (232.67 kg/tree and 4.33:1) in comparison to farmer's practice (192.43 kg/tree and 2.16:1).

Keywords: Evaluation; insecticide; mango leaf hopper and mango.

1. INTRODUCTION

Mango, *Mangifera indica* is considered as one of the most important fruit crop of tropical and subtropical region of India and is known as "king of fruits" for wide adaptability, attractive colour, delicious taste, exotic flavour, high nutritive value, richness in variety, attractive appearance and popularity among the people. In India the production of mango fruits is 20946.3 thousand metric tonnes with area of 2370.8 thousand hectare in 2021-22 (India stat, 2021). Both fresh fruits and processed products made from mangoes are in high demand. As a result, there is now a need to boost both mango fruit productivity and quality. The insects are one of the limiting factors for higher production of good quality fruits. Among different insect-pests, mango leaf hoppers are recorded as major, most serious and widespread sucking pests throughout the year in mango ecosystem. Three species of mango leaf hoppers, *Amritodes atkinsoni*, *Idioscopus clypealis* and *Idioscopus nitidulus* remain active throughout the year and damage each crop stage from emergence of new flush to fruiting stages [1-4] and cause upto 100 per cent yield losses. The leaf hoppers cause a loss of 20-100 per cent of inflorescences Both nymphs and adults of leaf hoppers suck the sap from the young leaves, tender shoots, inflorescences, panicles; branches and rachis of the young fruits which causes non-setting of flowers and dropping of the immature fruits. Leaf hoppers also excrete huge quantities of honey dew resulting in growth of sooty mould formation, thus affecting the photosynthetic activity of the plant, ultimately in poor fruit set and leads to reduction in yield. Dalvi et al. [5] considered it as major pest of mango, and is directly responsible in reducing the yield qualitatively and quantitatively. Considering the importance and the damage potential of the pest. The present study was carried out to know the efficacy of some newer insecticide molecules against the mango leaf hoppers.

2. MATERIALS AND METHODS

The investigation was conducted in the mango orchard (cv. Maldah) of the five different farmer's villages in Siwan district of Bihar during 2020 and 2021. The experiment was carried out in Randomized Block Design with ten treatments including control. Uniformly flowering 10 to 12 years old trees was selected for imposing the treatments and the observations were taken on them considering one tree as one replication. Ten panicles were randomly selected/tree from all directions of lower part of the tree canopy during panicle initiation stage and tagged for recording observations. In each treatment except control, the need based applications of insecticides were given when the hopper population reached to five or excluding five (nymphs and adults) on each randomly selected panicles of experimental tree. Spray fluid was prepared by mixing measured quantity of water and insecticides. Twenty liters of spray solution were used per tree. The respective insecticides were applied as a foliar spray on the tree with the help of tractor mounted power sprayer. An untreated check was also maintained for comparison. Population of mango leaf hoppers (nymphs and adults) were recorded visually on ten tagged panicles/tree. Leaf hopper population were counted one day before spraying and 7th, 10th and 15th days after insecticidal application of each treatment and with the 1st and 2nd spray, respectively.

3. RESULTS AND DISCUSSION

Different new insecticide molecules were tried to check their comparative efficacy against mango hopper during 2020 and 2021, and the results obtained are presented (Table 1 Pooled). Analysis of data on mango hopper population recorded before treatments indicated non-significant results suggested that the hopper population was homogeneous. The hopper population was recorded after 7 days of spraying showed that the superiority of treatment Thiomethoxam by registering least number of hoppers/panicle (1.15) and it was at par

with Imidacloprid, Acetamiprid, Difentheuron and Clothianidin noted hopper population of 1.23, 1.58, 1.72 and 1.78/panicle, respectively. Cypermethrin exhibited 2.87 hoppers/panicle and it was statistically equally effective as letter Buprofezin (2.94/panicle). On the other hand, the highest number of hopper population per panicle was recorded in the treatment of Azadirachtin (3.38) but it was at par with Dimethoate (farmer's practice). After 10 days of insecticidal application, Thiomethoxam, Difentheuron, Clothianidin and Imidacloprid were statistically equally effective against mango hoppers and exhibited hopper population of 2.24 to 3.13/panicle, respectively. Acetamiprid, Cypermethrin, Buprofezin and Azadirachtin were also statistically equally effective (Table 1). At 10 DAS the maximum number of hopper population per panicle was recorded in Dimethoate (6.86) but significantly superior over the untreated check (10.48). Among all the tested insecticides, Thiomethoxam found statistically superior against mango hoppers after 15 days of insecticidal application by recording 3.06 hoppers/panicle. All the insecticidal treatments were significantly at par with them. Azadirachtin and Dimethoate treated trees exhibited 8.47 and 9.14 hoppers/panicles, respectively. The mean result of first spray showed that the treatment with Thiomethoxam registered lower incidence of hopper/panicle (2.15) and it was at par with Imidacloprid, Clothianidin, Difentheuron and Acetamiprid exhibited 2.79, 3.34, 3.50 and 3.98 hoppers/panicle and they were at par with each other (Table 1).

The efficacy of insecticidal treatments on second spray against hopper population pooled data was given in Table 1. All the insecticidal treatments showed an increasing trend of hopper population at 15 DAS. The data recorded on seventh days after the second spray revealed that Thiomethoxam treated tree recorded minimum hopper population (1.17), which was significantly at par with Imidacloprid (1.47) and followed by Clothianidin (1.73), Cypermethrin (1.84), Difentheuron (1.93), respectively. Buprofezin and Azadirachtin were found next effective treatment against mango hopper. The maximum hopper population was recorded in treatment of Dimethoate as farmer's practice (3.53). In contrast to above 15.37 hoppers/panicle recorded in untreated trees. After ten days of insecticidal application of treatments, the results showed that the treatment comprises of Thiomethoxam was found statistically superior in controlling the pest and the recorded 1.23 hoppers/panicle. Next

effective treatments the Clothianidin, Difentheuron, Imidacloprid and Acetamiprid and Cypermethrin, which were found statistically at par with each other and showing average surviving population of 1.82, 2.03, 2.10, 2.16 and 2.38 panicle, respectively. On 15th DAS, it was found that Thiamethoxam was the superior treatment (1:26 hoppers/panicle). Similarly Kapadia, et al. [6] and Patel, et al. [7] reported similar result. Next promising treatments were Clothianidin, Difentheuron, Imidacloprid and Acetamiprid recorded 1.86, 2.10, 2.19 and 2.24 hopper/panicle, respectively. Whereas, Buprofezin, Cypermethrin, Dimethoate recorded 2.33, 2.89 and 3.74 hopper/panicle, respectively. The maximum hoppers/panicle (3.83) recorded in the treatment of Azadirachtin, but significantly superior with respect to untreated trees (19.23 hoppers/panicle). The mean analysis of second spray, presented in Table 1, showed higher effectiveness of Thiomethoxam by resisting lower hopper population (1.22) and it was at par with Imidacloprid, Clothianidin and Difentheuron, which recorded 1.69, 1.84 and 2.02 hoppers/panicle, respectively. These four treatments were significantly more effective than rest of the treatments. Acetamiprid, Cypermethrin and Buprofezin registered 2.16, 2.37 and 2.58 hoppers per panicle, respectively and significantly more effective than Azadirachtin and Dimethoate (Farmer's practice). Similar results were observed by Singh et al. [8].

The overall mean values computed for two sprays, clearly indicated that the treatment of Thiomethoxam (1.69 hoppers/panicle and 87.74% reduction over control) and Imidacloprid (1.92 hoppers/panicle and 86.08% reduction over control) were found statistically superior against mango hopper in comparison to rest of the insecticides tested. Difentheuron, Clothianidin and Acetamiprid were next best effective treatments and exhibited 2.76, 2.59 and 3.07 hoppers/panicle and 79.98%, 81.22% and 77.73% reduction over control, respectively. Cypermethrin noted 3.56 hoppers/panicle and 77.73% reduction over control and it was at par with Buprofezin (3.73 hoppers/panicle and 72.95% reduction over control). The highest population of hoppers (4.39) was noticed in trees treated with Azadirachtin (68.16% reduction over control) and it was statically equally effective with Dimethoate as farmer's practice (5.06 hoppers/panicle and 63.31% reduction over control). These two treatments were found poor in their effectiveness against mango hopper. In past, among evaluated different insecticides

Table 1. Efficacy of newer insecticides against mango leaf hoppers during 2020 and 2021 (pooled)

Treatment	Dose g/ml/L	Mean number of mango hoppers/panicle										% reduction over control	
		First Spray					Second Spray						
		1DBS	7 DAS	10 DAS	15 DAS	Mean	1 DBS	7 DAS	10 DAS	15 DAS	Mean		Over all mean
Clothianidin 50 WDG	0.25 ml	5.28 (2.40)	1.78 (1.51)	3.13 (1.90)	5.11 (2.37)	3.34 (1.96)	6.36 (2.62)	1.73 (1.49)	1.82 (1.52)	1.86 (1.54)	1.84 (1.53)	2.59 (1.76)	81.22
Imidacloprid 17.8 SL	0.30 ml	5.67 (2.48)	1.23 (1.32)	3.02 (1.88)	4.12 (2.15)	2.79 (1.81)	6.03 (2.56)	1.47 (1.40)	2.10 (1.61)	2.19 (1.64)	1.69 (1.48)	1.92 (1.56)	86.08
Difentheuron 50 WP	0.50gm	6.35 (2.62)	1.72 (1.49)	2.91 (1.85)	5.87 (2.52)	3.50 (2.00)	6.63 (2.67)	1.93 (1.56)	2.03 (1.59)	2.10 (1.61)	2.02 (1.59)	2.76 (1.81)	79.98
Thiomethoxam 25 WDG	0.30gm	5.28 (2.40)	1.15 (1.28)	2.24 (1.67)	3.06 (1.89)	2.15 (1.63)	5.57 (2.46)	1.17 (1.29)	1.23 (1.32)	1.26 (1.33)	1.22 (1.31)	1.69 (1.48)	87.74
Acetamiprid 20 SP	0.50gm	4.96 (2.34)	1.58 (1.44)	3.94 (2.11)	6.41 (2.63)	3.98 (2.12)	6.01 (2.55)	2.08 (1.61)	2.16 (1.63)	2.24 (1.66)	2.16 (1.63)	3.07 (1.89)	77.73
Cypermethrin 10 EC	1.50 ml	6.14 (2.58)	2.87 (1.84)	4.18 (2.16)	7.24 (2.78)	4.76 (2.29)	7.23 (2.78)	1.84 (1.53)	2.38 (1.70)	2.89 (1.84)	2.37 (1.69)	3.56 (2.01)	74.18
Buprofezin 25 EC	1.00 ml	5.82 (2.51)	2.94 (1.85)	4.35 (2.20)	7.36 (2.80)	4.88 (2.32)	7.36 (2.80)	2.18 (1.64)	3.24 (1.93)	2.33 (1.68)	2.58 (1.75)	3.73 (2.06)	72.95
Azadirachtin 3000 PPM	3.00 ml	5.76 (2.50)	3.38 (1.97)	3.64 (2.03)	8.47 (2.99)	5.16 (2.38)	8.42 (2.99)	3.46 (1.99)	3.60 (2.02)	3.83 (2.08)	3.63 (2.03)	4.39 (2.21)	68.16
Dimethoate 30 EC (Former's Practice)	2.00 ml	6.45 (2.64)	3.24 (1.93)	6.86 (2.71)	9.14 (3.10)	6.41 (2.63)	9.13 (3.10)	3.53 (2.01)	3.87 (2.09)	3.74 (2.06)	3.71 (2.05)	5.06 (2.36)	63.31
Untreated Check (Control)	-	6.59 (2.66)	8.26 (2.96)	10.48 (3.31)	11.65 (3.49)	10.13 (3.26)	12.63 (3.62)	15.37 (3.98)	17.73 (4.27)	19.23 (4.44)	17.44 (4.23)	13.79 (3.78)	0
SEm (±)	-	0.132	0.114	0.137	0.105	0.052	0.106	0.082	0.072	0.081	0.073	0.060	-
CD (P= 0.05)	-	NS	0.342	0.412	0.318	0.153	0.318	0.241	0.216	0.236	0.211	0.182	-

Figures in parentheses are $\sqrt{X + 0.5}$ transformation values.
 DBS = Days before spraying, DAS = Days after spraying

Table 2. Economics of newer insecticides against mango leaf hopper during 2020 and 2021 (pooled)

Treatments	Dose g/ml/L	Yield (kg/tree)	Increased Yield over control	Value of additional yield over control (Rs./tree)	Cost of treatment (Rs/tree)	Incremental benefit (Rs/tree)	BC ratio
Clothianidin 50 WDG	0.25 ml	225.53	100.84	4043.60	935.24	3098.36	3.31 : 1
Imidacloprid 17.8 SL	0.30 ml	228.24	103.55	4142.00	842.57	3299.43	3.92 : 1
Difentheuron 50 WP	0.50gm	224.31	99.62	3984.80	814.38	3170.42	3.89 : 1
Thiomethoxam 25 WDG	0.30gm	232.67	107.98	4319.20	810.65	3508.55	4.33 : 1
Acetamiprid 20 SP	0.50gm	218.45	93.76	3750.40	872.40	2878.00	3.30 : 1
Cypermethrin 10 EC	1.50 ml	210.18	85.49	3419.60	867.84	2551.76	2.94 : 1
Buprofezin 25 EC	1.00 ml	203.26	78.57	3142.80	875.52	2267.28	2.58 : 1
Azadirachtin 3000 PPM	3.00 ml	198.31	73.62	2944.80	843.37	2101.43	2.49 : 1
Dimethoate 30 EC (Former's Practice)	2.00 ml	192.43	67.74	2709.60	854.85	1854.75	2.16 : 1
Untreated Check (Control)	-	124.69	-	-	-	-	-
SEm (\pm)	-	5.653	-	-	-	-	-
CD (P= 0.05)	-	16.962	-	-	-	-	-

Market price of mango fruits @ Rs. 40.00 / kg.

were evaluated against hopper complex on mango, Thiomethoxam (0.0084%) was found most effective treatment noted by Kumar et al. [1] and Patel et al. [7]. Tumbada et al. [9] also observed Thiomethoxam 25 WG (0.0075%) and Acetamiprid 20 SP (0.005%) as the most effective treatments in checking the population of mango hopper. Munj and Rana [10] reported that Thiomethoxam 25 WG and Acetamiprid 20 SP were equally effective. Azadirachtin as least effective against mango hopper and Acetamiprid 20 SP were at par with each other whereas, Buprofezin 25 SC was found in next to order of toxicity against mango hoppers. Chaudhari et.al. [11] Concluded that Thiomethoxam 0.0025% showed superior effectiveness against mango hopper. Shawan et al. [12] found Acetamiprid 20 SP as on effective insecticide by controlling 93.33 per cent hopper population when these were applied at 0.0025% concentration. Karar et al., [13] found Thiomethoxam @ 10 g/100 litre of water as the most effective for management of mango hopper. Kadavkar et al., (2021) and Patel [7] in a study reported that Thiomethoxam 25 WG effective insecticide followed by Buprofezin 25% SC and Azadirachtin against mango hopper. All these earlier findings are more or less in contrast with the present finding and fully support the present result.

The yield data (Table 2) revealed significantly higher marketable mango fruit yield of 232.67 kg/tree in Thiomethoxam treated tree followed by Imidacloprid (228.24 kg/tree), Clothianidin (225.53 kg/tree) and Difentheuron (224.31 kg/tree), all being at par with each other. Dimethoate (Farmer's practice) treated tree produced lower marketable fruit yield (192.43 kg/tree), but significantly more than untreated check (124.69 kg/tree) The Cost-benefit analysis of different sate of treatments revealed that the maximum monitory benefit of Rs. 4319.20/tree accrued from Thiomethoxam treated tree. The most effective treatment in redacting the hopper incidence as well as fruit yield realised as per tree basis. Yet the highest benefit: cost ratio (4.33:1) was obtained in Thiomethoxam treated tree. This was followed by Imidacloprid (3.92:1), Difentheuron (3.89:1) Clothianidin (3.31:1) Acetamiprid (3.30:1) and rather moderately efficiently by Cypermethrin (2.94:1), Buprofezin (2.58:1) and Azadirachtin (2.49:1). In comparison the benefit cost ratio of 2.16:1 were lowest recorded in farmer practices (Dimethoate). Similar results were recorded by Dalvi et al. [5] and Prasad et al [14] reported B: C ratio in Imidacloprid (4.38:1) [15,16].

4. CONCLUSION

The conclude that thiomethoxam 25 WG (0.3 gm/L) and Imidacloprid 17.8 SL (0.3 ml/L) were found most effective against mango leaf hopper management and Clothianidin 50 WDG (0.25 gm/L), Difentheuron 50 WP and Acetamiprid 20 SP (0.5 gm/L) were gave also encouraging response in the management of the pest.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (Chat GPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

Authors are thankful to Senior Scientist and Head, Krishi Vigyan Kendra, Siwan, Dr. Rajendra Prasad Central Agricultural University, Pusa and ATATRI, Patna for financial assistance during my research tenure.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar SB, Raghvani R, Bhatt RI. Bio-efficacy of newer insecticide against hopper complex on alphonso mango in humid tropics of south Gujarat. J. Appl. Zool. Res. 2005;16(1):64-66.
2. Gundappa PD, Jayanthi K, Veirghese A. Migratory behaviour of mango hopper, *Idioscopus* spp. in relation to host plant flowering phenology; a synchronous shift. Bioscan. 2014;9(2):639-641.
3. Turkhade PD, Godase SK, Narangalkar AI, Dhekale JS, Haldankar PM. *In-vitro* efficacy of Entomo-pathogenic fungi against mango hopper," *Idioscopus niveosparus* (L). Ann. Pl. Protec. Sci. 2015;23:390-417.
4. Bana JK, Singh P, Makwana A. Influence of abiotic factors and crop stages on population dynamics of hoppers, *Idioscopus* spp. in mango ecosystem. Ann. pl. protec. Sci. 2016;24(2): 286-289.
5. Dalvi MB, Pushpa D, Patil, Raut SP. Pre-harvest fungicidal application for control of *Colletotrichum gloeosporioides* in alphonso

- mango fruit. Ann. pl. protec. Sci. 2010; 18(1):361-362.
6. Kapadia MN, Bhutani PG, Jeuthava DM, Varani VR, Beria NN. Integrated management of mealy bugs in custard appl. Ann. Pl. Protec. Sci. 2009;17(2):372-375.
 7. Patel T, Shukla A, Patel SR. Bio-efficacy of chemical and bio-rational insecticides against hopper, *Idioscop usuitidulus* infesting mango. Ann. Pl. Protec. Sci. 2021;29(2):88-92.
 8. Singh, Rajmdra Prasad CS. Tiwari GN. Efficacy of botanicals, bio pesticides and insecticides on mango mealy bug. Ann. Pl. Protec. Sci. 2011;18(2):311–314.
 9. Tumbada RD, Pokhrkar DS, Datkhile RV. Efficacy of entomopathogenic fungi against mungo hoppers. J. Pharmacogn and Phytochem. 2018;7(2):3198-3202.
 10. Munj AY, Rana BS. Population dynamics and management of mango hopper *Idioscopus niveasparsus* Lith. Under Konkarn region of Maharashtra. Thesis Ph.D. (Agriculture). Maharana Pratap University of Agriculture and Technology, Udaipur (Raj.); 2016.
 11. Chaudhari AV, Sridharan S, Sundar Singh SD. Management of mango hopper with newer molecules and bio-pesticides under ultra-high density planting system. Entomol. and Zool Stud. 2017;5(6):454-458.
 12. Shawan SI, Rashed BU, Mitra AS, Jahan. Efficacy of different chemical and botanical insecticides in Controlling mango hopper (*Amritodus atkinsoni* L.) Adv. Plants Agric. Res. 2018; 8(2):127-131.
 13. Karar H, Khan AH, Kiran S, Iqbal A, Vlah H. Efficacy of various insecticidal modules against mango hopper, *Idioscopus clypealis* Lethierry (*Hemiptera: Cicadellidae*) Samar Bahisht Chaunsa and their impact on yield. Pure Appl. Biol. 2020;9(3):1791-1759.
 14. Prasad Surendra, Mandal, S.K. and Kumar Manoj Management of mango leaf hopper, *Amritodes atkinsoni* Leth on mango in Saran district (Bihar). The Pharma Innovation Journal. 2023;SP-12(8):934-936.
 15. Haseeb M. Occurrence of fruit sucking bug on mango. Ann. pl. protec. Sci. 2006; 14:218-219.
 16. Samanta A, Gosh A, Hembran TK, Patra S, Somchowdhury AK. Efficacy of insecticides against mango hoppers and fruit yield. Ann. Pl. Protec Sci. 2009; 17(1):225-274.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121333>