



Screening of Mungbean Germplasms against Root-Knot Nematode, *Meloidogyne javanica* to Find Out the Source of Resistance

Seema Yadav ^{a*}, S.P. Bishnoi ^b, Hemraj Gurjar ^b, Parwati ^a,
B.S. Chandrawat ^a, Rajkumar Verma ^c and Ramavath Abhi ^c

^a Department of Nematology, SKN College of Agriculture, SKNAU, Jobner, Jaipur-303329, India.

^b Division of Nematology, Rajasthan Agricultural Research Institute, Durgapura, Jaipur, SKNAU, Jobner- 302018, India.

^c Department of Nematology, Rajasthan college of Agriculture, MPUAT, Udaipur, Rajasthan- 313001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101469>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124178>

Original Research Article

Received: 22/07/2024

Accepted: 25/09/2024

Published: 28/09/2024

ABSTRACT

Root-knot nematodes (*Meloidogyne* spp.) have been widely recorded in mung bean growing areas. Screening for resistant genotypes has reduced the damage caused by the root-knot nematodes. An experiment was conducted to find out the Screening of mungbean germplasms against root-knot nematode, *Meloidogyne javanica* to find out the source of resistance. Total forty-one cultivars

*Corresponding author: E-mail: ssy.seema@gmail.com;

Cite as: Yadav, Seema, S.P. Bishnoi, Hemraj Gurjar, Parwati, B.S. Chandrawat, Rajkumar Verma, and Ramavath Abhi. 2024. "Screening of Mungbean Germplasms Against Root-Knot Nematode, *Meloidogyne Javanica* to Find Out the Source of Resistance". *Journal of Advances in Biology & Biotechnology* 27 (10):462-68. <https://doi.org/10.9734/jabb/2024/v27i101469>.

screened in which three cultivars i.e., IPM 1603-3, MH 1142 (Ch.) and MH 1772 were found to be moderately resistant, twenty-eight cultivars recorded as susceptible and ten recorded highly susceptible ones.

Keywords: Cultivars; gall index; *Meloidogyne javanica*; mung bean; screening.

1. INTRODUCTION

Root-knot nematode, *Meloidogyne javanica* is a major threat to mung bean (*Vigna radiata* L.) cultivation. Root-knot nematode is the major pathogen causes reduction in plant growth parameters and loss of grain yield. Pulses production is most vulnerable to the attacks of pests and diseases causing yield losses [1]. Root-knot nematodes (*Meloidogyne* spp.) are one of the most destructive pests and causes severe economic losses [2,3]. Root-knot nematodes belonging to genus *Meloidogyne* are serious menace to crop production around the globe [4]. Mungbean root-knot nematode (*Meloidogyne javanica*) has emerged as a major threat throughout the world and it has occupied a place of 'National Pest' owing to its severity [5]. Buildup of inoculum of the nematode and repeated cultivation of same cultivars in the same land every year is the prime reason for yield losses by root-knot nematodes [6]. Hostplant resistance is effective management tool that increases yield in spite of nematode population densities that exceed the damage threshold [7]. There are several biotic and abiotic stresses including root-knot nematodes which are responsible for low productivity of mungbean [8]. Root-knot nematodes have a major impact on mungbean productivity, causing losses of 18-90% [8,9]. Root-knot nematodes parasitize plant root systems, directly absorption of water and essential nutrients. These are serious parasites which attack wide varieties of crop plants including pulses and are responsible for substantial economic losses [10]. During the infective stage, root-knot nematodes feed on the epidermal cells of root and penetrate through the newly formed tissues present above the meristematic zone [11]. At the initial stage of the infestation by root-knot nematode in the plant

tissue there is cell enlargement with rapid cell division in the pith and vascular bundle followed by transformation of cortex into gall [12]. Consequently, with the increment of infestation, growth and development of infested plants become stunted along with yellowing of leaf [13]. More than 80 species of *Meloidogyne* have been reported in different parts of the world. Out of them, two species *M. incognita* and *M. javanica* have been reported in pulse-based cropping systems [9,14].

2. MATERIALS AND METHODS

The experiment was conducted in root-knot nematode infected sick field at Division of Nematology, Rajasthan Agricultural Research Institute, Durgapura, Jaipur. In sick field maintained the minimum inoculum level @ 2 larvae/g soil (2000 J₂/kg soil). Total forty-one germplasm of mung bean were sown in 2.5 m rows and each line was flanked with susceptible check line. Plants were uprooted 60 days after sowing and observations recorded on number of galls per plant. Mung bean cultivar categorized as highly resistant, resistant, moderately resistant, susceptible, and highly susceptible based on root-knot index on a 1 to 5 scale (with 1 = no galls or no egg mass and 5 = greater than 101 galls or egg mass per plant) Table 1. Reaction of entries was recorded on the basis of root-knot index (RKI) (1-5 scale) according to Hartman and Sasser [15].

3. RESULTS AND DISCUSSION

In the present investigation, the result revealed that all the cultivar of mung bean showed varying degree of reaction against root-knot nematode, *M. javanica*. (Table-2 and Fig. 1). A total forty-one (41) mung bean germplasm were screened

Table 1. Root-knot index for resistant varieties

S. No.	No. of Galls/Plant	Reaction of Variety
1	No Galls	Highly Resistant (HR)
2	1-10 Galls	Resistant (R)
3	11-30 Galls	Moderate Resistant (MR)
4	31-100 Galls	Susceptible (S)
5	101 and Above Galls	Highly Susceptible (HS)

(Table 2). The mung bean cultivars were categorized highly resistant, resistant, moderately resistant, susceptible and highly susceptible on the basis of root-knot index (1-5 scale) given by Hartman and Sasser, [15]. The data presented in Tables 2 and 3 showed that among the screened forty-one germplasm, only three varieties were found as moderately resistance *i.e.*, IPM 1603-3, MH 1142 (Ch.) and MH 1772, and having 11-30 galls per plant. However, twenty-eight germplasm of mung bean *i.e.*, AKM 12-28, AKM 8802 (Ch.), BM 2019-10, BM 4 (Ch.), GJM 1701, IPM 02-14 (Ch.), IPM 02-3 (Ch.), IPM 13-6, IPM 410-3 (Ch), IPM 512-1 (Ch), IPMD 1603-7, JLPM 702-1, LGG 610, MH 1142 (Ch), MH 2-15 (Ch.), MHBC-20-2, MI 750-1, MI 98-64, ML 2506, ML 818 (Ch), OBG 104, OUM 11-5 (Ch), Pusa 0672 (Ch.), Pusa M 2171, SKNM 1904, SML 1839, VBN 4 (Ch), VCG 180-21 showed the susceptible reaction against root-knot nematode having (31-100) galls per plant. Whereas, ten cultivar of mung bean *i.e.*, COGG 912 (Ch.), PANT M4 (Ch.), PANT M6 (Ch.), PKV AKM 4 (Ch.), PM 1711, PM 1727, PUSA 1731 (Ch.), RMG 1166, SML 2015, Varsha (Ch.) were found the highly susceptible having (102-128) galls per plant against root knot nematode in mung bean. No cultivar was found highly

resistant and resistant against root-knot nematode, *Meloidogyne javanica* in mung bean.

The data are similar in accordance with Bora et al. [16] who screened 282 varieties of green gram against *Meloidogyne incognita* and observed that all the screened varieties showed susceptible reaction or highly susceptible to *Meloidogyne incognita*. Hassan and Devi [17] tested 72 chickpea, 34 pea germplasm for their resistance against root-knot nematode (*Meloidogyne incognita*) in culture plots. Amongst 72 chickpea varieties evaluated, 34 varieties showed highly resistant, 25 resistant, 9 moderately resistant and only three genotypes *viz.* C-41-42, C-42-0, C-44-2 were found susceptible. Out of 34 varieties of pea, 29 genotypes were found highly resistant, 4 resistant, 1 genotype FP-21-02 and found moderately resistant. Devi et al. [18] screened twenty-eight germplasms of mung bean against root-knot nematode (*Meloidogyne incognita*). Out of twenty-eight genotypes, 24 were susceptible and 4 were highly susceptible to *Meloidogyne incognita*. No resistant or moderately resistant genotypes recorded. Similarly, Panday and Nayak [19] screened thirty - eight varieties of

Table 2. Reaction of mung bean germplasm against root-knot nematode

RKI Scale	Cultivar (41)	Number of germplasm	Galls / Egg masses / Plant (Range)	Reaction
1	NIL	0	0	Highly Resistant (HR)
2	NIL	0	0	Resistant (R)
3	IPM 1603-3, MH 1142 (Ch.), MH 1772	3	11-30	Moderately Resistant (MR)
4	AKM 12-28, AKM 8802 (Ch.), BM 2019-10, BM 4 (Ch.), GJM 1701 IPM 02-14 (Ch.), IPM 02-3 (Ch.), IPM 13-6, IPM 410-3 (Ch), IPM 512-1 (Ch), IPMD 1603-7, JLPM 702-1, LGG 610, MH 1857, MH 2-15 (Ch.), MHBC-20-2, MI 750-1, MI 98-64, ML 2506, ML 818 (Ch), OBG 104, OUM 11-5 (Ch), Pusa 0672 (Ch.), Pusa M 2171, SKNM 1904, SML 1839, VBN 4 (Ch), VCG 18-021	28	31-100	Susceptible (S)
5	COGG 912 (Ch.), PANT M4 (Ch.), PANT M6 (Ch.), PKV AKM 4 (Ch.), PM 1711, PM 1723, PUSA 1731 (Ch.), RMG 1166, SML 2015, Varsha (Ch.)	10	101-128	Highly Susceptible (HS)




Fig. 1. Reaction of mung bean germplasm against root-knot nematode

Table 3. Reaction of mung bean germplasm against root-knot nematode

S. No.	Cultivar	Number of galls per plant	Root-knot index	Reaction
1.	AKM 12-28	57.2	4	S
2.	AKM 8802 (Ch.)	44.0	4	S
3.	BM 2019-10,	38.4	4	S
4.	BM 4 (Ch.)	54.6	4	S
5.	COGG 912 (Ch)	113.7	5	HS
6.	GJM 1701	43.8	4	S
7.	IPM 02-14 (Ch.)	34.7	4	S
8.	IPM 02-3 (Ch.)	48.3	4	S
9.	IPM 13-6	51.4	4	S
10.	IPM 1603-3	24.0	3	MR
11.	IPM 410-3 (Ch)	41.8	4	S
12.	IPM 512-1 (Ch)	47.4	4	S
13.	IPMD 1603-7	56.6	4	S
14.	JLPM 702-1	61.2	4	S
15.	LGG 610	55.0	4	S
16.	MH 1142 (Ch)	14.6	3	MR
17.	MH 1772	21.8	3	MR
18.	MH 1857	56.1	4	S
19.	MH 2-15 (Ch.)	50.5	4	S
20.	MHBC-20-2	46.9	4	S
21.	MI 750-1	47.3	4	S
22.	MI 98-64	55.6	4	S
23.	ML 2506	59.8	4	S
24.	ML 818 (Ch)	62.2	4	S
25.	OBGG 104	60.7	4	S
26.	OUM 11-5 (Ch)	63.2	4	S
27.	Pant M 4 (Ch)	107.3	5	HS
28.	Pant M 6 (Ch.)	115.4	5	HS
29.	PKV AKM 4 (Ch.)	109.7	5	HS
30.	PM 1711	118.3	5	HS
31.	PM 1723	123.4	5	HS
32.	Pusa 0672 (Ch.)	40.3	4	S
33.	Pusa 1371 (Ch)	121.2	5	HS
34.	Pusa M 2171	58.3	4	S
35.	RMG 1166	108.4	5	HS
36.	SKNM 1904	48.4	4	S
37.	SML 1839	57.3	5	HS
38.	SML 2015	105.7	5	HS
39.	Varsha (Ch)	113.8	5	HS
40.	VBN 4 (Ch)	60.2	4	S
41.	VCG 18-021	59.3	4	S

green gram against *Meloidogyne incognita*, in which thirty-four varieties found resistant reaction with 3-6 number of galls per plant, while four varieties showed moderately resistant reaction with 10-11 number of galls per plant. However, Kumar et al. [20] screened 14 mung bean genotypes against *M. javanica*, none of the

germplasm was found resistant, however, four genotypes showed, moderately resistant reaction. Devindrappa et al. [21] evaluated 24 pigeon pea genotypes against *M. javanica* and observed 5 genotypes viz., DPPA 85-12, DPPA 85-8, DPPA 85-1, DPPA 85-13 and IPA 15-1 as moderately resistant reaction, 13 showed

moderately susceptible, 3 susceptible and 2 genotypes were found in the highly susceptible reactions. The investigation clearly showed that different varieties showed different reactions to *Meloidogyne javanica*. This occurs due to genetic variation, initial inoculum level and other climatic or ecological conditions during the action and potential against the nematode.

4. CONCLUSION

Root-knot nematode, *Meloidogyne javanica* is the major pathogen causes reduction in plant growth parameters and loss of grain yield. Screening for resistant genotypes has reduced the damage caused by the root-knot nematodes. Out of 41 varieties, three were recorded as moderately resistance, ten as highly susceptible and twenty-eight varieties recorded as susceptible.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

I hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENT

The authors are grateful to Head of Department for providing facilities and support for carrying out the research experiment in Division of Nematology at RARI Durgapura and Department of Nematology at SKNAU Jobner Jaipur.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Root knot nematodes in India- A Comprehensive Monograph. AICRP on Plant Parasitic Nematodes with Integrated Approach for their Control, IARI, New Delhi, India. 2014;3.
2. Kalele DN, Affokpon A, Coosemans J, Kimenju JW. Suppression of root-knot nematodes in tomato and cucumber using biological control agents. African Journal of Horticultural Sciences. 2010;3: 72–80.
3. Collange B, Navarrete M, Peyer G, Mateille T, Tchamitchian M. Root-knot nematode (*Meloidogyne*) management in vegetable crop production: The challenge of an agronomic system analysis. Crop Protection. 2011;30:1251–1262.
4. Ali MA, Abbas A, Azeem F, Javed N, Bohlmann H. Plant-nematode interactions: from genomics to metabolomics. – International Journal of Agriculture and Biology. 2015;17(6):1071-1082.
5. Joshi V, Kumar S, Rawat S. Study on infection and development of root-knot nematode, *Meloidogyne javanica* on mungbean. J Entomol. 2020;8:1621.
6. Hussain MA, Mukhtar T, Kayani MZ. Reproduction of *Meloidogyne incognita* on resistant and susceptible okra cultivars. Pakistan Journal of Agricultural Sciences. 2016;53(2).
7. Sharma AS, Akhtar Haseeb AH, Syed Abuzar SA. Screening of field pea (*Pisum sativum*) selections for their reactions to root-knot nematode (*Meloidogyne incognita*); 2006.
8. Ali SS, Bansa Singh BS. An overview of nematode constraints in pulses and their management; 2007.
9. Khan MR, Jain RK, Ghule TM, Pal S. Root knot nematodes in India. A comprehensive monograph. All India Co-ordinated Research Project on Plant Parasitic Nematodes with Integrated Approach for their control. Indian Agricultural Research Institute, New Delhi. 2014;29: PP78.
10. Williamson VM, Hussey RS. Nematode pathogenesis and resistance in plants. Te Plant Cell. 1996;8:1735–1745.
11. Dasgupta DR, Gaur HS. The root-knot nematodes *Meloidogyne* spp. in India; 1986.
12. Taylor AL, Sasser JN. Biology, identification and control of root-knot nematodes. North Carolina State University Graphics. 1978;111.
13. Ahmed N, Abbasi MW, Shaukat SS, Zaki MJ. Physiological changes in leaves of mungbean plants infected with *Meloidogyne javanica*. Phytopathologia mediterranea. 2009;48(2):262-268.
14. Suresh P, Poornima K, Sivakumar M, Subramanian S. Current status of root knot nematodes (*Meloidogyne* spp.) in Tamil Nadu. J. Entom. Zool. Studies. 2017;5: 610-615.

15. Hartman KM, Sasser JN. Identification of Meloidogyne species on the basis of differential host test and perineal pattern morphology. An advanced treatise on Meloidogyne. Raleigh: North Carolina State University. Graphics. 1985; 69-77.
16. Bora A, Choudhury BN, Rahman MF. Screening of green gram varieties for resistance against Meloidogyne incognita. Indian Journal Nematology. 2004;34(2): 205-238.
17. Hassan MG, Devi S. Evaluation of mung bean, urd bean, pea, lentil and chickpea germplasm for resistance to root-knot nematode (*Meloidogyne incognita*). Indian Journal Nematology. 2004;34(2): 228-229.
18. Devi G, Choudhary BN, Bhagawati B. Screening of mungbean variety/germplasm against root-knot nematode (*Meloidogyne incognita*) race-2. Indian Journal of Nematology. 2014;44(2):251-252.
19. Pandey RK, Nayak DK. Screening and evaluation of mung bean varieties/ cultivars against root-knot nematode, *Meloidogyne incognita*. International Journal of Research and Review. 2016; 3(6):1-6.
20. Kumar A, Patil JA, Yadav S, Ram S. Screening, confirmation and field evaluation of promising resistant germplasm of different pulses against root-knot nematode, *Meloidogyne javanica*. Journal of Environmental Biology. 2020;41: 1594-1598.
21. Devindrappa Naik, SJ Satheesh, Bohra Abhishek, Singh Bansa, Singh NP. Evaluation of pigeon pea (*Cajanus cajan* L. Milli spp.) genotype against root-knot nematode (*Meloidogyne javanica*). Journal of Food Legume. 2020;32(3):170-173.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124178>