



# Effect of Different Plant Growth Regulators Applied as Foliar Sprays on the Growth and Flowering of Brinjal (*Solanum melongena* L.) under Agro-Climatic Conditions of Chhattisgarh Plains, India

Samiksha Singh Dewangan<sup>a++\*</sup> and Nisha Jangre<sup>a#</sup>

<sup>a</sup> Department of Vegetable Science, College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, India.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i105069>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124753>

Original Research Article

Received: 02/08/2024  
Accepted: 05/10/2024  
Published: 09/10/2024

## ABSTRACT

Present investigation "Effect of different plant growth regulators applied as foliar sprays on the growth and flowering of Brinjal (*Solanum melongena* L.) under agro-climatic conditions of Chhattisgarh plains" in var. Kashi Taru is located at the farm of KVK (Raipur) affiliated with IGKV,

<sup>++</sup> M.Sc. Scholar;

<sup>#</sup> Assistant Professor;

<sup>\*</sup>Corresponding author: E-mail: [samhappytesu@gmail.com](mailto:samhappytesu@gmail.com);

**Cite as:** Dewangan, Samiksha Singh, and Nisha Jangre. 2024. "Effect of Different Plant Growth Regulators Applied As Foliar Sprays on the Growth and Flowering of Brinjal (*Solanum Melongena* L.) under Agro-Climatic Conditions of Chhattisgarh Plains, India". *International Journal of Plant & Soil Science* 36 (10):220-25. <https://doi.org/10.9734/ijpss/2024/v36i105069>.

Raipur, (C.G.) in *Rabi* season 2022-23. The experiment was conducted in RBD (Randomized Block Design) with 3 replications, consisting of 10 treatments of 3 plant growth regulators with different concentrations, including GA<sub>3</sub> (25, 50 and 75 ppm); IAA (25, 50 and 75 ppm) and NAA (25, 50 and 75 ppm) along with control as foliar application at pre flowering stage (30 DAT) and pre fruiting stage (45 DAT). Among the different treatments, the findings indicated that the application of Treatment T<sub>3</sub> i.e. GA<sub>3</sub> @ 75 ppm gave the maximum plant height (89.96 cm), primary branches (11.76) and secondary branches (25.17) per plant over control. The least days taken to 50% flowering were recorded (51.62 days) with GA<sub>3</sub>@ 75 ppm and maximum days taken to 50% flowering were found with control (58.87 days). The highest number of flowers per cluster was recorded in GA<sub>3</sub>@ 75 ppm (4.01), followed by NAA@75 ppm (3.97). Thus, a suitable level i.e., 75 ppm of GA<sub>3</sub>, is effective in enhancing brinjal growth and flowering in the agro-climatic conditions of the Chhattisgarh plains.

**Keywords:** Brinjal; growth; NAA; GA<sub>3</sub>; IAA.

## 1. INTRODUCTION

Brinjal, or eggplant, is a popular vegetable crop. It belongs to the family Solanaceae and is widely distributed in subtropical and tropical countries. According to Whitaker and Stommel [1], brinjal has all the essential vitamins, minerals, nutritional fibre, proteins, and antioxidants. It also contains certain phytochemicals that have scavenging properties. In India, Brinjal covered 7.47 lakh thousand ha with 12.982 million thousand MT of production and productivity of 17.36 thousand MT/ha [2], instead of Chhattisgarh, which has a production area of 37.768 thousand ha and a productivity rate of 18.21 thousand MT/ha, which produces 6.91245 lakh thousand MT [3].

PGR is a broad category of organic substances (other than vitamins and nutrients) that, in very small quantities, alter physiological processes [4]. Growth regulators are organic compounds other than nutrients, a small amount of which is capable of modifying growth [5].

Gibberellic acid (GA<sub>3</sub>), a plant hormone stimulating plant growth and development, is a tetracyclic diterpenoid compound. Gibberellins have been reported to affect almost all plant organs, but their spectacular effects are still on stem elongation [6].

Indole-3-acetic acid (IAA) is produced in the apical region and meristematic tissues of plants, primarily in young leaves. It is a plant bio regulator that controls one or more specific physiological processes within a plant [7]. Indole acetic acid (IAA) has been reported to increase the percentage of long-style flowers in *Solanum khasianum* [8].

The synthetic plant hormone NAA is from the auxin family. NAA is commonly used in

horticulture crops. The higher concentration of NAA inhibits growth and exerts toxic effects on the plant. So, the optimum concentration is the required for the beneficial effect of NAA [9]. Therefore, the present investigation was designed to find out the suitable plant growth regulators for increasing the growth and flowering attributes in brinjal.

## 2. MATERIALS AND METHODS

The present investigation entitled "Effect of different plant growth regulators applied as foliar sprays on the growth and flowering of Brinjal (*Solanum melongena* L.) under agro-climatic conditions of Chhattisgarh Plains" in *var.* Kashi Taru was conducted at the farm of Krishi Vigyan Kendra, Raipur, during the *rabi* season of 2022-2023 under Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.). The seeds of Brinjal variety Kashi Taru sown in nursery were obtained from Krishi Vigyan Kendra, Raipur. The seeds of brinjal after treating with carbendazim @ 3 g/kg were sown in pro-trays on 15 October 2022, which contained coco peat and vermiculite, with water sprinkled on a daily basis. On November 17, 2022, experimental material was transplanted. The number of plants cultivated in the field for each plot is 20, and the total number of plots is 30. Five competitive plants were chosen at random from each plot to be observed on various characteristics. The growth and flowering parameters were recorded, i.e. plant height (cm) at 30, 60 and 90 days after transplanting, number of primary branches per plant and number of secondary branches per plant at harvest, Number of flowers per cluster, days to first, and 50% flowering. The experimental material consisted of ten Brinjal treatments planted in a Randomized Block Design with three replications. The plant growth

regulators *i.e.*, GA<sub>3</sub>, IAA and NAA were used at different concentrations by using foliar sprays (at 30 and 45 DAT). The plants were transplanted in the field at 75 cm row to row and 60 cm plant to plant spacing, with a plot size of 3.5 x 3.2 m. The details of treatment comprised of GA<sub>3</sub> (25, 50 and 75 ppm), IAA (25, 50 and 75 ppm), NAA (25, 50 and 75 ppm) and control (distilled water). The experimental site is the farm of Krishi Vigyan Kendra, Raipur. Raipur is located in the central part of the Chhattisgarh, agro-climatologically known as the Chhattisgarh plains, at an altitude of 289.56 meters above mean sea level (MSL) between 21.25° N latitude and 81.62° E longitude. Raipur is characterized as a slightly moist-hot zone with an average annual rainfall of 1200-1400 mm mainly concentrated between June and September, with occasional light showers during the winter and summer season. Southwestern monsoon receives the largest part of rainfall. May and December are, respectively, the hottest and coolest months of the year. Overall, the highest daily temperature in the summer season runs as high as 46°C and in the winter at least as high as 10°C. The RH is high from July to October and wind velocity is high from May to month of August. In Raipur, the average amount of annual precipitation is 1290 mm.

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant Height (cm)

Result showed that at 30 days after transplanting, the maximum plant height was recorded in treatment T<sub>3</sub> *i.e.* GA<sub>3</sub>@75 ppm (29.15 cm), followed by T<sub>9</sub> *i.e.* NAA@75 ppm (28.98 cm) as compared to other treatments. The minimum plant height was observed in treatment T<sub>10</sub> (20.30 cm) under control. At 60 days after transplanting, significantly maximum plant height was registered in treatment T<sub>3</sub> (61.58 cm), followed by treatment T<sub>9</sub> (60.79 cm). However, minimum plant height was observed in treatment T<sub>10</sub> (46.93 cm) as control. While at 90 days after transplanting, the treatments T<sub>3</sub> (89.96 cm) showed maximum plant height followed by T<sub>9</sub> (88.13 cm) as compared to other treatments. However, treatment T<sub>10</sub> (62.26 cm) as control was recorded with minimum plant height. This may be due to foliar application of GA<sub>3</sub> influence stimulating plant growth and development which ultimately helped in improving the plant height. Similar findings have been reported by Sahu and Choudhary [10] in brinjal and Baby *et.al.* [11] in tomato.

#### 3.2 Number of Primary Branches per Plant

At harvest, the number of primary branches plant<sup>-1</sup> ranged from 8.56 in treatment T<sub>10</sub> (control) to 11.76 in treatment T<sub>3</sub> (GA<sub>3</sub> @ 75 ppm). Therefore, the maximum number of primary branches per plant is seen in Treatment T<sub>3</sub>(GA<sub>3</sub> @ 75 ppm). Similar findings were also observed by Singh *et al.* [12] in brinjal and Prasad *et al.* [13] in tomato.

#### 3.3 Number of Secondary Branches per Plant

The data showed that at harvest, significantly maximum numbers of secondary branches per plant were recorded at harvest in treatment T<sub>3</sub> *i.e.* GA<sub>3</sub> @75 ppm (25.17) followed by treatment T<sub>9</sub> *i.e.* NAA @75 ppm (24.80) as compared to other treatments. While, minimum number of secondary branches per plant observed in treatment T<sub>10</sub> (20.25) as control. Similar findings were also observed by Singh *et al.* [12] in brinjal and Prasad *et al.* [13] in tomato.

#### 3.4 Days to First Flower Initiation

The days to first flower initiation was significantly influenced by different treatments. The first flower initiation was recorded under treatment T<sub>3</sub> *i.e.* GA<sub>3</sub>@ 75 ppm (36.23 days) followed by treatment T<sub>9</sub> *i.e.* NAA @ 75 ppm (36.89 days) as compared to other treatments. While, the late flowering was recorded days in treatment T<sub>10</sub> *i.e.* control (40.28 days). The earliness in the flowering might be due to the fact that GA<sub>3</sub> application enhanced the translocation of food for development of floral primordia, which leads to the early flowering. The above results were in conformity with the application of GA<sub>3</sub> by Patel *et al.* [14] in Brinjal.

#### 3.5 Days to 50 Percent Flowering

The minimum days to 50 percent flowering was noted under treatment T<sub>3</sub> *i.e.* GA<sub>3</sub>@75 ppm (51.62) followed by treatment T<sub>9</sub> *i.e.* NAA@75 ppm (52.44) as compared to other treatments. While, the maximum days to 50 percent flowering was recorded in control in treatment T<sub>10</sub> (58.87). This may be due to GA<sub>3</sub> controls a variety of plant growth and development processes including flower initiation and flower and fruit development. The above results were in conformity with the application of GA<sub>3</sub> by Patel *et al.* [14] in Brinjal.

### 3.6 Number of Flowers per Cluster

A significantly highest number of flowers per cluster was noted in treatment T<sub>3</sub> i.e. GA<sub>3</sub>@ 75 ppm (4.01) followed by treatment T<sub>9</sub> i.e.

NAA@ 75 ppm (3.97) and lowest was observed under the treatment T<sub>10</sub> (2.50) as control. Similar results were found by Ranjeet et al. [15] in tomato and Arivazhagan et al. [16] in brinjal.

**Table 1. Effect of plant growth regulators on plant height in brinjal var. Kashi Taru**

Treatments	Plant Height (cm)		
	30 DAT	60 DAT	90 DAT
GA <sub>3</sub> @25 ppm	26.38	56.87	22.85
GA <sub>3</sub> @50 ppm	28.26	58.57	23.53
GA <sub>3</sub> @75 ppm	29.15	61.59	25.17
IAA @25 ppm	21.76	50.60	20.64
IAA @50 ppm	22.89	51.62	21.28
IAA @75 ppm	23.63	53.73	21.33
NAA @25 ppm	25.42	54.56	22.48
NAA @50 ppm	27.09	57.67	23.07
NAA @75 ppm	28.98	60.79	24.80
Control	20.30	46.93	20.25
SE(m)±	0.60	0.59	0.89
C.D (p=0.05)	1.81	1.78	2.68

\*C.D. = Critical Difference, SE(m) = Standard Error of Mean

**Table 2. Effect of plant growth regulators on growth in brinjal var. Kashi Taru**

Treatments	No. of Primary branches per plant	No. of Secondary branches per plant
GA <sub>3</sub> @25 ppm	10.40	22.85
GA <sub>3</sub> @50 ppm	11.03	23.53
GA <sub>3</sub> @75 ppm	11.76	25.17
IAA @25 ppm	8.80	20.64
IAA @50 ppm	9.03	21.28
IAA @75 ppm	9.53	21.33
NAA @25 ppm	9.23	22.48
NAA @50 ppm	10.16	23.07
NAA @75 ppm	11.13	24.80
Control	8.56	20.25
SE(m)±	0.37	0.26
C.D (p=0.05)	1.13	0.79

\*C.D. = Critical Difference, SE(m) = Standard Error of Mean

**Table 3. Effect of plant growth regulators on flowering parameters in brinjal var. Kashi Taru**

Treatments	Days to 1 <sup>st</sup> flowering	Days to 50% flowering	No. of flowers per cluster
GA <sub>3</sub> @25 ppm	37.29	52.78	3.52
GA <sub>3</sub> @50 ppm	37.54	53.45	3.92
GA <sub>3</sub> @75 ppm	36.23	51.62	4.01
IAA @25 ppm	40.69	58.32	2.56
IAA @50 ppm	39.56	57.08	2.73
IAA @75 ppm	39.07	55.87	2.88
NAA @25 ppm	38.21	54.66	3.22
NAA @50 ppm	38.33	53.80	3.84
NAA @75 ppm	36.89	52.44	3.97
Control	40.28	58.87	2.50
SE(m)±	1.23	0.63	0.49
C.D (p=0.05)	3.70	1.90	1.47

\*C.D. = Critical Difference, SE(m) = Standard Error of Mean



Fig. 1. General view of experimental plot



Fig. 2. Field view of Brinjal experiment during Rabi 2022-23

#### 4. CONCLUSION

In light of the present experimental findings summarized above, it may be concluded that the application of three plant growth regulators i.e. GA<sub>3</sub>, IAA and NAA, GA<sub>3</sub> @ 75 ppm application is very effective for enhancement of growth and flowering traits of brinjal var. Kashi Taru followed by NAA @ 75 ppm. Based on the finding, it is concluded that application of foliar spray of GA<sub>3</sub> @ 75 ppm at 30 DAT and 40 DAT in Brinjal produce significantly higher plant height, number of primary and secondary branches per plant, number of flowers per cluster and gives early flowering.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

I express my gratitude to my advisor Dr. NISHA JANGRE for constant support, guidance and for her valuable suggestions for improving the quality of this research work and to all the faculty members of Department of Vegetable Science, IGKV, Raipur, Chhattisgarh, India for providing necessary facilities, for their cooperation, encouragement and support.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Whitaker BD, Stommel JR Distribution of Hydroxycinnamic acid conjugates in fruit of commercial eggplant (*Solanum melongena* L.) cultivars. J. Agr. Food Chem. 2003;51(11): 3448-3454.

2. Anonymous. National Horticulture Board, Ministry of Farmers and Welfare, Government of India, Statistics, Area and production of horticulture crops, (2<sup>nd</sup> Advance Estimates,1); 2021.
3. Anonymous. Directorate Horticulture and Farm Forestry C.G., Department of Agriculture, Govt. of C.G. Raipur. Horticulture Statistics, Area and production district wise. 2020;5.
4. Wareing PF, Phillips IDJ. The control of growth and differentiation in plants, Edition 2, Oxford and New York, Pergamon. 1978;155-157.
5. Leopold AC. Auxins and Plant Growth Regulator. Berkeley and Los Angeles, University of California Press. 1963;5.
6. Low HK. Rate of gibberellins in root and shoot growth in gibberellins and plant and plant growth, H.N. Wiley Eastern Limited, New Delhi. 1975;101-111.
7. Waheed A, Hamid FS, Ahmad H, Abbassi FM, Aslam S, Shah AH, Ahmad N, Naheed N, Ali H, Khan N. Effect of Indole Butyric acid (IBA) on early root formation (tomato "Sahil" hybrid) cuttings. J. Mat. Env. Sci. 2014;6(1):272–279
8. Ravindran S. Radiobiological studies in Solasodine yielding Solanums. Doctoral thesis. North-Eastern Hill Univ., Shillong, India; 1981.
9. Basnet S, Yadav A, Singh GK, Shekhar. Effect of 1-naphthaleneacetic acid (NAA) on growth and yield of brinjal (*Solanum melongena* L.) cv. Pusa purple cluster Kanpur, India. The Pharma Innovation Journal. 2021;10(9):1235-1238.
10. Sahu NK, Choudhary AS. Effect of plant growth regulators on growth and yield attributes of brinjal (*Solanum melongena* L.). The Pharma Innovation Journal. 2022;11(12):347-349.
11. Baby R, Saravanan S, Prasad VM, Baby S, Geethu BL. Effect of GA<sub>3</sub> and NAA on plant growth and yield of cherry tomato (*Lycopersicon esculentum* var. *cerasiforme*) under polyhouse condition. The Pharma Innovation Journal. 2018;7(7):79-82.
12. Singh, SK, Kumar, A, Beer, K, Singh, VP, & Patel, SK. Effect of naphthalene acetic acid (NAA) and Gibberellic acid (GA<sub>3</sub>) on growth and fruit quality of tomato (*Lycopersicon esculentum* Mill.). Int. J. Curr. Microbiol. App. Sci. 2018;7(3):306-311.
13. Prasad RN, Singh SK, Yadav RB, Chaurasia SNS. Effect of GA<sub>3</sub> and NAA on growth and yield of Tomato. Veg. Sci. 2013;40(2):195-197.
14. Patel S, Verma S, Maurya BK, Maurya SK, Patel Y and Prakash S. Effects of GA<sub>3</sub> and NAA on Growth and Yield of Brinjal (*Solanum melongena* L.) cv. Kashi Sandesh. International Journal of Plant & Soil Science. 2022;6:71-76.
15. Ranjeet, Ram, RB, Prakash, J and Meena, ML. Growth, flowering, fruiting, yield and quality of tomato (*Lycopersicon esculentum* Mill.) as influenced plant bio regulators. Intl. J. Plant Sci. 2014;9(1) 67-71.
16. Arivazhagan, E, Kavitha, A and Kandasamy, R. Influence of plant growth regulators on yield and quality characters of Brinjal (*Solanum melongena* L.) cv. Annamalai. Asian J. Hort. 2018; 13(2): 45-49.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124753>