



Bangladeshi Physicians' Knowledge, Attitude and Perception of Nicotine Replacement Therapy

Samia Amin ^{a*}, Eida Jahra Jarin ^b, Tahsin Munajja ^c,
Farjana Masud ^d, Ishra Jalal ^e, Mashiat Faizah ^e,
Meer Sadad Billah ^f, Oishi Barua ^g and Sumaiya Monjur ^h

^a Macquarie University, Australia.

^b Sylhet Women's Medical College, Bangladesh.

^c BIRDEM General Hospital, Bangladesh.

^d Shaheed Ziaur Rahman Medical College, Bangladesh.

^e Khulna Medical College, Bangladesh.

^f New York Institute of Technology College of Osteopathic Medicine, United States.

^g Chattogram Maa-O Shishu Hospital Medical College, Bangladesh.

^h Dhaka Medical College & Hospital, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. Author SA conceptualized the study, performed the methodology, did formal analysis, wrote the original draft of the manuscript, reviewed and edited the manuscript, supervised the work. Author EJJ wrote, reviewed and edited the manuscript, performed the methodology and did Data collection. Authors TM performed the methodology and did Data collection. Authors FM, IJ, MF, MSB, OB and SM did data collection. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127260>

Original Research Article

Received: 21/09/2024

Accepted: 23/11/2024

Published: 28/11/2024

*Corresponding author: Email: drsamia27@gmail.com;

Cite as: Amin, Samia, Eida Jahra Jarin, Tahsin Munajja, Farjana Masud, Ishra Jalal, Mashiat Faizah, Meer Sadad Billah, Oishi Barua, and Sumaiya Monjur. 2024. "Bangladeshi Physicians' Knowledge, Attitude and Perception of Nicotine Replacement Therapy". *Asian Journal of Research in Nursing and Health* 7 (1):240-45. <https://journalajrn.com/index.php/AJRNH/article/view/175>.

ABSTRACT

Background: Smoking is a major public health concern worldwide, with significant morbidity and mortality. Nicotine Replacement Therapy (NRT) is an effective intervention for smoking cessation. However, in Bangladesh, the adoption of NRT has been limited, partly due to insufficient knowledge and varying attitudes among healthcare providers. This study aims to assess the knowledge, attitudes, and perceptions of Bangladeshi physicians regarding NRT.

Methodology: A cross-sectional survey was conducted among 59 physicians in Bangladesh using a structured questionnaire. The survey evaluated demographic information, knowledge of NRT products, attitudes and perceptions towards NRT as a smoking cessation tool. Data were analyzed using descriptive statistics to summarize responses.

Results: The majority of respondents were male (71.2%) and aged 31-40 years (67.8%). Most had practiced for 5-10 years (88.1%) and held advanced qualifications (MBBS + MD/MS: 79.7%). The average knowledge score was found to be 7.59 out of 15 (50.6%), indicating a moderate level of knowledge regarding NRT among the surveyed physicians. Support for NRT as a first-line treatment was high (77.9%), and 84.7% perceived NRT as safer than smoking. However, there was uncertainty about societal views on smoking cessation without NRT, with 39% unsure.

Implications: This study highlights a strong foundation of knowledge and positive attitudes towards NRT among Bangladeshi physicians, yet underscores the need for further education to address specific knowledge gaps and uncertainties. Furthermore, ensuring the availability of NRT in Bangladesh is crucial for supporting these efforts.

Keywords: *Nicotine replacement therapy; smoking cessation; quit; physician; knowledge; attitudes; perceptions; Bangladesh.*

1. INTRODUCTION

Smoking is a leading cause of preventable morbidity and mortality worldwide, contributing to a wide range of chronic diseases, including cardiovascular disease, respiratory illnesses, and various cancers [1,2]. Nicotine Replacement Therapy (NRT) is widely recognized as an effective tool to help individuals quit smoking by reducing withdrawal symptoms and cravings associated with nicotine addiction [3,4]. NRT includes products such as nicotine patches, gums, lozenges, inhalers, and nasal sprays, which provide a controlled dose of nicotine to ease the quitting process [5].

Despite the global recognition of NRT's efficacy, its adoption and utilization in Bangladesh remain limited. Previous studies have primarily focused on Western countries, where NRT is more readily available and widely used [4,6]. In Bangladesh, smoking cessation programs and the availability of NRT products are less developed, and there is a paucity of research on the knowledge, attitudes, and perceptions of healthcare providers regarding NRT. This gap is significant because physicians play a crucial role in advocating for and administering smoking cessation treatments. Understanding their perspectives is essential for designing effective interventions tailored to the Bangladeshi context.

This study is significant as it addresses the critical need to understand how Bangladeshi physicians perceive NRT, their level of knowledge about its use, and their attitudes towards recommending it to patients. By examining these factors, the study aims to identify potential barriers to consider NRT and provide its insights that could inform the development of targeted training programs and policy initiatives. Enhancing physicians' knowledge and positively shaping their attitudes towards NRT could significantly improve smoking cessation efforts in Bangladesh, and reduce the health burden associated with smoking-related diseases. The primary objective of this study was to assess the knowledge, attitudes, and perceptions of Bangladeshi physicians towards NRT.

2. METHODOLOGY

A descriptive cross-sectional study was conducted on a sample of 59 Bangladeshi physicians to examine the knowledge, attitude, and perception of NRT among Bangladeshi physicians. Utilizing a Google Form survey instrument, participants underwent a comprehensive assessment comprising a knowledge test, attitude evaluation, and perception analysis regarding NRT. The knowledge test aimed to gauge participants'

understanding of NRT's goals, mechanisms of action, product specifics, adverse effects, contraindications, and factors influencing successful cessation. Attitude questionnaire explored/test participants' beliefs about NRT as a primary treatment, opinions on its accessibility and promotion, confidence in recommending it, to influence media on their decision-making. Perception inquiries delved into the impact of NRT on cessation efforts, societal views, concerning the harm of smoking, health provides confidence in discussing NRT with patients, of its use between as in the Western countries. The knowledge test comprised 15 questions, while each of the attitude and perception domains included 5 questions. To calculate the average knowledge score, each correct answer was given 1 point, and each incorrect answer received 0 points. Scores from 0 to 5 were considered low, 6 to 10 were moderate, and 11 to 15 were high knowledge on NRT.

This survey was conducted from January to March 2024, targeting physicians practicing in Bangladesh. The survey was disseminated through professional networks and Bangladeshi medical professional social media (Facebook) groups to ensure a diverse representation of participants. Informed consent was obtained from all participants, and measures such as data encryption, secure password protected storage of the data, restricted access to identifiable information, and anonymization of participant identifiers were taken to ensure the confidentiality and anonymity of respondents' data.

Utilizing descriptive statistics in SPSS, the collected data will be analyzed to provide a comprehensive overview of the knowledge, attitudes, and perceptions of Bangladeshi physicians regarding NRT. Descriptive statistics such as means, frequencies, and percentages was used to summarize the responses to individual survey items and demographic characteristics of the participants.

3. RESULTS AND DISCUSSION

The majority of the respondents were male (71.2%) and between the ages of 31 and 40 years (67.8%). A significant proportion had practiced for 5 to 10 years (88.1%) and held advanced qualifications (MBBS + MD/MS: 79.7%). These demographics suggest that the surveyed group is experienced and well-qualified, which may contribute to their overall

understanding and application of NRT in clinical practice. Prior studies have indicated that physician demographics can influence attitudes towards smoking cessation interventions, with more experienced and specialized physicians likely being more supportive of such measures [7].

Regarding the knowledge test about NRT, Across the 59 respondents, a total of 448 answers were correct, while 437 answers were incorrect. The total score achieved by all respondents was 448 out of the maximum possible score of 885. By dividing the total score of 448 by the number of respondents (N=59), the average knowledge score was found to be 7.59 (50.6%) out of 15; suggesting a moderate level of knowledge regarding NRT among the surveyed physicians, with room for improvement through further education and training. This aligns with the findings of Zhong et al. [8], which emphasize that comprehensive knowledge of NRT mechanisms and adjunctive support significantly enhances the effectiveness of smoking cessation efforts.

The attitude findings suggest a positive inclination towards NRT. The majority of respondents (77.9%) agreed that NRT should be a first-line treatment for smoking cessation and 52.5% agreed that access to NRT should be improved in Bangladesh for effective smoking cessation. This support is critical, as physician endorsement has been shown to increase patient uptake and adherence to NRT [5]. In the absence of NRT availability, respondents believed social media intervention (42.4%) and health education/awareness campaigns (20.3%) should be promoted as alternative methods for smoking cessation in Bangladesh, highlighting the need for a multifaceted approach to smoking cessation. The strong support for NRT aligns with evidence suggesting that NRT, combined with behavioral interventions, is among the most effective smoking cessation strategies [4]. Around one-third of respondents (33.9%) were very confident in recommending NRT as a primary smoking cessation aid to their patients, assuming NRT availability in Bangladesh. Among respondents, 28.8% stated that videos, social media posts, or literature about NRT significantly influenced their decision to consider using NRT for smoking cessation in Bangladesh.

The perception findings highlight concerns about the detrimental impact of NRT unavailability on smoking cessation efforts in Bangladesh. The

majority of respondents (42.4%) perceived the unavailability of NRT as moderately detrimental to smoking cessation efforts in Bangladesh. Total 39% of respondents were unsure about the current societal view of smoking cessation in Bangladesh without access to NRT, while 22% perceived it as indifferent. An overwhelming 84.7% of respondents perceived NRT as safer compared to continued smoking. Nearly half of the respondents (49.2%) were moderately confident in discussing NRT with patients seeking smoking cessation options if NRT were available in Bangladesh. Total 66.1% of respondents perceived NRT as more successful and widely accepted in Western countries such as the United States and United Kingdom [9] compared to its potential use in Bangladesh.

Evidence suggest that senior-year medical students in Middle Eastern country [10-12] demonstrated greater knowledge and awareness of NRT compared to their junior counterparts. This finding could influence Bangladesh by highlighting the importance of incorporating targeted education on NRT into the medical curriculum, particularly for junior medical students. By ensuring that medical students in Bangladesh, regardless of their year, receive adequate training and awareness about NRT, future physicians will be better equipped to address smoking cessation effectively, contributing to improved public health outcomes.

4. STRENGTHS AND LIMITATIONS

A strength of this study is its focus on a diverse group of physicians with varying levels of experience and qualifications, providing insights into the knowledge, attitudes, and perceptions across different healthcare professionals in Bangladesh. However, the study has limitations, including the relatively small sample size and the potential for response bias, as the survey was self-reported. The study was limited to physicians in Bangladesh, which may limit the generalizability of the findings to other countries with different healthcare systems and cultural contexts. Comparative studies across different countries would be beneficial to understand the global applicability of these findings.

5. IMPLICATIONS

The findings of this study have important implications for smoking cessation efforts in Bangladesh. The moderate level of knowledge about NRT among physicians highlights the need

for comprehensive training and educational programs to enhance their understanding of NRT products, their mechanisms of action, and their appropriate use. Additionally, the positive attitudes towards NRT and the recognition of its potential benefits suggest that initiatives to improve access to NRT in Bangladesh could be well-received by healthcare professionals. Given the detrimental impact of NRT unavailability on smoking cessation efforts, Bangladesh should consider including NRT treatment as an available option to support smokers in their cessation journey. Providing access to NRT, along with educational campaigns and healthcare professional training, could significantly strengthen smoking cessation programs in the country.

6. FUTURE RESEARCH SCOPE

Given the high prevalence of smoking in Bangladesh, it is crucial to conduct further research to identify effective strategies for promoting smoking cessation, including the potential role of NRT as a viable cessation aid. Additionally, examining the barriers to the implementation of NRT from both the physician's and the patient's perspectives and addressing these barriers is crucial for improving the uptake and success of NRT. Such research could also inform the development of comprehensive smoking cessation programs tailored to the specific needs and cultural context of the Bangladeshi population.

7. CONCLUSION

This study highlights the critical role of Bangladeshi physicians in promoting smoking cessation through NRT. The findings indicate a strong foundation of knowledge and positive attitudes towards NRT among physicians, but also reveal areas of uncertainty that need to be addressed. By building on these insights, future interventions can be better tailored to support physicians in their efforts to reduce smoking prevalence and improve public health outcomes in Bangladesh. Ongoing education and support for healthcare providers are essential to ensure the successful adoption and implementation of NRT in clinical practice, ultimately leading to better health outcomes for patients.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

DATA AVAILABILITY STATEMENT

Available upon request from the corresponding author via email.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. WHO report on the global tobacco epidemic, 2019: Offer help to quit tobacco use [Internet]. Geneva: World Health Organization; 2020 [cited 2024 Nov 27]. Available: <https://www.who.int/publications/i/item/WHO-NMH-PND-2019.5>
2. U.S. Department of Health and Human Services. The health consequences of smoking—50 years of progress: A report of the Surgeon General [Internet]. Atlanta: Centers for Disease Control and Prevention; 2014 [cited 2024 Nov 27]. Available: <https://doi.org/10.1002/cncr.28695>
3. Cahill K, Stevens S, Perera R, Lancaster T. Pharmacological interventions for smoking cessation: An overview and network meta-analysis. *Cochrane Database Syst Rev.* 2013;(5):CD009329. Available: <https://doi.org/10.1002/14651858.CD009329.pub2>
4. Stead LF, Perera R, Bullen C, Mant D, Hartmann-Boyce J, Cahill K, et al. Nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev.* 2012;11:CD000146. Available: <https://doi.org/10.1002/14651858.CD000146.pub4>
5. Fiore MC, Jaén CR, Baker TB, et al. Treating tobacco use and dependence: 2008 update. Clinical practice guideline [Internet]. Rockville: U.S. Department of Health and Human Services; 2008 [cited 2024 Nov 27]. Available: <https://rc.rcjournal.com/content/resp/53/9/1217.full.pdf>
6. Hajek P, Stead LF, West R, Jarvis M, Hartmann-Boyce J, Lancaster T. Relapse prevention interventions for smoking cessation. *Cochrane Database Syst Rev.* 2013;(8):CD003999. Available: <https://doi.org/10.1002/14651858.CD003999.pub4>
7. Malik M, Javed D, Hussain A, Essien EJ. Smoking habits and attitude toward smoking cessation interventions among healthcare professionals in Pakistan. *J Family Med Prim Care.* 2019;8(1):166-70. Available: https://doi.org/10.4103/jfmpc.jfmpc_230_18
8. Zhong Q, An K, Wu Z, Zhang H, Li S, Zhang L, et al. Knowledge and awareness of nicotine, nicotine replacement therapy, and electronic cigarettes among general practitioners with a special interest in respiratory medicine in China. *Front Med (Lausanne).* 2024;10:1236453. Available: <https://doi.org/10.3389/fmed.2023.1236453>
9. Hammond D, Reid JL, Driezen P, Cummings KM, Borland R, Fong GT, et al. Smokers' use of nicotine replacement therapy for reasons other than stopping smoking: Findings from the ITC Four Country Survey. *Addiction.* 2008;103(10):1696-703. Available: <https://doi.org/10.1111/j.1360-0443.2008.02320.x>
10. Abusamak M, AlQato S, Alfooh HH, Altheeb R, Bazbaz L, Suleiman R, et al. Knowledge, attitudes, practices and barriers of medical research among undergraduate medical students in Jordan: A cross-sectional survey. *BMC Med Educ.* 2024;24(1):23. Available: <https://doi.org/10.1186/s12909-023-05002-9>
11. Almehmadi S, Almehmadi M, Alhajaji F, Alharbi F, Alqorashi S, Alqahtani M, et al. Awareness and perception toward nicotine replacement therapy among medical students at Umm Al-Qura University. *Cureus.* 2023;15(8):e44343.

- Available:<https://doi.org/10.7759/cureus.44343>
12. Ergin A, Uzun SU, Bozkurt AI. Knowledge and attitudes of health professional students on smoking cessation techniques in Turkey. Tob Prev Cessation. 2016; 2(June): 66. Available:<https://doi.org/10.18332/tpc/63821>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127260>