



Prevalence of Methicillin-Resistant *Staphylococcus aureus* on Paper Currency Notes

**Rikhi Ram Marasini^{1*}, Pratikshya Shrestha², Prabhat Dhakal³,
Sukra Raj Shrestha³, Sirjana Adhikari³ and Binita Koirala Sharma³**

¹Department of Microbiology, Prithivi Narayan Campus, Tribhuvan University, Pokhara, Nepal.

²Central Department of Microbiology, Tribhuvan University, Kathmandu, Nepal.

³Department of Microbiology, Janapriya Multiple Campus, Tribhuvan University, Pokhara, Nepal.

Authors' contributions

This work was carried out in collaboration among all authors. Authors PD, SRS, SA and RRM designed the study, performed the statistical analysis and wrote the protocol. Authors PS and RRM wrote the first draft of the manuscript and managed the analyses of the study. Author PD managed the literature searches. Author BKS supervised the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRID/2021/v7i230213

Editor(s):

(1) Dr. Win Myint Oo, SEGi University, Malaysia.

Reviewers:

(1) Shelina Nameirakpam, Saveetha Medical College, India.

(2) Abha Kumari, Amity University, India.

(3) Feleke Eriso Orbalo, Wachemo University, Ethiopia.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/69837>

Original Research Article

Received 14 April 2021

Accepted 19 June 2021

Published 06 July 2021

ABSTRACT

The main objective of this study was to determine the prevalence of Methicillin Resistant *Staphylococcus aureus* (MRSA) in paper currency. The paper currencies in circulation in Pokhara Metropolitan City were inspected. Bills of various denominations (Rs 5, 10, 20, 50, 100, 500 and 1000) were collected from five different locations; namely Food and Vegetable Shop, Bus conductor, Hospital Pharmacy, Butcher Shop and Grocery Shop. Collected sample were cultured and incubated for 24 hours at 37 °C in Brain Heart Infusion (BHI) Broth. The inoculums were further cultured on Mannitol Salt Agar (MSA) and Blood Agar (BA) media to obtain colonies, which were examined and evaluated for various parameters like gram staining and biochemical tests for identification. Then, antibiotic susceptibility test of the isolates was performed using standard procedures. A total of 35 sample of paper currency were processed, all of which showed positive growth. Out of 86 total isolates, 21 (24.42%) were *Staphylococcus aureus* followed by *Coagulase*

*Corresponding author: E-mail: Prashantnepsoul@gmail.com;

Negative *Staphylococci* 19 (22.09%), *Diphtheroids* 14 (16.3%), *Bacillus* spp 13 (15.11%), *Micrococci* 9 (10.46%), *Streptococcus pneumonia* 4 (4.65%), *Viridans Streptococcus* 4 (4.65%) and *Streptococcus pyogenes* 2 (2.32%). The total prevalence of MRSA in this study was 7 (33.33%). Paper currency contaminated with MRSA poses a high threat to those handling the bills as well as the community. Thus, this study suggests proper hygiene measures to be adopted after handling of paper currency to minimize the risk of contamination and emergence of diseases.

Keywords: Contamination; methicillin resistant *Staphylococcus aureus*; paper currency.

1. INTRODUCTION

Paper currency notes are extensively used globally to exchange the purchase of goods and services circulating within the largest population around its surrounding. Circulating and expose of these notes often contaminate from the environment or from the people's hand and unhygienic practice [1] including sneezing and coughing on palm, using mucus for note counting, improper hand washing after using a toilet which ultimately makes currency as an environmental vehicle for the transmission of microorganisms from one person to another [2, 3]. Infection with pathogens through contact with fomites and vehicles is widespread and more serious in the era of the COVID-19 Pandemic. Dirty and moist paper money provide substrate for micro flora which could be transmitted through butchers house, conductors and vendors are often overlooked as enteric disease reservoir [3-5] resulting in various infectious diseases [6-8]. In a developing country like Nepal, people often use paper money than digital payment that may result in the rapid spread of contamination every day leading to serious public health disasters in near future [7].

A previous study showed the presence of a high microbial load on paper currency and coins [3]. However, there is no evidence that the presence of potential microorganisms on paper currency or coins leads to infections. Therefore, this study aims to investigate the prevalence of Methicillin Resistance *Staphylococcus aureus* (MRSA) on paper notes in circulation in selected sites and sources of Pokhara.

2. MATERIALS AND METHODS

A descriptive cross-sectional study was conducted taking altogether 35 samples of paper notes (Rupees 5, 10, 20, 50, 100, 500, and 1000) excluding damaged, really dirty, or physically ripped/torn and polymer bills collected from different sources in Pokhara and analyzed at Microbiology laboratory of Janapriya Multiple

Campus, Pokhara, Nepal during January 2019 to May 2019. The samples were collected randomly from different users from different sources like a butcher shop, grocery shop, vegetable/fruit shop, bus conductors, and hospital pharmacy nearby Prithivi chowk and Janapriya Multiple Campus, Pokhara. During the collection, users were requested to keep those samples in sterile polythene bags. Then, the collected specimens were transported to the research laboratory as soon as possible and processed accordingly.

As reported previously, each sample was placed in 10 mL of BHI broth and subsequently incubated at 37 °C for 24 hours after shaking for 5-10 minutes [3]. After that, the pre-enriched samples were sub-cultured on Mannitol salt agar and Blood agar and incubated aerobically at 37 °C for 24 hours. Further isolation was done by sub-culturing on selective media and nutrient agar to obtain a pure culture.

Bacterial isolates were identified phenotypically based on colony morphology, gram reaction, spore staining, and biochemical tests including catalase, oxidase, urease, and coagulase test. The identified bacterial isolates were taken for the antibiotic susceptibility test by the disc diffusion method. The inoculum was prepared by suspending the organisms into 2 mL of sterile saline (0.9% w/v NaCl) and the turbidity of this inoculum was adjusted to 0.5 McFarland standards. The inoculum was cultured at 37 °C on Mueller Hinton agar (MHA) media with a sterile cotton swab. Bacterial isolates were tested against antibiotics such as Vancomycin (30 µg), Penicillin-G (10 µg), Oxacillin (10 µg), Cefoxitin (30 µg), Chloramphenicol (30 µg), Erythromycin (15 µg), Ampicillin (10 µg) and Co-Trimoxazole (25µg). The zones of inhibition (mm) were measured after 18-24 hours of incubation time. The antibiotic susceptibility was interpreted based on CLSI guidelines [9].

2.1 Screening and Confirmation of MRSA

Further antimicrobial susceptibility test was carried out for *S. aureus* isolates to test their

methicillin susceptibility, using Oxacillin (10µg) and Cefoxitin (30µg) disc placed on Muller-Hinton agar. The zones of inhibition were measured and interpreted according to the Clinical and Laboratory Standards Institute. Cefoxitin is considered more accurate than oxacillin (CLSI 2018). *Staphylococcus aureus*, which showed a zone of inhibition of ≤ 10 mm with Oxacillin (10µg) and zone of inhibition ≤ 21mm with Cefoxitin (30µg) is considered as MRSA on MHA after overnight incubation at 35 °C, were considered as MRSA [9].

3. RESULTS

Among the 35 samples subjected to laboratory investigation, all samples were found to be heavily contaminated by microbial flora. Discrete colonies were isolated and identified as gram-positive bacteria through gram staining. Among the total of 86 gram-positive isolates, 27 (31.4%) were found to be gram-positive rods and 59 (68.60%) were gram-positive cocci as shown in (Fig. 1).

3.1 Distribution of Different Organisms among Gram-Positive Isolates

Out of the total 86 gram-positive isolates, 21 were *Staphylococcus aureus*, 19 were coagulase-negative *Staphylococcus* (CONS), 14 were Diptheroids, 13 were *Bacillus* spp, 9 were *Micrococcus* spp, 4 were *Streptococcus pneumoniae*, 4 were *Viridans streptococcus* and 2 were *Streptococcus pyogenes*.

3.2 Prevalence of gram positive bacteria on different denominations from various sources

A total of 15 (17.45%) isolates were obtained from Rs 5 currency notes followed by 14 (16.28%) in Rs 10 bills, 13 (15.12%) in Rs 50 bills, 12 (16.28%) in Rs 20 bills, 12 (16.28%) in Rs 100 bills, 12 (16.28%) in Rs 500 bills and least 8 (9.30%) in Rs 1000 bills. Denomination of Rs 5 showed the highest gram-positive load while Rs 1000 showed the least.

Similarly, the highest numbers of gram-positive isolates were found in currency notes collected from the butcher shop and grocery shop while the least from the hospital pharmacy. Among the total number of gram-positive isolates, 19

(22.09%) were isolated from the butcher shop, 19 (22.09%) from the grocery shop, 18 (20.93%) from bus conductor, 16 (18.61%) from food and vegetable shop and 14 (16.28%) from hospital pharmacy respectively.

3.3 Antibiotic susceptibility test of Isolated *Staphylococcus Aureus*

Antibiotic susceptibility pattern was determined using the Kirby-Bauer disc diffusion method. An Antibiogram of *S. aureus* revealed a higher degree of susceptibility towards the tested antibiotics. Interestingly, all *S. aureus* isolates were sensitive to Vancomycin (100%), Chloramphenicol (100%), and Co-trimoxazole (100%) while resistant to Penicillin-G and Ampicillin (100%). Meanwhile, 90.48% were sensitive to Erythromycin and 66.67% to Cefoxitin and Oxacillin that means 33.33% *S. aureus* isolates were methicillin resistant isolates as shown in Table 4.

3.4 Distribution of MSSA and MRSA among isolated *S. aureus*

Out of a total of 21 *Staphylococcus aureus*, 14 (66.67%) were found to be methicillin-sensitive *Staphylococcus aureus* (MSSA) and 7 (33.33%) were methicillin-resistant *Staphylococcus aureus* (MRSA) as shown in (Fig. 2).

3.5 Distribution of MRSA According to Sources and Denominations

Out of 21 *S. aureus* isolates, the highest number was isolated from the butcher shop and grocery shop 5 (23.81%) followed by pharmacy and bus conductor 4 (19.1%). Meanwhile, 3 (14.29%) were isolated from food and vegetable shop. Among the 7 methicillin-resistant *S. aureus* (MRSA), 2 (9.52%) isolates were found to be isolated from each food and vegetable shop, butcher shop, and grocery shop whereas MRSA was not detected from pharmacy samples as shown in Table 5.

Similarly, among 7 MRSA isolates, 2 (9.25%) were detected from Rs 5, Rs 50, and Rs 100, 1 (4.76%) from Rs 10 while MRSA was not detected from Rs 20 and Rs 1000 as shown in Table 6.




Fig. 1. Percentage of Gram-positive rod and Gram-positive cocci bacteria among the total gram-positive isolates

Table 1. Distribution of different organisms from isolated gram positive colonies

| S. N | Organisms | % (n= No. of isolates) |
|------|--|------------------------|
| 1 | <i>Staphylococcus aureus</i> | 24.42 (n= 21) |
| 2 | Coagulase Negative Staphylococcus (CONS) | 22.09 (n= 19) |
| 3 | Diphtheroids | 16.30 (n= 14) |
| 4 | <i>Bacillus</i> spp | 15.11 (n= 13) |
| 5 | <i>Micrococci</i> | 10.46 (n= 9) |
| 6 | <i>Streptococcus pneumoniae</i> | 4.65 (n= 4) |
| 7 | <i>Viridans streptococcus</i> | 4.65 (n= 4) |
| 8 | <i>Streptococcus pyogenes</i> | 2.32 (n= 2) |
| | Total | 86 (100%) |

Table 2. Distribution pattern of gram-positive bacteria isolated in paper currency notes of different denominations

| Name of organisms | Denomination and No of isolates | | | | | | |
|---------------------------------|---------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|
| | Rs 5 | Rs 10 | Rs 20 | Rs 50 | Rs 100 | Rs 500 | Rs 1000 |
| <i>Staphylococcus aureus</i> | 5 | 3 | 4 | 4 | 3 | 2 | - |
| CONS | 4 | 2 | 1 | 2 | 2 | 5 | 3 |
| <i>Diphtheroids</i> | 2 | 2 | 3 | 1 | 3 | 1 | 2 |
| <i>Bacillus</i> spp | - | 2 | 2 | 3 | 1 | 3 | 2 |
| <i>Micrococcus</i> | 1 | 3 | 1 | 2 | 1 | - | 1 |
| <i>Streptococcus pneumoniae</i> | 2 | 1 | 1 | - | - | - | - |
| <i>Viridans streptococcus</i> | 1 | - | - | 1 | 1 | 1 | - |
| <i>Streptococcus pyogenes</i> | - | 1 | - | - | 1 | - | - |
| Total | 17.5 (N= 15) | 16.3 (N=14) | 13.9 (N=12) | 15.1 (N=13) | 13.9 (N=12) | 13.9 (N=12) | 9.3 (N=8) |

Table 3. Distribution pattern of gram positive bacteria isolated in paper currency notes collected from different sites

| Name of organisms | Sites and No of isolates | | | | |
|---------------------------------|--------------------------|--------------------|--------------------|--------------------|--------------------|
| | Food and Vegetable Shop | Bus Conductor | Hospital Pharmacy | Butcher Shop | Grocery Shop |
| <i>Staphylococcus aureus</i> | 3 | 4 | 4 | 5 | 5 |
| CONS | 5 | 3 | 4 | 3 | 4 |
| <i>Diphtheroids</i> | 3 | 4 | 2 | 3 | 2 |
| <i>Bacillus spp</i> | 3 | 3 | 1 | 3 | 3 |
| <i>Micrococcus</i> | 1 | 2 | 1 | 2 | 3 |
| <i>Streptococcus pneumoniae</i> | - | 1 | 1 | 1 | 1 |
| <i>Viridans streptococcus</i> | 1 | - | - | 2 | 1 |
| <i>Streptococcus pyogenes</i> | - | 1 | 1 | - | - |
| Total | 16 (18.61%) | 18 (20.93%) | 14 (16.28%) | 19 (22.09%) | 19 (22.09%) |

Table 4. Antibiotic Susceptibility Pattern of *Staphylococcus aureus*

| Antibiotic used | Susceptibility pattern | |
|---------------------------------------|------------------------|---------------|
| | Sensitive (%) | Resistant (%) |
| Vancomycin (VA ₃₀) | 21 (100) | 0 |
| Erythromycin (E ₁₅) | 19 (90.48) | 2 (9.52) |
| Penicillin-G (P _{10 units}) | 0 (0) | 21 (100) |
| Ampicillin (AMP ₁₀) | 0 (0) | 21 (100) |
| Co-Trimoxazole (COT ₂₅) | 21 (100) | 0 |
| Chloramphenicol (C ₃₀) | 21 (100) | 0 |
| Cefoxitin (CX ₃₀) | 14 (66.67) | 7 (33.33) |
| Oxacillin (OX ₁₀) | 14 (66.67) | 7 (33.33) |




Fig. 2. Distribution of MSSA and MRSA among isolated *S. aureus*

Table 5. Distribution of MRSA according to different sample sites

| S. N | Sampling site | <i>Staphylococcus aureus</i> | MRSA |
|-------------|-------------------------|-------------------------------------|-------------------|
| 1. | Food and Vegetable shop | 3 (14.29%) | 2 (9.52%) |
| 2. | Bus Conductor | 4 (19.05%) | 1 (4.76%) |
| 3. | Pharmacy | 4 (19.05%) | 0 (0%) |
| 4. | Butcher shop | 5 (23.81%) | 2 (9.52%) |
| 5. | Grocery Shop | 5 (23.81%) | 2 (9.52%) |
| | Total | 21 (100%) | 7 (33.33%) |

Table 6. Distribution of MRSA according to different denominations

| S. N | Denominations | <i>S. aureus</i> | MRSA |
|-------------|----------------------|-------------------------|-------------------|
| 1. | Rs 5 | 5 (23.81%) | 2 (9.25%) |
| 2. | Rs 10 | 3 (14.29%) | 1 (4.76%) |
| 3. | Rs 20 | 4 (19.05%) | - |
| 4. | Rs 50 | 4 (19.05%) | 2 (9.25%) |
| 5. | Rs 100 | 3 (14.29%) | 2 (9.25%) |
| 6. | Rs 500 | 2 (9.52%) | - |
| 7. | Rs 1000 | - | - |
| | Total | 21 (100%) | 7 (33.33%) |

4. DISCUSSION

In this study, often all Nepali paper currency had contaminants. This finding was similar to studies conducted in Saudi Arabia [7]. The result was evident that the paper currency notes act as a significant carrier of human pathogens and transfer harmful pathogens through them.

This study demonstrated the high level of bacterial contamination on paper notes with the higher number of gram-positive bacteria. Similar findings were also recorded on Indian rupees [8], Bangladesh Teka [10], Ghanaian [11], Nepal [12], Pakistan [13], Uganda[14], and Saudi Riyal [7]. This study demonstrates the highest number of bacterial isolates as well as *S. aureus* on samples collected from the butcher shop and Grocery shop 19 (22.09%), which is also supported by the study conducted in India [2,15], and Bangladesh [10]. The variation in the number of positive samples from different sources may be connected with differences in the hygiene of the currency owners and sanitary conditions in the environment. The high level of bacterial contamination of currency notes could be attributed to the fact that they are being used frequently by a very high number of people [14].

The prevalence of MRSA on Butcher and vegetable shop 2 (9.52%) was consistent with

the study conducted by the Khanal group [16]. Meanwhile, detection of MRSA in Rs 5, 10, and 100 but not in Rs 20, Rs 500, and Rs 1000 may be due to more circulation and frequent use for the trade of these denominations than the latter one among the people.

The findings of this research work provide an important clue of dissemination of MRSA in the environment through the circulating paper money posing a public health threat. Therefore, further study should be done to validate the results of the present study and need to take action to reduce the spread of pathogenic micro flora through paper currency.

5. CONCLUSION

The presence of a high load of microbial contamination and Methicillin Resistance *Staphylococcus aureus* in paper currency suggests that banknotes play an important role in the causation and spreading of potentially harmful diseases. This also clearly indicates that pathogenic microorganisms are circulating among people through paper currency in day-to-day life. This is an important finding as the spread of pathogenic bacteria could be minimized through possible preventive measures such as maintaining good hygienic behavior among money handlers.

CONSENT

It is not applicable

ETHICAL APPROVAL

It is not applicable

ACKNOWLEDGMENTS

We are grateful to Janapriya Campus, Pokhara, Nepal for providing microbiology laboratory facilities during this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Basavarajappa KG, Rao PNS, Suresh K. Study of bacterial, fungal and parasitic contamination of currency notes in circulation. Indian J Pathol Microbiol. 2005;48(2):278-279.
2. Barolia SK, Verma S, and Verma BK. Coliform contamination on different paper currency in Ajmer, Rajasthan, India. Univers J Environ Res Technol. 2011;1(4):552-556.
3. Lamichhane J, Adhikary S, Gautam P, Maharjan R, Dhakal B. Risk of handling paper currency in circulation chances of potential bacterial transmittance. Nepal J Sci Technol. 1970;10:161-166.
4. Staley C, Dunny GM, Sadowsky MJ. Environmental and animal-associated enterococci. Adv Appl Microbiol. 2014;87:147-186.
5. Sharif M and Ansari F. Assessment of bacterial load on Pakistani currency notes and coins in circulation. European J Biomed Pharm Sci. 2016;3(8):133-145.
6. Ayandele SA and Adeniyi AA. Prevalence and Antimicrobial resistance pattern of microorganisms isolated from naira notes

- in Ogbomoso North, Nigeria. Journal of Research in Biology. 2011;1(8):587-593.
7. Al-Ghamdi AK, Abdelmalek SM, Bamaga MS, Azhar EI, Wakid MH, Alsaied Z. Bacterial contamination of Saudi 'one' Riyal paper notes. Southeast Asian J Trop Med Public Health. 2011;42(3):711-716.
 8. Bhat N, Bhat N, Bhat S, Asawa K, Agarwal A. An Assessment of Oral Health Risk Associated with Handling of Currency Notes. Int J Dent Clin. 2010;2(3):16.
 9. Clinical and laboratory standards institute. Performance standards for antimicrobial susceptibility testing; twenty-eight informational supplement. CLSI Document M100-S8. Wayne, PA 19087, USA. 2018;8(3):27-39.
 10. Shakir M, Ahmed U, Parveen S, Nasreen T, Feroza B. Evaluation of the microbial contamination of bangladesh paper currency notes (taka) in circulation. Adv Biol Res. 2010;4(5):266-271.
 11. Alemu A. microbial contamination of currency notes and coins in circulation: A Potential Public Health Hazard. Biomed Biotechnol. 2014;2(3):46-53.
 12. Prasai T, Yami KD, Joshi DR. Microbial Load on Paper/Polymer Currency and Coins, Nepal J Sci Technol. 1970;9:105-109.
 13. Ejaz H, Javeed A, Zubair M. Bacterial contamination of Pakistani currency notes from hospital and community sources. Pak J Med Sci. 2018;34(5):1225-1230.
 14. Allan M, Atuhaire C, Nathan M, Ejobi F, Cumber SN. Bacterial contamination of Ugandan paper currency notes possessed by food vendors around mulago hospital complex, Uganda. Pan Afr Med J. 2018;31:1-7.
 15. Sruthi S, Panchmal GS, Shenoy RP, Kumar V, Jodalli P, Somaraj V. Assessment of microbial contamination of indian currency notes in circulation – An In vitro study. J Indian Assoc Public Health Dent. 2020;18(2):179.
 16. Khanal LK, Adhikari RP, Guragain A. Prevalence of methicillin resistant staphylococcus aureus and antibiotic susceptibility pattern in a tertiary hospital in Nepal. J Nepal Health Res Council. 2018;16(2):172-174.

© 2021 Marasini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/69837>*