



Assessment of Hematological and Serum Electrolytes Effects of Intermittent Fasting on Mice

**David Chibuikwe Ikwuka^{a,b*}, Bond Ugochukwu Anyaehie^b, Eghosa E. Iyare^b,
Princewill Ugwu^b, Amechi Uchenna Katchy^c, Gabriel Ejike Igbokwe^d,
Jide Uzowulu Uzoigwe^b, Emmanuel Nonso Ezeokafor^a
and Kelechi Love Ezeudensi^a**

^a *Department of Human Physiology, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Nigeria.*

^b *Department of Physiology, University of Nigeria Enugu Campus, Enugu, Nigeria.*

^c *Department of Anatomy, University of Nigeria Enugu Campus, Enugu, Nigeria.*

^d *Department of Applied Biochemistry, Nnamdi Azikiwe University Awka, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors DCI and BUA designed the study, wrote the protocol and the first draft of the manuscript. Authors DCI, ENE and KLE were involved in the laboratory analysis. Authors DCI, PU, JUU and ENE performed the statistical analysis.

Authors EEI, AUK, GEI and PU managed the analysis of the study. All authors contributed to the preparation of the final draft of the manuscript, read and approved the final manuscript for publication.

Article Information

DOI: 10.9734/JPRI/2021/v33i62B35188

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80415>

Original Research Article

Received 05 November 2021

Accepted 28 December 2021

Published 29 December 2021

ABSTRACT

Introduction: There is growing interest on trying to explore the importance of fasting to health as it has been practiced as religious exercises for centuries. This study assessed the haematological and serum electrolytes effects of intermittent fasting (IF) on mice.

Methods: Fifty (50) male mice were randomly assigned into five (5) groups A, B, C, D and E (n=10). Group A (control) was fed normal chow *ad libitum*, experimental groups B,C,D and E were fasted seven weeks intermittently for 12, 18, 24 and 48 hours respectively. Blood samples were collected and analysed for haematological and electrolyte indices. Data obtained were analyzed using IBM SPSS Statistics 25.

Results: Our findings revealed no significant differences in the RBC, HGB, HCT, MCV, MCH and MCHC between the different fasting groups and the control. There was increase significantly in

group C for Red density width-coefficient of variance (RDW-CV) and group C and E for Red density width-standard deviation (RDW-SD). Platelet count decreased, plateletcrit increased significantly in group C. There were no significant differences in the mean platelet volume (MPV), platelet distribution width (PDW), platelet-large cell count (P-LCC) and Platelet-large cell ratio (P-LCR). Sodium ion decreased significantly in group C and chloride ion decreased significantly in group B, C and E. There were no significant differences in serum potassium ion and bicarbonate ion.

Conclusion: 12, 24 and 48 hours IF are safe and do not negatively influence hematological indices and electrolyte levels but 18 hours IF could have a slight negative effect on platelet count, plateletcrit and sodium ion levels.

Keywords: Intermittent fasting; diseases; weight; electrolytes; platelet; red blood cell.

1. INTRODUCTION

In recent times, lifestyle changes and behavioural modifications have been suspected to have long lasting effects on health and personal well-being. The type of the food we eat and its frequency, the quality of sleep we get, and exercise habits could all influence our general body physiology ranging from the cardiovascular, to gastrointestinal, to musculoskeletal, to respiratory to neurophysiology and to defence physiology. Dietary changes is a very important category of lifestyle changes which can help one loss excess weight and reduce the risk of developing diseases.

Lots of interest, time and resources have been channelled towards how to eliminate excess fat, prevent unhealthy weight gain and contain our body weight due to the link associating excess body weight to exacerbation or complication of certain diseases and as a predisposing factor for diseases. One of the useful and practicable ways to prevent and eliminate excess body fats is through fasting. Hippocrates correctly said, "When you are ill and eat, you are giving fuel to the raging sickness within you" [1]. Saint Nikolai of Zicha [2] stated that fasting makes man joyful and courageous whereas, gluttony makes him gloomy and fearful.

The human body has been designed by the Creator to fast and man has advanced at times of food scarcity to become a product of feast and famine [2]. Living organisms generally alternate between periods of feeding and fasting throughout their existence [3]. Humans in particular, often fast during night sleep periods between 6 to 12 hours daily. However, longer periods of fasting has been widely practised for various purposes globally for thousands years [4]. Fasting is described as a wilful or voluntary abstinence from feeding for certain duration [5],

while, intermittent fasting (IF) is the pattern of alternating feeding and fasting. Alternate day fasting, routine periodic fasting and daily time-restricted feeding are forms of IF which has been studied as a practice to improve metabolic health and many other physiological and molecular markers of health and longevity [6]. Intermittent fasting as a dieting concept is achieved by restricting food intake for a period of time. It has been an important religious practice for Muslims, Christians and other religions during the Ramadan, Lent and other religious seasons.

The most common researched IF is the Ramadan fasting. Ramadan is the ninth month of the Islamic calendar and is observed as a month of fasting for Muslims adult and about a billion Muslims faithful globally observe this injunction by abstaining from eating, drinking and other physical needs such as smoking and sexual intercourse daily during Ramadan from sunrise to sunset [4,7,8]. In similar vein, Christians globally observe Lent in preparation for Easter. Lent is a 40 days period which begins on Ash Wednesday and ends the night before Easter Sunday. Amongst the activities during the Lent includes fasting, praying, almsgiving etc [9]. Fasting during the Ramadan month is the most popular and extensively studied IF for its numerous health benefits.

Haematological indices are important in diagnosing the structural and functional status of animals due to its sensitivity to environmental and physiological changes [10]. Several haematological changes affecting red blood cell (RBCs), white blood cells (WBCs), and coagulation factors have been shown to be linked with obesity associated comorbidities [11]. Wojciak [12] posited that short term fasting significantly decreases iron concentrations in hair and serum as well as levels of ferritin, haematocrit, haemoglobin and red blood cells.

There is growing interest on trying to explore the importance of fasting to health and researchers have identified fasting as a strong non-pharmacological, non-genetic tool to enhance multi-systemic protection against stress and to delay several aspects related to the onset, rate and progression of age-related chronic diseases, mainly due to its anti-inflammatory and antioxidant activities [13]. However, despite established potential health benefits of fasting in ageing-related disease prevention and promoting physiological effects, little has been known about the effect of different fasting duration regimen on haematological indices and immunological function. Therefore, the study will aim to determine the effect of different intermittent fasting regimen on haematological indices, electrolyte levels and immunological function in mice.

Intermittent fasting has been reported to ameliorate diabetic retinopathy and increase insulin sensitivity [14-16]; improve varied indicators of cardiovascular health [17-22]; prevent nephrotoxicity [23]; lower the risk of developing age-related diseases [24-25]; increase lifespan in a number of organisms [24]; good in preventing neurodegeneration in the brain [26-32]; and attenuate and purge precancerous and cancerous cells [24,33-35].

A few studies on IF have been able to evaluate red blood cell parameter and white blood cell parameters among fasters during the month of Ramadan [36-38] but the reports are however conflicting and controversial, hence, the need for this study.

2. MATERIALS AND METHODS

2.1 Research Design

Animal experimental research design was used for this study.

2.2 Experimental Animals

Fifty (50) five to six weeks old Matured male pathogen-free mice weighing between 17-20g were purchased from the Vivarium of Department of Zoology & Environmental Biology, University of Nigeria Nsukka. They were housed in standard polypropylene cages with wood chip bedding at $25\pm 2^{\circ}\text{C}$ and $60\pm 5\%$ relative humidity under a 12-hour light/ 12-hour dark cycle (6am to 6pm Light). Prior to the experiment initiation, they

were acclimatized for two week so as to effectively recover from the stress of transport and change in environment. The general health statuses of the mice were evaluated via weight measurement. The polypropylene cages and accessory equipment including feeders and watering devices were washed and autoclaved regularly to keep them clean and free from contamination before use.

2.3 Experimental Procedure

The experiment was conducted in the Animal house of the Faculty of Basic Medical Sciences, College of Medicine, University of Nigeria Enugu Campus. After the acclimatization, they were weighed and distributed randomly ($n = 10$ per group) into Control groups (A) and IF groups (B, C, D and E). To investigate the impact of different IF regimen on haematological and electrolyte indices, the mice in the fasting group were divided into four sub-groups according to the duration of the fasting: 12hr, 18hr, 24hr and 48hr ($n = 10$ per group). Group A (control) was fed normal chow *ad libitum*, group B was fasted for 12 hours daily, group C was fasted for 18 hours daily, group D was on 24 hour alternate day fasting, and group E was fed one day out of every three days. The time for feeding and fasting were based on group regimen. After the seven weeks of intermittent fasting, fasting was stopped. The animal cages were cleaned every seven days to keep the animals dry and clean throughout the experiment duration.

After the experimental duration, all the mice were anesthetized and blood samples collected via retro-orbital sinus and analysed for haematological and electrolyte indices.

2.4 Full Blood Count Testing

The blood was processed for routine haematological testing immediately after sample collection using the Symex XN-1000 Hematologic Analyser (Symex Corporation) in the Molecular Laboratory of Nnamdi Azikiwe University Awka, Nigeria. The parameters tested included, haemoglobin concentration, hematocrit level, erythrocyte count, leucocyte count, leucocyte differential count (neutrophils, lymphocytes, basophils, monocytes, eosinophils and large unstained cell counts) and platelet count.

2.5 Data Analysis

Data were presented in means \pm standard error of mean (SD \pm SEM). The differences in body weight, red blood cell parameters, white blood cell counts, platelet parameters and serum electrolyte levels were determined by one-way Analysis of Variance (ANOVA), followed by Tukey's post hoc test for multiple comparisons (IBM SPSS version 25). Our statistically significant value was set at 2-sided $p < 0.05$.

3. RESULTS

Prior to the experiment, body weight did not differ between groups. At the end of the experiment, mortality was higher in the control group than the IF groups.

Table 1 show the mean values of the different intermittent fasting groups from week one to week 6. From the result, there is no observable difference of the control group from the experimental groups.

From Table 2 above showing the results of mean red blood cell count, haemoglobin concentration, hematocrit, mean corpuscular volume, mean corpuscular haemoglobin, mean corpuscular haemoglobin concentration, red density width-coefficient of variance and red density width-standard deviation for groups A, B, C, D and E, it is observed that there is no significant difference in the mean RBC between group A (Control) and groups B, C, D and E. The result also shows that there is no significant difference in mean haemoglobin concentration between group A and groups B, C, D and E, no significant difference in mean hematocrit between group A and groups B, C, D and E, no significant difference in mean corpuscular volume between group A and groups B, C, D and E, no significant difference in mean corpuscular haemoglobin between group A and groups B, C, D and E, and no significant difference in mean corpuscular haemoglobin concentration between group A and groups B, C, D and E. However, for red density width-coefficient of variance and red density width-

standard deviation there was an increase which was statistically significant in experimental group C as compared to group A.

From the result in Table 3, there is no statistically significant difference in the MPV, PDW, P-LCC and P-LCR between group A and groups B, C, D and E. The result from Table 3 shows a decrease in PLT in experimental groups (B, C, D and E) which was only statistical significant in group C. The result also shows decrease in PCT in the experimental groups (B, C, D and E) which was only statistically significant in group C.

Data in Table 4 presents the mean values of the serum electrolytes of Groups A, B, C, D and E respectively. From the Table, there was decrease in serum sodium ion which was significant only in group C relative to the control. There was also decrease in chloride ion which was significant in groups B, C and D respectively in relative to the control group. There were no significant differences between the mean serum levels of potassium ion and bicarbonate ion of the experimental groups (B, C, D and E) as compared to the control (A).

4. DISCUSSIONS

Research on intermittent fasting (IF) is growing in popularity globally. Few evidences have shown the adaptive effects of physiological fasting in promoting health and in the treatment and prevention of metabolic diseases [39]. This study appears to be the first study to investigate the effects of different IF regimen on haematological parameters, and electrolytes level in mice. Lifestyle modifications have been found to have significant impact on human health, decreasing the risk for cardiovascular related issues, inhibiting growth of malignant conditions and impacting ageing. On the effects of different IF regimen on mean mice weight, the results showed that there is no statistically significant difference. Although other reports had declared decrease in body weight gain during fasting period [3,40-41], it is unclear as to why we did not record a significant decrease following extended period of IF.

Table 1. Showing the effect of different fasting regimen on mice weight

Weight	Group A Mean \pm SEM	Group B Mean \pm SEM	Group C Mean \pm SEM	Group D Mean \pm SEM	Group E Mean \pm SEM
Initial	19.83 \pm 1.01	21.63 \pm 1.24	21.40 \pm 0.97	23.90 \pm 1.37	22.50 \pm 0.53
Final	36.93 \pm 0.42	32.56 \pm 0.88	29.70 \pm 0.77	36.29 \pm 1.13	34.66 \pm 1.11

Table 2. Showing the effect of different fasting regimen on Red Blood Cell parameters

Parameter	Groups	Mean±SEM	ANOVA (Sig)	Post hoc (Tukey)	F-ratio
RBC ($\times 10^{12}/L$)	A	9.70±0.04	0.680		0.581
	B	9.36±0.15		0.993	
	C	8.91±0.48		0.861	
	D	8.98±0.49		0.898	
	E	8.57±1.07		0.638	
HGB (g/dl)	A	13.30±0.17	0.959		0.153
	B	12.70±0.33		0.992	
	C	12.52±1.29		0.978	
	D	12.98±0.74		0.999	
	E	12.34±1.53		0.954	
HCT (%)	A	48.32±0.50	0.878		0.294
	B	47.28±0.95		0.999	
	C	45.12±3.95		0.953	
	D	46.60±2.60		0.995	
	E	43.92±5.32		0.867	
MCV (fl)	A	49.80±0.32	0.264		1.419
	B	50.48±0.63		0.997	
	C	50.24±2.05		0.999	
	D	53.90±2.03		0.257	
	E	51.50±0.81		0.904	
MCH (pg)	A	13.72±0.12	0.510		0.851
	B	13.58±0.32		0.999	
	C	13.90±0.82		0.998	
	D	14.46±0.28		0.770	
	E	14.46±0.38		0.770	
MCHC (g/dl)	A	27.54±0.16	0.613		0.681
	B	26.92±0.40		0.915	
	C	27.56±0.60		1.000	
	D	27.30±0.83		0.997	
	E	28.10±0.36		0.939	
RDW-CV (%)	A	18.90±0.21	0.001		7.729
	B	19.78±0.69		0.893	
	C	23.58±1.03*		0.001	
	D	19.24±0.14		0.997	
	E	19.46±0.89		0.978	
RDW-SD (fl)	A	29.70±0.23	0.004		5.575
	B	31.66±0.91		0.218	
	C	33.76±0.71*		0.002	
	D	31.70±0.44		0.202	
	E	32.54±0.64*		0.033	

RBC=Red blood cell, HGB=Hemoglobin, HCT=Hematocrit, MCV=Mean corpuscular volume, MCH=Mean corpuscular haemoglobin, MCHC=Mean corpuscular haemoglobin concentration, RDW-CV=Red density width-coefficient of variance, RDW-SD= Red density width-standard deviation. *= Statistically significant

In this study, we observed no significant difference in the RBC, HGB, HCT, MCV, MCH and MCHC between the different fasting groups in comparison with the control. This finding is in line with the reports of Akrami et al. [42], Plumelle et al. [43] and Turan et al. [44] who observed no statistically significant difference between fasting and postprandial levels of these haematological parameters. However, the findings of this study disagrees with the report of Nematy et al. [45] who observed higher RBC count following Ramadan fasting and Naderi et

al. [46], and Koscielnak et al. [47] who posited that fasting decreased RBC counts, Hb and HCT. The study also disagrees with the findings of Attarzadeh [37] and Ahmed [38] who reported increase significantly for Hb and HCT in Ramadan type IF. There was increase significantly in group fasted for 18 hours for Red density width-coefficient of variance (RDW-CV) and groups fasted intermittently for 18 and 48 hours for Red density width-standard deviation (RDW-SD). Ahmed [38] similarly reported an increase in RDW-CV and RDW-SD during

Table 3. Showing the effect of different fasting regimen on platelets

Parameter	Groups	Mean±SEM	ANOVA (Sig)	Post-hoc (Tukey)	F-ratio
PLT ($\times 10^9/L$)	A	1106.60±30.30	0.023	0.188	3.608
	B	736.60±131.40			
	C	592.20±49.3*			
	D	695.60±82.24			
	E	997.00±193.97			
MPV (g/dl)	A	6.16±0.07	0.112	0.228	2.147
	B	7.14±0.19			
	C	7.32±0.67			
	D	7.12±0.06			
	E	6.70±0.14			
PDW (%)	A	15.54±0.08	0.612	0.999	0.682
	B	15.68±0.08			
	C	15.68±0.30			
	D	15.86±0.07			
	E	16.42±0.88			
PCT (fl)	A	0.70±0.02	0.028	0.383	3.399
	B	0.52±0.09			
	C	0.37±0.01*			
	D	0.59±0.07			
	E	0.66±0.11			
P-LCC (pg)	A	38.00±4.21	0.703	0.993	0.548
	B	46.00±9.86			
	C	65.00±24.60			
	D	45.40±13.53			
	E	45.80±4.49			
P-LCR (g/dl)	A	3.42±0.30	0.368	0.973	1.135
	B	6.08±0.74			
	C	12.12±6.82			
	D	7.88±0.64			
	E	5.16±0.97			

PLT=Platelet, MPV=Mean platelet volume, PDW=Platelet distribution width, PCT=Plateletcrit, P-LCC=Platelet-large cell count, P-LCR=Platelet-large cell ratio, *= Statistical significant

Table 4. Showing the effect of different fasting regimen on serum electrolyte levels

Parameter	Groups	Mean±SEM	ANOVA (Sig)	Post-hoc (Tukey)	F-ratio
Sodium ion- Na^+ (mE/L)	A	162.82±7.64	0.041	0.282	3.050
	B	151.42±3.80			
	C	144.48±0.70*			
	D	148.02±1.97			
	E	152.44±0.80			
Potassium ion (K^+)	A	7.38±1.66	0.564	0.959	0.759
	B	6.36±0.52			
	C	6.04±0.24			
	D	8.05±1.52			
	E	5.91±0.54			
Bicarbonate ion (HCO_3^-)	A	24.58±1.75	0.384	0.298	1.100
	B	20.20±0.24			
	C	22.48±1.45			
	D	21.56±1.54			
	E	22.84±2.07			
Chloride ion (Cl)	A	122.59±2.90	0.004	0.025	5.442
	B	114.00±2.30*			
	C	111.30±0.70*			
	D	114.98±1.18			
	E	113.98±1.06*			

*=statistically significance

Ramadan fast. High RDW may be indicative of nutrient deficiency such as a deficiency of iron, folate, or vitamin B-12. This translates that exposure to IF without specialized nutrient rich diet intake at feeding intervals could predispose one to having high red cell distribution width.

From our findings, we found out that there was decrease in the platelet count of the fasting groups which was only significant in the group fasted intermittently for 18 hours. Akrami et al. [42], Koscielnak et al. [47] and Turan et al. [44] similarly reported that fasting decreased platelet count after Ramadan fasting. There was increase in plateletcrit value which was significant only in group fasted intermittently 18 hours daily for seven weeks. Our findings also showed that there is no significant difference in the mean platelet volume (MPV), platelet distribution width (PDW), platelet-large cell count (P-LCC) and Platelet-large cell ratio (P-LCR) between the control group and fasting groups. Recent researches have suggested that platelet may have diagnostic and prognostic roles use in certain diseases [48]. Our finding identified decrease platelet count and increase in plateletcrit in an IF group and Gao et al. [49] suggested that PCT is nonlinearly correlated to platelet count.

The findings for serum electrolytes showed a slight decrease in sodium ion level and chloride ions. This decrease in sodium ion is significant in the group fasted for 18 hours and the decrease in chloride ion is only significant in groups fasted intermittently for 12, 18 and 48 hours respectively. Findings reported in literature on influence of fasting on sodium and chloride ion levels have been inconsistent and contrasting. Sedaghat et al. [50] similarly found a significant decrease in sodium ion following Ramadan fasting while; Kenenisa et al. [40] reported that sodium ion levels increased however insignificantly and chloride ion increased significantly following Ethiopian Orthodox Christians fasting. There was no significant difference in serum potassium ion and bicarbonate ion between the control and fasting groups. Literature on serum potassium ion during fasting is however conflicting. Sedaghat et al. [50] indicated that serum potassium ion decreased during Ramadan fasting. In a conflicting report, Kenenisa et al. [40] found that serum potassium ion increased following Ethiopian Orthodox Christian fasting.

5. CONCLUSION

This study thus concludes that different intermittent fasting regimen does not impact RBC results but may affect its distribution width, does not impact MPV, PDW, P-LCC and P-LCR but 18 hours IF could decrease platelet count and plateletcrit. Intermittent fasting for 18 hours was shown decrease sodium and chloride ion concentration. The present study has demonstrated that 12, 24 and 48 hours IF are safe and do not negatively influence hematological indices and electrolyte levels but 18 hours IF could have a slight negative effect on platelet count, plateletcrit and sodium ion levels. However, a larger study using specific nutrient rich diet and extended duration are required to duplicate our findings is recommended.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

Not applicable

ETHICAL APPROVAL

The study protocol was approved and conducted in accordance with the rule for animal care as stipulated by the College of Medicine Ethics Committee, University of Nigeria Enugu Campus, Enugu, Nigeria (050/07/2021) which is in compliance with the experimental guidelines of the U.S. National Institute of Health (NIH) and Institutional Animal Ethics Committee (IAEC) on the care and use of laboratory animals.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tiller JB. Weight loss through fasting, use of spices and tasty diets. E-book. 2008; 104.

2. Mosley M, Spencer M. The fast diet. 2nd ed. London. Short book. 2019;210.
3. Toledo FW, Grundler F, Goutzourelas N, Tekos F, Vassi E, Mesnage R, Kouretas D. Influence of long-term fasting on blood redox status in humans. *Antioxidants*. 2020; 9(496):1-15. Available:<http://dx.doi.org/10.3390/antiox9060496>
4. Li L, Su Y, Li F, Wang Y, Ma Z, Li Z, Su J. The effects of daily fasting hours on shaping gut microbiota in mice. *BMC Microbiol*. 2020;20:65. Available: <https://doi.org/10.1186/s12866-020-01754-2>
5. Hannan MA, Rahman MA, Rahman MS, Sohag AAM, Dash R, Hossain S, Farjana M, Uddin MJ. Intermittent fasting, a possible priming tool for host defense against SARS-CoV-2 infection: crosswalk among calorie restriction, autophagy and immune response. *Immunology Letters*. 2020;226:38-45. Available:<https://doi.org/10.1016/j.imlwt.2020.07.001>
6. Horne BD, Muhlestein JB, Anderson JL. Health effects of intermittent fasting: hormesis or harm? A systematic review. *Am J Clin Nutr*. 2015;102:464–70.
7. Adawi M, Diamiani G, Bragazzi NL, Bridgewood C, Pacifico A, Conic RRZ, et al. The Impact of Intermittent Fasting (Ramadan Fasting) on Psoriatic Arthritis Disease Activity, Enthesitis, and Dactylitis: A Multicentre Study. *Nutrients*. 2019; 11:601 Available:<http://dx.doi.org/10.3390/nu11030601>
8. Abunada T, Abunada H and Zayed H. Fasting Ramadan During COVID-19 Pandemic: Immunomodulatory Effect. *Front. Nutr*. 2020;7:557025. DOI: 10.3389/fnut.2020.557025
9. Little Shoots Deep Roots. Christie Thomas; 2021. 40 Lent activities for families (that are simple and fun). [Updated 2021 March 1, cited 2021 Oct 17]; Available:<https://littleshootsdeeproots.com/kids-ideas-blog/40-lent-ideas-kids/>
10. Ikwuka DC, Nwobodo Ed, Anyaehie BU, Umegbolu EI. Hematological and histological effect of IRACARP® (fractionated neem leaf extract) in healthy wistar rats model. *Physiol and Pharmacol*. 2020;24:314-321. Available:<http://dx.doi.org/10.32598/ppj.24.4.60>
11. Biadgo B, Melku U, Abebe SM, Abebe M. Hematological indices and their correlation with fasting blood glucose level and anthropometric measurements in type 2 diabetes mellitus patients in Gonder, Northwest Ethiopia. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 2016;9:91-99. Available:<https://doi.org/10.2147/DMSO.S97563>
12. Wojciak RF. Effect of short-term food restriction on iron metabolism, relative well-being and depression symptoms in healthy women. *Eat Weight Disord*. 2014;19:321–327. Available: <https://doi.org/10.1007/s40519-013-0091-2>
13. Rojas-Morales P, Tapia E, León-Contreras JC, González-Reyes S, Jiménez-Osorio AS, Trujillo J, et al. Mechanisms of fasting-mediated protection against renal injury and fibrosis development after ischemic acute kidney injury. *Biomolecules*. 2019; 9:404 Available:<https://doi.org/10.3390/biom9090404>
14. Wan R, Camandola S, Mattson MP. Intermittent food deprivation improves cardiovascular and neuroendocrine responses to stress in rats. *J. Nutr*. 2003; 133:1921–1929.
15. Furmli S, Elmasry R, Ramos M, Fung J. Therapeutic use of intermittent fasting for people with type 2 diabetes as an alternative to insulin. *BMJ Case Rep*. 2018;bcr2017221854.
16. Sutton EF, Beyl R, Early KS, Cefalu WT, Ravussin E, Peterson CM. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes. *Cell Metab*. 2018;27: 1212–1221.
17. Harvie MN, Wright C, Pegington M, McMullan D, Mitchell E, Martin B, et al. The effect of intermittent energy and carbohydrate restriction v. Daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br. J. Nutr*. 2013;110:1534–1547.
18. Moro T, Tinsley G, Bianco A, Marcolin G, Pacelli QF, Battaglia G, et al. Effects of eight weeks of time-restricted feeding (16/8) on basal metabolism, maximal strength, body composition, inflammation, and cardiovascular risk factors in

- resistance-trained males. *J. Transl. Med.* 2016;14:1–10.
19. Catenacci VA, Pan Z, Ostendorf D, Brannon S, Gozansky WS, Mattson MP, et al. A randomized pilot study comparing zero-calorie alternate-day fasting to daily caloric restriction in adults with obesity. *Obesity.* 2016;24:1874–1883.
 20. Antoni R, Johnston KL, Collins AL, Robertson MD. Intermittent v. continuous energy restriction: Differential effects on postprandial glucose and lipid metabolism following matched weight loss in overweight/obese participants. *Br. J. Nutr.* 2018;119:507–516.
 21. Stekovic S, Hofer SJ, Tripolt N, Aon MA, Royer P, Pein L, et al. Alternate day fasting improves physiological and molecular markers of aging in healthy, non-obese humans. *Cell Metab.* 2019;30:462–476.
 22. Pinto AM, Bordoli C, Buckner LP, Kim C, Kaplan PC, Del Arenal IM, et al. Intermittent energy restriction is comparable to continuous energy restriction for cardiometabolic health in adults with central obesity: A randomized controlled trial; the Met-IER study. *Clin. Nutr.* 2020;39:1753–1763.
 23. Gunebakan E, Yalcin E, Dulger EC, Yigitbasi A, Ates N, Caglayan A, et al. Short-term diet restriction but not alternate day fasting prevents cisplatin-induced nephrotoxicity in mice. *Biomedicines* 2020;8:23. Available:<https://doi.org/10.3390/biomedicines8020023>
 24. Longo VD, Panda S. Fasting, Circadian Rhythms, and Time-Restricted Feeding In Healthy Lifespan. *Cell Metab.* 2016;23:1048–1059.
 25. Longo VD, Mattson MP. Fasting: molecular mechanisms and clinical applications. *Cell Metab.* 2014;19:181–192.
 26. Halagappa VKM, Guo Z, Pearson M, Matsuoka Y, Cutler RG, Laferla FM, Mattson MP. Intermittent fasting and caloric restriction ameliorate age-related behavioral deficits in the triple-transgenic mouse model of Alzheimer's disease. *Neurobiol. Dis.* 2007;26:212–220.
 27. Madorsky I, Opalach K, Waber A, Verrier JD, Solmo C, Foster T, et al. Intermittent fasting alleviates the neuropathic phenotype in a mouse model of Charcot-Marie-Tooth disease. *Neurobiol. Dis.* 2009;34:146–154.
 28. Tajes M, Gutierrez-Cuesta J, Folch J, Ortun˜o-Sahagun D, Verdaguer E, Jimenez A, et al. Neuroprotective role of intermittent fasting in senescence-accelerated mice P8 (SAMP8). *Exp. Gerontol.* 2010;45:702–710.
 29. Singh R, Lakhanpal D, Kumar S, Sharma S, Kataria H, Kaur M, Kaur G. Late-onset intermittent fasting dietary restriction as a potential intervention to retard age-associated brain function impairments in male rats. *Age (Dordr.)* 2012;34:917–933.
 30. Shimazu T, Hirschey MD, Newman J, He W, Shirakawa K, Le Moan N, et al. Suppression of Oxidative Stress by β -Hydroxybutyrate, an Endogenous Histone Deacetylase Inhibitor. *Science.* 2013;339:211–214.
 31. Hu D, Xie Z, Ye Y, Badijri S, Chen M. The beneficial effects of intermittent fasting: an update on mechanism, and the role of circadian rhythm and gut microbiota. *Heepato Biliary Surg Nutr*; 2020. Available:<http://dx.doi.org/10.21037/hbsn-20-317>
 32. Mattson MP, Moehl K, Ghena N, Schmaedick M, Cheng A. Intermittent metabolic switching, neuroplasticity and brain health. *Nat. Rev. Neurosci.* 2018;19:63–80.
 33. Meynet O, Ricci JE. Caloric restriction and cancer: Molecular mechanisms and clinical implications. *Trends Mol. Med.* 2014;20:419–427.
 34. Mattison JA, Colman RJ, Beasley TM, Allison DB, Kemnitz JW, Roth GS, et al. Caloric restriction improves health and survival of rhesus monkeys. *Nat. Commun.* 2017;8:14063
 35. Nencioni A, Caffa I, Cortellino S, Longo VD. Fasting and cancer: Molecular mechanisms and clinical application. *Nat. Rev. Cancer.* 2018;18:707–719.
 36. Unalacak M, Kara IH, Baltaci D, Erdem O, Bucaktepe PG. Effects of Ramadan Fasting on Biochemical and Hematological Parameters and Cytokines in Healthy and Obese Individuals. *Metab Syndr Relat Disord.* 2011;9:156-61.
 37. Attarzadeh Hosseini SR, Hejazi K, Nikroo H. The Effects of Ramadan Fasting and Physical Activity on Blood Hematological-Biochemical Parameters. *Iran J Basic Med Sci.* 2013;16:845-9.
 38. Ahmed OA. Effect of ramadan fasting on red and white blood cell parameters in healthy females. *GMJ.* 2019;30:1-3.

- Available:<http://dx.doi.org/10.12996/gmj.2019.01>
39. Mayer AL, Zhang Y, Feng EH, Higgins CB, Adenakan O, Pietka TA, et al. Enhanced hepatic PPAR α activity links GLUT8 deficiency to augmented peripheral fasting responses in male mice. *Endocrinology*. 2018;159(5):2110-2126. Available:<https://doi.org/10.1210/en.2017-03150>
40. Kenenisa ECH, Degef Teklemariam M, Mekonen W, Haile Wondimu D, Sisay T, Genet Gebre S. The Effect of Ethiopian Orthodox Christians 'AbiyTsom' (Lent fasting) on Metabolic Syndrome Indices and Serum Electrolytes. *J Nutrition Fasting Health*. 2018;6(2):60-70. Available:<https://doi.org/10.22038/JNFH.2018.35602.1153>
41. Wang S, Huang M, You X, Zhao J, Chen L, Wang L, et al. Gut microbiota mediates the anti-obesity effect of calorie restriction in mice. *Sci. Rep*. 2018;8:13037.
42. Akrami MF, Ahmadi Z, Hassanshahi G, Akrami ME, Ravari A, Ghalebi SR. Dose ramadan fasting affects inflammatory responses: evidences for modulatory roles of this unique nutritional status via chemokine network. *Iran J Basic Med Sci*. 2013;16:1217-22.
43. Plumelle D, Lombard E, Nicolay A, Portugal H. Influence of diet and sample collection time on 77 laboratory tests on healthy adults. *Clin Biochem*. 2014;47:31-7.
44. Turan E, Tunç R, Turan Y. Comparison of fasting and postprandial levels of commonly used biochemical and hematological parameters. *Arch Clin Exp Med*. 2019;4(2):66-71.
45. Nematy M, Alinezhad-Namaghi M, Rashed MM, Mozhdehifard M, Sajjadi SS, Akhlaghi S, et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J*. 2012;10:11-69.
46. Naderi Ghalenoie Z, Hessami Arani S, Kerachian M. Fasting and experimental animals, detrimental or positive? *Journal of Nutrition, Fasting and Health*, 2015;3(4): 169-176. Available:<https://doi.org/10.22038/jfh.2015.6073>
47. Kościelniak BK, Charchut A, Wójcik M, Sztefko K, Tomasiak PJ. Impact of fasting on complete blood count assayed in capillary blood sample. *Laboratory Medicine* 2017;48:4:357-361. Available:<https://doi.org/10.1093/labmed/lmx044>
48. Budak YU, Polat M, Huysal K. The use of platelet indices, plateletcrit, mean platelet volume and distribution width in emergency non-traumatic abdominal surgery: A systematic review. *Biochemia Med (Zagreb)*. 2016;26(2):178-193. Available:<https://doi.org/10.11613/BM.2016.020>
49. Gao Y, Li Y, Yu X, Guo S, Ji X, Sun T, et al. The impact of various platelet indices as prognostic markers of septic shock. *PLoS One*. 2014;9(8):e10376.
50. Sedaghat MR, Askarizadeh SF, Heravian J, Rakhshandadi T, Nematy M, Mahmoodi Z, et al. The Effects of Islamic Fasting on Blood Hematological Biochemical Parameters. *J Fasting Health*. 2017;5(2): 56-62. Available:<http://dx.doi.org/10.22038/jfh.2017.22778.1085>

© 2021 Ikwuka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/80415>