



Comparative Assessment of Farmer's Brinjal Variety in Different Agro-Climatic Conditions

**Hardev Choudhary^{a*}, Parthkumar P. Dave^a,
Noushad Parvez^a, Mahesh B. Chodvadiya^a,
Prajana Ritaparna^a, Laishram Yelhounganba Khuman^a,
Chriss Ipe^a and Leelaram Sahu^a**

^a National Innovation Foundation-India (Autonomous Body of the Department of Science and Technology, Government of India), Amrapur, Gandhinagar-382650, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i21660

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96567>

Original Research Article

Received: 21/11/2022

Accepted: 29/01/2023

Published: 23/02/2023

ABSTRACT

In order to evaluate the performance and suitability of the long brinjal variety 'Niranjan Bhata' developed by an innovative farmer of Chhattisgarh in different agro-climatic conditions of the country, the comparative field experiment conducted under Gandhinagar, Gujarat condition was accommodated in a randomized complete block design comprising five replications of four treatments, i.e. *Niranjan Bhata*, *Pusa Shamla* as National Check (NC) and two Local Check (LC) during June - March 2016. The plant height of *Niranjan Bhata* (115.50 cm) was recorded at par with LC-1 (118.33 cm). However, the maximum fruit length (45.80 cm) was recorded in *Niranjan Bhata* (45.80 cm). The significant highest fruit yield was produced by *Niranjan Bhata* (31.40 t/ha). Farmer managed trials (FMTs) were also carried out in five states between 2017-18 and 2020-21 at

*Corresponding author: E-mail: hardevnif@gmail.com, hardev@nifindia.org;

68 locations (loc.) during the *Kharif* season. The test variety '*Niranjan Bhata*' reported superior performance in Chhattisgarh with the highest plant height (161.17 cm), fruit length (59.67 cm), and marketable fruit yield (9.81 kg/plant). The variety also outperformed with similar characters in Gujarat and Maharashtra states over checks. As per the growers' feedback, the variety was found to be high-yielding. The long fruits containing lesser seeds and soft pulp having good market preference too. Thus the evaluation of variety with the farmers' participation was found most suitable approach which assured the varietal preference and its suitability for large-scale dissemination.

Keywords: *Brinjal; eggplant; Solanum melongena; Niranjan Bhata; farmer-managed trials; long fruit.*

1. INTRODUCTION

Brinjal (*Solanum melongena* L.) is the most important solanaceous crop cultivated in subtropical and tropical areas. Globally, India is the second largest producer after China contributing 24.5% with 12.9 Million tonnes in global production [1]. A range of wild species of Eggplant (*Solanum melongena*) is found abundantly in the country, considered to be a rich source of genetic diversity for breeding programs, particularly for the characteristics associated with climate change adaptability [2]. The indigenous genotypes and local landraces of brinjal were superior in yielding to other economic traits and resistance to pests and diseases, which are considered the most desirable traits for crop improvement programs [3,4]. The replacement of traditional, regionally adapted crop varieties with high-yielding and hybrid varieties threatens global agrobiodiversity, resulting in collective losses of the crop, variety, and allele diversity [5]. Despite this fact, wild relatives of crops are highly under-represented in the worldwide eggplant genetic resource conservation system. Farmers require improved eggplant varieties for sustainable production and adaptability against the various challenges due to climate change [6,7].

Although such studies so far, very less substantial work has been undertaken to promote potential traditional landraces and farmers' developed brinjal varieties at farmers' fields. As a result, in light of the importance of such studies, comparative performance evaluation trials at Gandhinagar, Gujarat and farmer-managed trials to evaluate the performance of a long-fruited farmer's brinjal variety '*Niranjan Bhata*' with ascertaining farmers' assessment of the variety for desirable traits in different agro-climatic zones of the country were undertaken. The test variety was developed through the mass

selection method based on fruit length with other characteristics i.e. lesser number of seeds, prickles on the stem, and calyx with resistance to major insect pests and diseases by an innovative farmer of Dhamtari district of Chhattisgarh from a traditional cultivar conserved by his forefathers for the last 70 years [8]. The innovator for his significant contribution to the crop improvement of brinjal has been conferred with an award by then Hon'ble President of India Shri Pranab Mukherjee at Rashtrapati Bhavan, New Delhi during the Festival of Innovation-2017 organized by National Innovation Foundation (NIF) - India. The application for registration of the variety under PPV&FR, Act 2001 has also been filed (REG/2017/1797) at Protection of Plant Varieties and Farmers' Rights Authority, New Delhi.

2. MATERIALS AND METHODS

The field trial was conducted at NIF's research farm, Gandhinagar district of Gujarat, situated at 23.223°N latitude and 72.650°E longitude conditions for the performance evaluation of farmer's brinjal variety '*Niranjan Bhata*' in comparison with a national check (NC) and two local checks (LC) during June - March 2016. The trial was accommodated in a randomized complete block design comprising four treatments and five replications. The local checks were procured from the agri-input shop from Gandhinagar after interaction with the progressive farmers of the region, while a national check (*Pusa Shyamla*) was collected from Seed Division, ICAR-Indian Institute of Agricultural Research, New Delhi. The cultivation practices of the brinjal as recommended for a particular area for a good crop stand were adopted.

In order to evaluate the performance and ascertain farmers' participation for the assessment of variety on its claimed traits viz.

fruit length with plant height, and fruit yield per plant in different agro-climatic conditions, the criteria defined by Witcombe et al. [9] for the Farmer-managed trials (FMTs) with slight modifications were adopted for performance and suitability assessment of the variety. The FMTs were also undertaken by the NIF team from the headquarter, Gandhinagar (Gujarat), NIF-Cell, Guwahati (Assam) for North East Region and NIF-cell, Bhubneswar (Odisha) at 68 locations covering five states Chhattisgarh, Gujarat, Odisha, Kerala and Manipur during *Kharif* 2016-17 and 2017-18. Similarly, comparative FMTs were also conducted at ten locations in each Gujarat and Maharashtra states during *Kharif* 2020-21. Under the supervision of NIF's experts, the seeds of the test variety (*Niranjan Bhata*) are only being produced and maintained by the innovator. NIF provided seeds of the variety to selected potential brinjal growers and acquainted them with its cultivation practices. The observations were recorded from ten randomly selected plants and/or plant parts and subjected to statistical analysis using standard protocols. This FMTs study can be used to facilitate multi-location on-farm trials by involving more farmers to generate reliable data on its uniformity and stability for further crop improvement programs. Such trials will also help us to devise appropriate strategies for its dissemination in suitable agro-climatic conditions on a large scale.

3. RESULTS AND DISCUSSION

The comparative study conducted at NIF research farm, Gujarat during June - March 2016 revealed that the *Niranjan Bhata* variety reported a higher plant height of 115.50 cm, found statistically at par with LC-1 recorded with a maximum plant height of 118.33 cm among all test varieties. However, the fruit length of *Niranjan Bhata* variety (45.80 cm) was recorded as significantly superior. In the same way, the significantly highest marketable fruit yield was recorded in *Niranjan Bhata* with 31.40 t/ha (Table 1).

The higher fruit yield of *Niranjan Bhata*, brinjal variety might be attributed also by higher plant height and more fruit length. The fruit yield has a significant and positive correlation with plant height [10,11], completely support the present result obtained. A similar significant and positive association of plant height and fruit length with fruit yield was also revealed in

some studies [12,13] which is also consistent with the present result obtained. In the study of Bhushan et al. [14], the highest fruits yield (per plant & per hectare) was obtained in *Pusa Shyamla*, indicating the higher yield of farmer's variety and partially supporting the present findings.

3.1 Farmer-Participatory on-Farm Trials

The data generated and analyzed from the research trial revealed that the brinjal variety, *Niranjan Bhata* fulfilled multiple requirements, including agronomic performance and quality production during the trials. Therefore, in view of the above, it was envisaged to evaluate the variety at farmers' fields to check its performance and suitability in various locations across India. Thus, the data recorded and analyzed from the farmer-managed trials conducted during *Kharif* 2016-17 and 2017-18 at a total of 68 locations (Loc.) covering five states, e.g. Manipur (Loc 4), Chhattisgarh (Loc 48), Gujarat (Loc 08), Kerala (Loc 3) and Odisha (Loc 5) is depicted in Table 2 and Annexure. NIF provided seeds of the test variety to selected farmers who voluntarily agreed to participate in the study and the farmers adopted cultivation practices of the respective region, including plant protection measures. The researchers posted in different locations by NIF collected the observations and farmers' feedback from the trials. The data presented showed that the *Niranjan Bhata*, brinjal variety outperformed in Chhattisgarh with the highest plant height (161.17 cm), fruit length (59.67 cm), and yield (9.81 kg/plant). Similarly, Chhattisgarh recorded highest fruit length (59.67 cm) and followed by Odisha, Manipur, Kerala, Gujarat, and Maharashtra. On the same way, the maximum fruit yield was recorded in Chhattisgarh (9.81 kg/plant) followed by Maharashtra (6.46 kg/plant), Gujarat (5.83 kg/plant), Odisha (5.10 kg/plant), Manipur (4.00 kg/plant), and Kerala (2.50 kg/plant). In unavailability of long-fruited local check at test locations, the trait-specific observations for the farmer's variety were recorded and presented.

Where in the comparative results of FMTs in Gujarat and Maharashtra showed the superiority of *Niranjan Bhata* over local checks. It was recorded 9.39%, 55.10%, and 34.81% higher plant height, fruit length, and fruit yield, respectively as compared to local check in Gujarat, and a similar trend was also observed in Maharashtra with 16.40%, 49.00%, and

35.45% increase in plant height, fruit length and fruit yield in Maharashtra (Table 3). These results revealed that the *Niranjan Bhata* variety is most suitable for *river basins Agro-eco region and Hot Moist/Dry Sub-humid Transitional ESR region* of Chhattisgarh, which gave a better result in the *hot arid/ semi-arid eco-region* including plains of Maharashtra and Gujarat states. It was also found suitable for the *hot moist sub-humid eco sub-region* of Odisha. However, it performed lower in the *North-Eastern Hills Warm Perhumid Eco-sub*

regions of Manipur. It may not perform better in the *hot subhumid - humid to perhumid eco-subregion* of Kerala (ICAR-Agro-Ecological Sub Region).

The performance of variety at farmers' fields mainly depends upon the various agro-climatic conditions of the respective area. Therefore, the variation in plant height, fruit length, and fruit yield might be influenced by the weather and growing environment in respective states at different locations.

Table 1. Performance evaluation of *Niranjan Bhata* in comparison to checks under the Gandhinagar condition

Brinjal varieties	Plant height (cm)	Length of fruit (cm)	Fruit yield (t/ha)
Niranjan Bhat	115.50	45.80	31.40
<i>Pusa shyamla</i>	86.60	19.30	26.98
Local check-1	118.33	22.50	22.62
Local check-2	75.23	17.20	22.56
SEm \pm	1.27	0.71	2.55
CD (p=0.05)	3.91	2.19	7.85
CV%	2.87	6.07	2.20

Note: The data presented are the mean of five replications and recorded from 10 plants/plant parts; CD: critical difference; SEm: standard error of the mean; CV: coefficient of variation

Table 2. Performance of *Niranjan Bhata* brinjal variety under FMTs conducted in five states in the *Kharif* season between 2016-17 and 2017-18

States	Plant height (cm)	Fruit length (cm)	Yield per plant (kg)
Manipur (Loc 4)	108.75	53.00	4.00
Chattisgarh (Loc 48)	161.17	59.67	9.81
Gujarat (Loc 08)	97.17	41.83	5.83
Kerala (Loc 3)	109.67	38.00	2.50
Odisha (Loc 5)	107.80	54.00	5.10
Mean	116.91	49.30	5.45
SD	25.25	9.04	2.74
SEm \pm	11.29	4.04	1.23

Note: The data presented were the mean of data recorded from 10 plants/plant parts from respective locations

Table 3. Comparative assessment of *Niranjan Bhata* brinjal variety under FMTs conducted at ten locations in each Gujarat and Maharashtra state during *Kharif* 2020-2021

States	Plant height (cm)			Fruit length (cm)			Yield per plant (kg)		
	<i>Niranjan bhatan</i>	Local check	PIOC	<i>Niranjan bhatan</i>	Local check	PIOC	<i>Niranjan bhatan</i>	Local check	PIOC
Gujarat	99.12	89.81	9.39	35.30	15.85	55.10	5.85	3.82	34.81
Maharashtra	103.47	86.5	16.40	37.06	18.9	49.00	6.46	4.17	35.45
Mean	101.30	88.16		36.18	17.38		6.16	3.99	
SD	3.08	2.34		1.24	2.16		0.43	0.25	
SEm \pm	2.18	1.66		0.88	1.53		0.30	0.18	

Note: The data presented were the mean of data recorded from 10 plants/plant parts from respective locations where Loc.: Location; PIOC: Percent Increase Over Control

i. Comparative study of *Niranjan Bhata* conducted during *Kharif* 2016-17 under Gandhinagar (GJ) condition.



Photo 1. View of the fruit length & pulp appearance of *Niranjan Bhata* at experimental field

ii. FMTs in different agro-climates



Photo 2. FMTs in Gujarat 2016-17 and



Photo 3. FMTs in Chattishgarh-2017-18

A similar finding of the significant role of temperature and relative humidity in biometric and yield parameters such as plant height, number of leaves, leaf area index, fruit length, girth, and weight, including soil fertility improvement which favours growth and productivity of the crop, was revealed by Shikha et al. [15] which partially supports variation in yield found in the present findings.

As per the feedback recorded from growers, the farmer's brinjal variety, *Niranjan Bhata* found a high yielder and bears more fruits per plant. The longer fruits contained lesser seeds with soft pulp and were found to be most suitable for the 'sabzi' and "Sambhar" preparation due to their good taste. The consumer preference for the brinjal fruits is based on shape, size, colour and flavour,

including ripeness with less seeded fruit pulp [16]. The fruits of this variety also fetch a higher market price due to long fruit preferences. The beneficiaries of Maharashtra, Odisha, and Chhattisgarh have been able to enhance their income due to attractive fruit appearance, higher yield, and long fruit size. Local landraces and traditional cultivars still contribute one-third of the cultivation of brinjal

crops in India, which might be due to locale specification performance, and consumer-friendly attributes of the cultivars [17] found consistent with present findings. Indigenous vegetable cultivars provide nutritional and financial security for small farmers and women. The result is inconsistent with the findings of Singh et al. [18].



Photo 4. FMTs in Manipur



Photo 5. FMTs in Kerala 2016-17



Photo 6. FMTs in Odisha

iii. FMTs in Gujarat and Maharashtra



Photo 7. FMTs in Gujarat



Photo 8. FMTs in Maharashtra 2020-21

The present study gives a choice to the farmers that they can adopt the improved farmers' developed crop varieties, which are better adapted to wider climatic conditions. A similar finding was revealed by Kaushik et al. [2] for wild species of eggplant, which was found in concordance with the present findings.

As per the phenotypic observation of the farmers' field and noticed by the growers of this variety that *Niranjan Bhata* possesses long fruits having spines on shoot and calyx was reported to lower the incidence of pests and diseases. The similar morphological characteristics prevented the insect from egg-laying on shoot and fruits, as reported by Gautam et al. [19] and the Local long fruited cultivars *Pusa Purple Long - 74*, including *Navkiran* were observed with the low shoot and fruit infestation by Mathur et al. [20] which is in line with the present study.

Hence, it is a thrust to work on Host Plant Resistance and the identification of markers for this brinjal variety, *Niranjan Bhata*. It shall also

be a prominent variety for developing high-yielding, pests-resistant varieties. The local landraces and farmers' plant varieties are a rich source of genetic material for the trait-specific development of varieties for researchers. They would facilitate the improvement of local genotypes with greater adaptability to local conditions with the capability of performing better with higher yield [21].

4. CONCLUSION

Superiority in desirable traits fruit length and yield of the farmers developed brinjal variety - *Niranjan Bhata* in comparison to Pusa Shyamla and local checks makes it potential variety for wider dissemination. The FMTs also concluded the variety as most preferred for different agro-climatic conditions i.e. Chhattisgarh, Maharashtra, Gujarat, and Odisha. It also outperformed during comparative FMTs conducted at the tribal and hilly regions of Gujarat and Maharashtra. It made is a more reliable and promising planting material for such resource-poor farmers for additional

income generation from small land holdings. As per the growers' feedback, the variety had minimum pest occurrence where lesser seeds with soft pulp made it more preferred for cooking purpose. The performance recorded with such preferred characteristics can increase the varietal adoption among small landholders and make it a potential landrace for crop improvement programs too.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. IndiaStat. Available: <https://www.indiastat.com/table/agriculture/area-production-productivity-brinjal-india-1987-19/14880> Access on 15.1.2023
2. Kaushik P, Prohens J, Vilanova S, Gramazio P, Plazas M. Phenotyping of eggplant wild relatives and interspecific hybrids with conventional and phenomics descriptors provides insight for their potential utilization in breeding. *Front. Plant Sci.* 2016;7:677.
3. Rajan N, Debnath S, Dutta AK, Pandey B, Singh AKR. Characterization of indigenous brinjal (*Solanum melongena* L.) lines using morphological traits under Jharkhand condition. *Ann. Plant Soil Res.* 2020;22(4):425-431.
4. Sowmya E, Pradeep S. Screening of Traditional Brinjal (*Solanum melongena* L.) varieties against Cotton Leaf Hopper, *Amrasca bigutulla bigutulla* (Ishida). *Int. J. Curr. Microbiol. App. Sci.* 2020;9(08): 3513-3524.
5. Ficiciyan AM, Loos J, Tschardtke T. Similar yield benefits of hybrid, conventional, and organic tomato and sweet pepper varieties under well-watered and drought-stressed conditions. *Front. Sustainable. Fd. System.* 2021;5: 628537.
6. Taher D, Solberg SO, Prohens J, Chou Yu-yu, Rakha M, Wu Tien-hor. World vegetable center eggplant collection: Origin, composition, seed dissemination and utilization in breeding. *Front. Plant Sci.* 2017;8:1484.
7. Kumar N, Dave PP, Mir PA, Zulapi L, Choudhary H. Dissemination of elite brinjal variety hzkb-1 under organic cultivation in Uttarakhand and Jammu & Kashmir through farmers' participation. *Int. J. Environ. Clim. Change* 2022;12(12):1202-1207.
8. NIF - India. Innovation of the day. *Niranjan Bhata*: A longer and sweeter variety of brinjal developed by a grassroots innovator. 2017. Available: <https://nif.org.in/Innovationofday/Niranjan-Bhata-A-longer-sweeter-variety-of-brinjal-developed/1> Accessed on 18.12.2021
9. Witcombe JR, Joshi A, Joshi KD and Sthapit BR. Farmer participatory crop improvement. I: varietal selection and breeding methods and their impact on biodiversity. *Exp. Agric.* 1996;32(4):445-460.
10. Dasmohapatra A, Sharma D. Correlation and path coefficient analysis in long fruited brinjal (*Solanum melongena* L.). *Int. J. Pure Appl. Biosci. SPI:* 2018;6(3): 400-406.
11. Onyia VN, Chukwudi UP, Ezea AC, Atugwu AI, Ene CO. Correlation and path coefficient analyses of yield and yield components of eggplant (*Solanum melongena*) in a coarse-textured Ultisol. *Info. Processing Agr.* 2020(7): 173-181.
12. Angadi P, Indiresh KM, Rao AM. Correlation studies for fruit yield and its attributing characters in brinjal (*Solanum melongena* L.). *Int. J. Curr. Microbiol. Appl. Sci.* 2017;6(12):1007-1012.
13. Gurve VR, Waskar DP, Khandare VS, Mehtre SP, More DG. Studies on correlation and path analysis for yield and its contributing traits in brinjal (*Solanum melongena* L.) *Int. J. Curr. Microbiol. Appl. Sci.* 2020;9(10): 179-188.
14. Bhushan A, Samnotra RK, Kumar S, Chopra S, Kumar S, Kumar M. Effect of Seasons on Fruit Yield, major insect pests and diseases in selected cultivars of brinjal under jammu conditions (J&K). *Int. J. Curr. Microbiol. App. Sci.* 2020;11: 3148-3159.
15. Shikha E, Kunnathadi M, Shahida K, Sreelekshmi S, Jubail E. Assessment of environmental factors on growth of brinjal under open and protected conditions. *Int. J. Curr. Microbiol. Appl. Sci.* 2020;9(4): 380-386.
16. Sitorus LT, Henriques PD, Lucas MR. Consumer Preferences of tomato and

- eggplant in Dili, East Timor. Int. J. Biz. Mktg. and Mgmt. (IJBMM). 2020;5(3): 118-128.
ISSN: 2456-4559
Available:www.ijbmm.com
17. Tripathy B, Tripathy P, Sahu GS, Dash SK, Pradhan B, Sahu P, Nayak NJ, Pradhan P, Sourav S, Mishra S. Evaluation of brinjal (*Solanum melongena* L.) landraces of Odisha for fruit yield and its components. J. Crop Weed. 2020;16(1):151-154.
 18. Singh K, Gupta K, Tyagi V, Rajkumar S. Plant genetic resources in India: Management and utilization. Vavilovskii Zhurnal Genetiki i Seleksii = Vavilov Journal of Genetics and Breeding. 2020;24(3):306-314.
 19. Gautam M, Kafle S, Regmi, Thapa G, Paudel S. Management of brinjal fruit and shoot borer (*Leucinodes orbonalis* Guenee) in Nepal. Acta Scientific Agr. 2019;3(9):188-195.
 20. Mathur A, Singh NP, Singh S. Management of brinjal shoot and fruit borer: Dilemma of adopting *Bt* brinjal over integrated pest management technology. International Conf. on Clean and Green Energy. IPCBEE. 2012;27:93-97.
 21. Breseghello F. Traditional and modern plant breeding methods with examples in rice (*Oryza sativa* L.). J. Agric. Fd. Chem. 2021;61:8277-8286.
 22. Agriculture contingency plan. Ministry of Agriculture & Farmers Welfare, Department of Agriculture & Farmers Welfare. Govt. of India. 2022. Available:<https://agricoop.nic.in/agriculture-contingency-plan-listing>
 23. NARP Zones: State agriculture plan and state infrastructure development plan (SAP & SIDP). Gujarat, Dept. Agri., Farmers Welfare and Co-operation Govt. of Gujarat, Gandhinagar. 2017-18 to 2019-20;271.

ANNEXURE

Site and agro-climatic descriptions of the various locations for the performance evaluation trials of farmer's brinjal variety 'Niranjan Bhata'

State	Districts covered	Agro-Climatic Region (ICAR-Agro Ecological Sub Region)	Agro Climatic Zone (NARP)
Manipur	Thoubal	Zone II: Eastern Himalayan Region (17.2 North-Eastern Hills (Purvachal), Warm Perhumid Eco-sub region)	Sub-Tropical Zone (NEH-4)
Chhattisgarh	Dhamtari	Zone-VII: Eastern plateau and hills (11.0 Chhattisgarh/Mahanadi Basin Agro-eco region)	Chhattisgarh plain zone
	Durg	Zone VII: Eastern plateau and hills region (11.0 Moderately To Gently Sloping Chattisgarh Mahanadi Basin, Hot Moist/Dry Sub-humid Transitional ESR With Deep Loamy To Clayey Red And Yellow Soils)	Chhattisgarh plain zone (MP-1)
Gujarat	Junagadh	Zone XIII: Gujarat Plains & Hills Region (5.1 Arid western Plains)	South Saurashtra Zone (GJ.7)
	Kheda	Zone XIII: Gujarat plains and hill region (5.2 Central Highlands (Malwa), Gujarat Plain And Kathiawar Peninsula, Semi-Arid Eco-Region)	Middle Gujarat Agro Climatic zone(GJ-3), North Gujarat Zone GJ-4
	Gandhinagar*	Zone XII: Gujarat Plains and Hills Region (4.2 Northern Plain (And Central Highlands) Including Aravallis, Hot Semi-Arid Eco- Region)	North Gujarat Zone (GJ-4)
	Bharuch	Zone XIII: Gujarat plains and hills region (5.2 Central (Malva) Highlands, Gujarat Plains and Kathiawar, Peninsula Ecoregion)	South Gujarat Zone (GJ-2)
	Narmada	Zone XIII: Gujarat Plains and hills region (5.2 Central highlands, Malwa, Gujarat Plain)	South Gujarat Zone (GJ-2)
Kerala	Kozhikode	Zone XII: West Coast Plains and Ghat Region (19.3 Konkan, Karnataka and Kerala Coastal plain, hot humid to perhumid eco-subregion)	Northern Zone (KE-1)
	Idukki	Zone XII: Western Plain and Ghat Region (19.2 Western Ghats And Coastal Plain, Hot-Humid Region)	High altitude zone (KE-4)
	Thrissur	Zone XII: West Coast Plains and Ghat Region (19.2 Konkan, Karnataka and Kerala Coastal plain, hot humid to perhumid eco-subregion (19.3) Central and south Sahyadris, hot moist, subhumid to humid eco-subregion)	Central Zone (KE 3)
Odisha	Kalahandi	Zone VII: Eastern plateau and hill region (12.1 Gujrat Hills, Dandakaranya and Eastern Ghats hot moist sub-humid eco sub-region)	Western undulating zone (OR-8)

State	Districts covered	Agro-Climatic Region (ICAR-Agro Ecological Sub Region)	Agro Climatic Zone (NARP)
Maharashtra	Nandurbar	Zone IX: Western Plateau & hills region (6.1 Deccan Plateau, hot semi-arid ecoregion)	Western Maharashtra Plain Zone (MH-6)

Source: Agriculture Contingency Plan. Ministry of Agriculture & Farmers Welfare, Department of Agriculture & Farmers Welfare. Govt. of India. 2022. [22,23]; * SAP & SIDP 2017-18 to 2019-20 Gujarat, Dept. Agri., Farmers Welfare and Co-operation Govt. of Gujarat, Gandhinagar

© 2023 Choudhary et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96567>